

Trawling Data in August 2021

| Group | Species Name | INA | | | | | | | | | | | | | | |
|-------|----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | INA-1 | | | INA-2 | | | INA-3 | | | INA-4 | | | INA-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| Fish | <i>Arnoglossus waiti</i> | | | | | | | | | | | | | | | |
| | <i>Anchoviella commersonii</i> | | | | | | | | | | | | | | | |
| | <i>Apogon kiensis</i> | 1 | 7.0 | | 6 | 29.0 | | 2 | 12.0 | | | | | | | |
| | <i>Apogon lineatus</i> | | | | | | | | | | | | | | | |
| | <i>Arothron hispidus</i> | | | | | | | | | | | | | | | |
| | <i>Arius chinensis</i> | | | | | | | | | | | | | 3 | 190.0 | |
| | <i>Butis koilomatodon</i> | | | | | | | | | | | | | | | |
| | <i>Callionymus curvicornis</i> | | | | | | | | | | | | | | | |
| | <i>Caranx kalla</i> | | | | | | | | | | | | | | | |
| | <i>Chrysochir aureus</i> | | | | | | | | | | | | | | | |
| | <i>Clupanodon thrissa</i> | | | | | | | | | | | | | | | |
| | <i>Coilia mystus</i> | | | | | | | | | | | | | | | |
| | <i>Collichthys lucidus</i> | | | | | | | | | | | | | | | |
| | <i>Cryptocentrus filifer</i> | | | | | | | | | | | | | | | |
| | <i>Cynoglossus arel</i> | | | | | | | | | | 1 | 10.0 | | | | |
| | <i>Cynoglossus puncticeps</i> | | | | | | | | | | 1 | 13.0 | | | | |
| | <i>Cynoglossus roulei</i> | | | | | | | | | | | | | | | |
| | <i>Dasyatis akaiei</i> | 4 | 760.0 | | 1 | 104.0 | | 1 | 60.0 | | | | | | | |
| | <i>Dasyatis zugei</i> | | | | 1 | 670.0 | | | | | | | | | | |
| | <i>Dendrophysa russelli</i> | | | | | | | 1 | 10.0 | | | | | | | |
| | <i>Drepane punctata</i> | 2 | 346.0 | | 3 | 990.0 | | | | | | | | 1 | 490.0 | |
| | <i>Epinephelus awoara</i> | | | | | | | | | | | | | | | |
| | <i>Gerres abbreviatus</i> | | | | | | | | | | | | | | | |
| | <i>Gerres lucidus</i> | | | | | | | | | | | | | | | |
| | <i>Grammoplites scaber</i> | 1 | 10.0 | | 2 | 21.0 | | 5 | 55.0 | | 1 | 13.0 | | | | |
| | <i>Gymnura japonica</i> | | | | | | | | | | | | | | | |
| | <i>Harpadon nehereus</i> | | | | | | | | | | | | | | | |
| | <i>Ilisha elongata</i> | 5 | 27.0 | | 5 | 20.0 | | | | | | | | | | |
| | <i>Inimicus japonicus</i> | 1 | 158.0 | | | | | | | | | | | | | |
| | <i>Johnius belangerii</i> | 4 | 70.0 | | 7 | 100.0 | | 5 | 180.0 | | 1 | 12.0 | | 4 | 100.0 | |
| | <i>Konosirus punctatus</i> | | | | | | | | | | | | | | | |
| | <i>Lagocephalus inermis</i> | 1 | 250.0 | | | | | 1 | 80.0 | | | | | | | |
| | <i>Lagocephalus sceleratus</i> | | | | | | | | | | | | | | | |
| | <i>Larimichthys polyactis</i> | | | | | | | | | | | | | | | |
| | <i>Leiognathus nuchalis</i> | | | | | | | | | | 1 | 20.0 | | 2 | 40.0 | |
| | <i>Lisa macrolepis</i> | | | | | | | | | | | | | | | |
| | <i>Lutianus sp.</i> | | | | | | | | | | | | | | | |
| | <i>Muraenesox cinereus</i> | | | | | | | | | | | | | | | |
| | <i>Nemipterus japonicus</i> | | | | | | | | | | 3 | 48.0 | | 1 | 12.0 | |
| | <i>Oxyurichthys tentacularis</i> | | | | | | | | | | | | | | | |
| | <i>Paramonacanthus polynema</i> | | | | | | | | | | | | | | | |
| | <i>Pennahia anea</i> | 2 | 30.0 | | 6 | 40.0 | | 6 | 50.0 | | 2 | 18.0 | | | | |
| | <i>Pisodonophis boro</i> | | | | | | | | | | | | | | | |
| | <i>Platycephalus indicus</i> | | | | | | | | | | | | | 11 | 120.0 | |
| | <i>Pseudorhombus cinnamomeus</i> | | | | | | | | | | | | | | | |
| | <i>Pseudorhombus oligodon</i> | | | | | | | 2 | 60.0 | | | | | | | |
| | <i>Saurida elongata</i> | | | | | | | | | | | | | | | |
| | <i>Scatophagus argus</i> | | | | | | | | | | | | | | | |
| | <i>Sebastiscus marmoratus</i> | 2 | 140.0 | | | | | 2 | 92.0 | | | | | 1 | 14.0 | |
| | <i>Siganus fuscescens</i> | | | | | | | | | | | | | 4 | 80.0 | |
| | <i>Sillago sihama</i> | 1 | 80.0 | | | | | 1 | 18.0 | | 2 | 130.0 | | | | |
| | <i>Solea ovata</i> | | | | | | | | | | | | | | | |
| | <i>Stephanolepis cirrhifer</i> | | | | | | | 2 | 250.0 | | | | | | | |

Trawling Data in August 2021

| Group | Species Name | INA | | | | | | | | | | | | | | |
|-----------------|-----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | INA-1 | | | INA-2 | | | INA-3 | | | INA-4 | | | INA-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Takifugu niphobles</i> | | | | | | | | | | | | | | | |
| | <i>Takifugu oblongus</i> | | | | 1 | 330.0 | | 2 | 340.0 | | | | | 2 | 380.0 | |
| | <i>Terapon jarbua</i> | | | | | | | | | | | | | | | |
| | <i>Terapon theraps</i> | 2 | 60.0 | | 4 | 73.0 | | 4 | 63.0 | | | | | | | |
| | <i>Thryssa kammalensis</i> | | | | | | | | | | | | | | | |
| | <i>Trachicephalus uranoscopus</i> | | | | | | | | | | | | | | | |
| | <i>Trypauchen vagina</i> | | | | | | | | | | | | | 1 | 13.0 | |
| Prawn or shrimp | <i>Alpheus brevirostratus</i> | | | | | | | | | | | | | | | |
| | <i>Alpheus distinguendus</i> | | | | | | | | | | | | | | | |
| | <i>Alpheus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Metapenaeopsis barbata</i> | 1 | 4.0 | | 2 | 5.0 | | 2 | 8.0 | | 1 | 5.0 | | | | |
| | <i>Metapenaeus affinis</i> | | | | | | | | | | | | | | | |
| | <i>Metapenaeus ensis</i> | | | | | | | | | | | | | | | |
| | <i>Metapenaeus joyneri</i> | | | | | | | | | | | | | | | |
| | <i>Parapeneopsis hardwickii</i> | | | | | | | | | | | | | | | |
| | <i>Parapeneopsis hungerfordii</i> | | | | | | | | | | | | | | | |
| | <i>Parapeneopsis tenella</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus japonicus</i> | | | | | | | | | | 1 | 22.0 | | | | |
| | <i>Penaeus merguensis</i> | | | | | | | | | | | | | 2 | 33.0 | |
| | <i>Penaeus penicillatus</i> | | | | | | | | | | | | | | | |
| | <i>Solenocera crassicornis</i> | | | | | | | | | | | | | | | |
| | c.f. <i>Thenus</i> sp. | 1 | 3.2 | | | | | | | | | | | | | |
| Crab | <i>Calappa philargius</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis affinis</i> | | | | 1 | 30.0 | | 3 | 45.0 | | 3 | 35.0 | | 5 | 80.0 | |
| | <i>Charybdis annulata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis cruciata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis hellerii</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis japonica</i> | 1 | 90.0 | | 1 | 18.0 | | 6 | 180.0 | | | | | 4 | 110.0 | |
| | <i>Charybdis truncata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis variegata</i> | | | | | | | | | | 1 | 12.0 | | | | |
| | <i>Charybdis spp.</i> | 6 | 47.2 | | 5 | 41.2 | | 7 | 51.4 | | | | | 3 | 31.0 | |
| | <i>Doclea spp.</i> | | | | | | | 2 | 53.0 | | 3 | 87.0 | | 7 | 181.0 | |
| | <i>Dorippe dorsipes</i> | 12 | 174.0 | | 24 | 283.0 | | 13 | 196.0 | | 11 | 161.0 | | 9 | 112.0 | |
| | <i>Dorippe polita</i> | | | | | | | | | | | | | | | |
| | <i>Eucrate crenata</i> | | | | | | | | | | | | | | | |
| | <i>Eucrate solaris</i> | | | | | | | | | | | | | | | |
| | <i>Galene bispinosa</i> | | | | | | | | | | | | | | | |
| | <i>Maorophthalmus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Parapanope euagora</i> | | | | | | | | | | | | | 1 | 8.1 | |
| | <i>Parthenope validus</i> | 4 | 39.7 | | 5 | 43.8 | | 6 | 51.8 | | 2 | 21.6 | | 16 | 159.0 | |
| | <i>Porcellanella picta</i> | | | | | | | | | | | | | | | |
| | <i>Portunus hastatoides</i> | | | | | | | | | | | | | | | |
| | <i>Portunus pelagicus</i> | | | | | | | | | | | | | 1 | 96.0 | |
| | <i>Portunus trituberculatus</i> | | | | | | | | | | | | | | | |
| | <i>Scylla paramamosain</i> | | | | | | | | | | | | | | | |
| | <i>Typhlocarcinus</i> sp. | | | | | | | | | | | | | | | |
| Mantis shrimp | <i>Dictyosquilla foveolata</i> | | | | | | | | | | | | | | | |
| | <i>Harpisquilla harpax</i> | | | | | | | | | | | | | | | |
| | <i>Oratosquilla interrupta</i> | | | | | | | | | | 1 | 6.0 | | 1 | 20.0 | |
| | <i>Oratosquilla kempii</i> | | | | | | | | | | | | | | | |
| | <i>Oratosquilla oratoria</i> | | | | | | | | | | 1 | 10.0 | | 3 | 30.0 | |
| Gastropod | <i>Calyptraea</i> sp. | | | | | | | | | | | | | | | |

Trawling Data in August 2021

| Group | Species Name | INA | | | | | | | | | | | | | | |
|------------|----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | INA-1 | | | INA-2 | | | INA-3 | | | INA-4 | | | INA-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Drupa musiva</i> | | | | | | | | | | | | | | | |
| | <i>Lophiotoma leucotropis</i> | | | | | | | | | | | | | | | |
| | <i>Murex trapa</i> | | | | | | | | | | | | | 8 | 66.4 | |
| | <i>Nassarius hepaticus</i> | | | | | | | | | | | | | | | |
| | <i>Nassarius succinctus</i> | | | | | | | | | | | | | | | |
| | <i>Siphopatella walshi</i> | 3 | 1.2 | | 10 | 9.4 | | | | | | | | 4 | 2.1 | |
| | <i>Toma sulcosa</i> | | | | | | | | | | | | | | | |
| | <i>Turritella terebra</i> | | | | | | | 3 | 27.0 | | | | | 4 | 31.0 | |
| | UNID Nudibranch | | | | | | | | | | | | | | | |
| Bivalve | <i>Anomia chinensis</i> | | | | | | | | | | | | | 1 | 1.8 | |
| | <i>Anomalocardia squamosa</i> | | | | | | | | | | | | | | | |
| | <i>Chlamys spp.</i> | | | | 3 | 26.0 | | 5 | 39.0 | | | | | 6 | 74.5 | |
| | <i>Paphia undulata</i> | | | | | | | | | | | | | | | |
| | <i>Perna viridis</i> | 2 | 12.6 | | | | | | | | | | | 2 | 33.4 | |
| | <i>Pteria sp.</i> | | | | 1 | 4.1 | | | | | | | | | | |
| | <i>Ruditapes philippinarum</i> | | | | | | | | | | | | | | | |
| | <i>Tegillarca nodifera</i> | | | | | | | | | | | | | | | |
| Cephalopod | <i>Lolingid squid</i> | 1 | 40.0 | | | | | 1 | 40.0 | | | | | | | |
| | <i>Sepiid cuttlefish</i> | 1 | 100.0 | | 1 | 140.0 | | 3 | 360.0 | | | | | | | |
| | <i>Octopus</i> | | | | | | | | | | | | | | | |
| Echinoderm | <i>Acaudina molpadioides</i> | | | | | | | | | | | | | | | |
| | <i>Comanthus sp.</i> | | | | | | | | | | | | | | | |
| | <i>Styela plicata</i> | | | | 7 | 34.4 | | | | | | | | | | |
| | <i>Temnopleurus toreumaticus</i> | 26 | 290.0 | | 18 | 202.0 | | 25 | 280.0 | | 30 | 310.0 | * | 40 | 420.0 | * |
| Cnidaria | <i>Carcinacis ichikawai</i> | | | | | | | | | | | | | | | |
| | <i>Cavernularia habereri</i> | | | | | | | 3 | 14.0 | | | | | | | |
| | <i>Ellisella laevis</i> | | | | | | | | | | | | | | | |
| | <i>Pteroeides chinense</i> | | | | | | | 4 | 86.0 | | | | | 2 | 44.0 | |
| | <i>Gorgonia spp.</i> | | | | | | | 1 | 36.0 | | | | | | | |
| | Jellyfish | 1 | 120.0 | | | | | | | | | | | | | |
| | <i>Virgularia gustaviana</i> | | | | | | | 2 | 7.2 | | | | | | | |
| Cirripedia | <i>c.f. Balanus spp.</i> | 30 | 18.0 | * | 40 | 21.0 | * | 30 | 18.0 | * | 30 | 18.0 | * | 40 | 21.0 | * |

Trawling Data in August 2021

| Group | Species Name | INB-1 | | | INB-2 | | | INB | | | INB-4 | | | INB-5 | | |
|-------|----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| Fish | <i>Arnoglossus waitei</i> | | | | | | | | | | | | | | | |
| | <i>Anchoviella commersonii</i> | | | | | | | | | | | | | | | |
| | <i>Apogon kiensis</i> | 4 | 22.0 | | 1 | 6.0 | | 3 | 22.0 | | 1 | 12.0 | | 1 | 10.0 | |
| | <i>Apogon lineatus</i> | | | | | | | | | | | | | | | |
| | <i>Arothron hispidus</i> | | | | | | | | | | | | | | | |
| | <i>Arius chinensis</i> | | | | | | | | | | | | | | | |
| | <i>Butis koilomatodon</i> | | | | 1 | 4.0 | | | | | | | | | | |
| | <i>Callionymus curvicornis</i> | | | | | | | | | | | | | | | |
| | <i>Caranx kalla</i> | | | | | | | | | | 1 | 20.0 | | | | |
| | <i>Chrysochir aureus</i> | | | | | | | | | | | | | | | |
| | <i>Clupanodon thrissa</i> | | | | | | | | | | | | | | | |
| | <i>Coilia mystus</i> | | | | | | | | | | | | | | | |
| | <i>Collichthys lucidus</i> | | | | | | | | | | | | | | | |
| | <i>Cryptocentrus filifer</i> | | | | | | | | | | | | | | | |
| | <i>Cynoglossus arel</i> | | | | 1 | 28.0 | | | | | | | | | | |
| | <i>Cynoglossus puncticeps</i> | 1 | 22.0 | | | | | | | | 1 | 14.0 | | | | |
| | <i>Cynoglossus roulei</i> | | | | | | | | | | | | | | | |
| | <i>Dasvatis akajei</i> | 1 | 160.0 | | | | | | | | | | | | | |
| | <i>Dasvatis zugei</i> | | | | | | | | | | | | | | | |
| | <i>Dendrophysa russelli</i> | | | | | | | | | | | | | | | |
| | <i>Drepane punctata</i> | 1 | 450.0 | | | | | | | | | | | | | |
| | <i>Epinephelus awoara</i> | | | | | | | | | | | | | | | |
| | <i>Gerres abbreviatus</i> | | | | | | | | | | | | | | | |
| | <i>Gerres lucidus</i> | | | | | | | | | | | | | | | |
| | <i>Grammolites scaber</i> | 3 | 50.0 | | 1 | 10.0 | | 4 | 50.0 | | | | | | | |
| | <i>Gymnura japonica</i> | | | | | | | | | | | | | | | |
| | <i>Harpadon nehereus</i> | | | | | | | | | | | | | | | |
| | <i>Ilisha elongata</i> | | | | | | | | | | | | | | | |
| | <i>Inimicus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Johnius belangerii</i> | 11 | 130.0 | | 16 | 270.0 | | 15 | 270.0 | | 6 | 90.0 | | 7 | 103.0 | |
| | <i>Konosirus punctatus</i> | | | | | | | | | | | | | | | |
| | <i>Lagocephalus inermis</i> | | | | | | | | | | | | | | | |
| | <i>Lagocephalus sceleratus</i> | | | | | | | | | | | | | | | |
| | <i>Larimichthys polyactis</i> | | | | | | | | | | | | | | | |
| | <i>Leiognathus nuchalis</i> | | | | | | | | | | | | | | | |
| | <i>Lisa macrolepis</i> | | | | | | | | | | | | | | | |
| | <i>Lutjanus sp.</i> | | | | | | | | | | | | | | | |
| | <i>Muraenesox cinereus</i> | | | | | | | | | | | | | | | |
| | <i>Nemipterus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Oxyurichthys tentacularis</i> | | | | | | | | | | | | | | | |
| | <i>Paramonacanthus polynema</i> | | | | | | | | | | | | | | | |
| | <i>Pennahia anea</i> | | | | 1 | 8.0 | | | | | 1 | 10.0 | | 1 | 10.0 | |
| | <i>Pisodonophis boro</i> | | | | | | | | | | | | | | | |
| | <i>Platycephalus indicus</i> | | | | | | | | | | | | | | | |
| | <i>Pseudorhombus cinnamomeus</i> | | | | | | | | | | | | | | | |
| | <i>Pseudorhombus oligodon</i> | | | | | | | | | | | | | | | |
| | <i>Saurida elongata</i> | | | | | | | | | | | | | | | |
| | <i>Scatophagus argus</i> | | | | | | | 1 | 150.0 | | | | | | | |
| | <i>Sebastes marmoratus</i> | 1 | 18.0 | | 2 | 22.0 | | 2 | 20.0 | | | | | | | |
| | <i>Siganus fuscescens</i> | | | | | | | | | | | | | | | |
| | <i>Sillago sihama</i> | | | | | | | | | | | | | | | |
| | <i>Solea ovata</i> | | | | | | | | | | | | | | | |
| | <i>Stephanolepis cirrhifer</i> | | | | | | | | | | | | | | | |

Trawling Data in August 2021

| Group | Species Name | INB-1 | | | INB-2 | | | INB | | | INB-4 | | | INB-5 | | |
|-----------------|-----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Takifugu niphobles</i> | | | | | | | | | | | | | | | |
| | <i>Takifugu oblongus</i> | | | | | | | | | | | | | | | |
| | <i>Terapon jarbua</i> | | | | | | | | | | | | | | | |
| | <i>Terapon theraps</i> | | | | 1 | 30.0 | | | | | | | | | | |
| | <i>Thryssa kammalensis</i> | | | | | | | | | | | | | | | |
| | <i>Trachicephalus uranoscopus</i> | | | | | | | | | | | | | | | |
| | <i>Trypauchen vagina</i> | | | | | | | | | | | | | | | |
| Prawn or shrimp | <i>Alpheus brevicristatus</i> | | | | 1 | 3.0 | | | | | | | | | | |
| | <i>Alpheus distinguendus</i> | | | | | | | | | | | | | | | |
| | <i>Alpheus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Metapenaeopsis barbata</i> | 2 | 9.0 | | | | | 1 | 4.0 | | | | | | | |
| | <i>Metapenaeus affinis</i> | | | | | | | 1 | 10.0 | | 1 | 10.0 | | 3 | 10.0 | |
| | <i>Metapenaeus ensis</i> | | | | | | | | | | | | | | | |
| | <i>Metapenaeus jayneri</i> | | | | | | | | | | | | | | | |
| | <i>Parapeneopsis hardwickii</i> | | | | | | | | | | | | | | | |
| | <i>Parapeneopsis hungerfordii</i> | | | | | | | | | | | | | | | |
| | <i>Parapeneopsis tenella</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus merguensis</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus penicillatus</i> | | | | | | | | | | | | | | | |
| | <i>Solenocera crassicornis</i> | | | | | | | | | | 1 | 4.0 | | 1 | 7.0 | |
| | <i>c.f. Thenus sp.</i> | | | | | | | | | | | | | | | |
| Crab | <i>Calappa philargius</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis affinis</i> | 50 | 880.0 | | 17 | 303.0 | | 43 | 880.0 | | 15 | 190.0 | | 17 | 270.0 | |
| | <i>Charybdis annulata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis cruciata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis hellerii</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis japonica</i> | 5 | 190.0 | | 8 | 360.0 | | 5 | 250.0 | | | | | | | |
| | <i>Charybdis truncata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis variegata</i> | | | | 2 | 50.0 | | 5 | 90.0 | | 6 | 120.0 | | | | |
| | <i>Charybdis spp.</i> | | | | 5 | 42.0 | | 8 | 64.0 | | 18 | 170.0 | | 11 | 107.0 | |
| | <i>Doclea spp.</i> | | | | | | | | | | | | | 20 | 370.0 | * |
| | <i>Dorippe dorsipes</i> | 40 | 420.0 | * | 2 | 16.0 | | 30 | 390.0 | * | 2 | 21.0 | | | | |
| | <i>Dorippe polita</i> | | | | | | | | | | | | | | | |
| | <i>Eucrete crenata</i> | | | | | | | 20 | 230.0 | | | | | | | |
| | <i>Eucrete solaris</i> | | | | | | | | | | | | | | | |
| | <i>Galene hispinosa</i> | | | | | | | | | | | | | | | |
| | <i>Maorophthalmus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Parapanope euagora</i> | | | | | | | 1 | 8.4 | | | | | | | |
| | <i>Parthenope validus</i> | 30 | 330.0 | * | 7 | 23.0 | | 30 | 340.0 | * | 6 | 76.0 | | | | |
| | <i>Porcellanella picta</i> | | | | | | | | | | | | | | | |
| | <i>Portunus hastatooides</i> | | | | | | | | | | | | | | | |
| | <i>Portunus pelagicus</i> | 1 | 80.0 | | | | | 2 | 260.0 | | 2 | 210.0 | | | | |
| | <i>Portunus trituberculatus</i> | | | | | | | | | | | | | | | |
| | <i>Scylla paramamosain</i> | | | | | | | | | | | | | | | |
| | <i>Typhlocarcinus sp.</i> | | | | | | | | | | | | | | | |
| Mantis shrimp | <i>Dictyosquilla foveolata</i> | | | | | | | | | | | | | | | |
| | <i>Harpisquilla harpax</i> | 1 | 50.0 | | 1 | 45.0 | | 1 | 42.0 | | | | | | | |
| | <i>Oratosquilla interrupta</i> | 1 | 3.0 | | | | | 1 | 20.0 | | 2 | 20.0 | | 1 | 30.0 | |
| | <i>Oratosquilla kempii</i> | | | | | | | | | | | | | | | |
| | <i>Oratosquilla oratoria</i> | 1 | 13.0 | | | | | | | | | | | | | |
| Gastropod | <i>Calyptreaea sp.</i> | | | | | | | 1 | 0.3 | | | | | | | |

Trawling Data in August 2021

| Group | Species Name | INB-1 | | | INB-2 | | | INB | | | INB-4 | | | INB-5 | | |
|------------|----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Drupa musiva</i> | 20 | 130.0 | * | | | | | | | | | | | | |
| | <i>Lophiotoma leucotropis</i> | | | | 2 | 10.2 | | | | | | | | | | |
| | <i>Murex trapa</i> | | | | | | | 4 | 21.0 | | | | | | | |
| | <i>Nassarius hepaticus</i> | | | | | | | 1 | 6.2 | | | | | | | |
| | <i>Nassarius succinctus</i> | | | | | | | | | | | | | | | |
| | <i>Siphonotella walshi</i> | | | | | | | | | | | | | | | |
| | <i>Tonna sulcosa</i> | | | | | | | | | | | | | | | |
| | <i>Turritella terebra</i> | 5 | 42.0 | | 7 | 62.0 | | 3 | 23.0 | | 7 | 63.0 | | 5 | 36.0 | |
| | <i>UNID Nudibranch</i> | | | | | | | | | | | | | | | |
| Bivalve | <i>Anomia chinensis</i> | | | | | | | | | | | | | | | |
| | <i>Anomalocardia squamosa</i> | | | | | | | | | | | | | | | |
| | <i>Chlamys spp.</i> | 40 | 210.0 | * | 20 | 106.0 | * | 40 | 250.0 | * | | | | 5 | 43.0 | |
| | <i>Paphia undulata</i> | | | | | | | | | | | | | | | |
| | <i>Perna viridis</i> | 1 | 4.3 | | | | | | | | | | | | | |
| | <i>Pteria sp.</i> | | | | | | | | | | | | | | | |
| | <i>Ruditapes philippinarum</i> | 20 | 82.0 | * | 20 | 82.0 | * | | | | 4 | 21.0 | | 7 | 52.0 | |
| | <i>Tegillarca nodifera</i> | | | | | | | | | | | | | | | |
| Cephalopod | <i>Lolingid squid</i> | | | | | | | | | | | | | | | |
| | <i>Sepiid cuttlefish</i> | | | | 1 | 140.0 | | | | | | | | | | |
| | <i>Octopus</i> | | | | 1 | 130.0 | | | | | | | | | | |
| Echinoderm | <i>Acaudina molpadioides</i> | | | | | | | | | | | | | | | |
| | <i>Comanthus sp.</i> | | | | | | | | | | 1 | 6.2 | | | | |
| | <i>Styela plicata</i> | | | | | | | | | | | | | | | |
| | <i>Temnopleurus toreumaticus</i> | 90 | 860.0 | * | 110 | 1070.0 | * | 80 | 810.0 | * | 110 | 1090.0 | * | 60 | 630.0 | * |
| Cnidaria | <i>Carcinactis ichikawai</i> | | | | | | | | | | | | | | | |
| | <i>Cavernularia habereri</i> | 5 | 22.0 | | 12 | 58.0 | | 4 | 18.0 | | | | | 7 | 59.0 | |
| | <i>Ellisella laevis</i> | | | | 2 | 18.0 | | 1 | 30.0 | | 1 | 6.0 | | | | |
| | <i>Pteroeides chinense</i> | 6 | 150.0 | | 30 | 830.0 | * | 5 | 130.0 | | 2 | 52.0 | | | | |
| | <i>Gorgonia spp.</i> | | | | | | | | | | | | | | | |
| | Jellyfish | 2 | 230.0 | | 3 | 1460.0 | | | | | 13 | 2240.0 | | 4 | 480.0 | |
| | <i>Virgularia gustaviana</i> | | | | 1 | 9.2 | | 3 | 20.4 | | | | | | | |
| Cirripedia | <i>c.f. Balanus spp.</i> | 60 | 52.0 | * | 90 | 54.0 | * | 80 | 46.0 | * | 90 | 48.0 | * | 70 | 41.0 | * |

Trawling Data in August 2021

| Group | Species Name | TNA-1 | | | TNA-2 | | | TNA-3 | | | TNA-4 | | | TNA-5 | | |
|-------|----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| Fish | <i>Arnoglossus waitei</i> | | | | | | | | | | | | | | | |
| | <i>Anchoviella commersonii</i> | | | | | | | | | | | | | | | |
| | <i>Apogon kiensis</i> | 3 | 23.0 | | | | | | | | | | | 2 | 11.0 | |
| | <i>Apogon lineatus</i> | | | | | | | | | | | | | | | |
| | <i>Arothron hispidus</i> | | | | | | | | | | | | | | | |
| | <i>Arius chinensis</i> | | | | | | | | | | | | | | | |
| | <i>Butis koilomatodon</i> | | | | | | | | | | | | | | | |
| | <i>Callionymus curvicornis</i> | | | | | | | | | | | | | | | |
| | <i>Caranx kalla</i> | | | | | | | | | | | | | | | |
| | <i>Chrysochir aureus</i> | | | | | | | | | | | | | | | |
| | <i>Clupanodon thrissa</i> | | | | | | | | | | | | | | | |
| | <i>Coilia mystus</i> | | | | 1 | 20.0 | | | | | 2 | 54.0 | | | | |
| | <i>Collichthys lucidus</i> | 1 | 20.0 | | | | | | | | 3 | 80.0 | | | | |
| | <i>Cryptocentrus filifer</i> | | | | | | | | | | | | | | | |
| | <i>Cynoglossus arel</i> | 10 | 360.0 | | 2 | 12.0 | | 3 | 20.0 | | 8 | 60.0 | | 9 | 130.0 | |
| | <i>Cynoglossus puncticeps</i> | | | | | | | | | | | | | | | |
| | <i>Cynoglossus roulei</i> | | | | | | | | | | | | | | | |
| | <i>Dasvatis akajei</i> | | | | | | | | | | | | | | | |
| | <i>Dasvatis zugei</i> | 1 | 490.0 | | 3 | 140.0 | | | | | 3 | 648.0 | | 3 | 130.0 | |
| | <i>Dendrophysa russelli</i> | 3 | 70.0 | | 7 | 90.0 | | 1 | 10.0 | | 5 | 100.0 | | | | |
| | <i>Drepane punctata</i> | | | | | | | | | | | | | | | |
| | <i>Epinephelus awoara</i> | | | | | | | | | | | | | | | |
| | <i>Gerres abbreviatus</i> | | | | | | | | | | | | | | | |
| | <i>Gerres lucidus</i> | | | | | | | | | | | | | | | |
| | <i>Grammolites scaber</i> | | | | | | | | | | | | | | | |
| | <i>Gymnura japonica</i> | | | | | | | | | | | | | 1 | 202.0 | |
| | <i>Harpadon nehereus</i> | | | | | | | | | | 1 | 170.0 | | | | |
| | <i>Ilisha elongata</i> | | | | | | | | | | | | | | | |
| | <i>Inimicus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Johnius belangerii</i> | 6 | 50.0 | | 9 | 120.0 | | 3 | 20.0 | | 5 | 70.0 | | 9 | 220.0 | |
| | <i>Konosirus punctatus</i> | | | | | | | | | | | | | | | |
| | <i>Lagocephalus inermis</i> | | | | | | | | | | | | | | | |
| | <i>Lagocephalus scleratus</i> | | | | | | | | | | | | | | | |
| | <i>Larimichthys polyactis</i> | | | | | | | | | | | | | | | |
| | <i>Leiognathus nuchalis</i> | | | | | | | | | | | | | | | |
| | <i>Lisa macrolepis</i> | | | | | | | | | | | | | | | |
| | <i>Lutjanus sp.</i> | | | | | | | | | | | | | | | |
| | <i>Muraenesox cinereus</i> | | | | | | | | | | | | | | | |
| | <i>Nemipterus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Oxyurichthys tentacularis</i> | | | | | | | | | | | | | | | |
| | <i>Paramonacanthus polynema</i> | | | | | | | 3 | 22.0 | | | | | | | |
| | <i>Pennahia anea</i> | 30 | 330.0 | | 40 | 440.0 | | 15 | 150.0 | | 8 | 80.0 | | 26 | 320.0 | |
| | <i>Pisodonophis boro</i> | | | | | | | | | | | | | | | |
| | <i>Platycephalus indicus</i> | | | | | | | | | | | | | | | |
| | <i>Pseudorhombus cinnamomeus</i> | | | | | | | | | | | | | | | |
| | <i>Pseudorhombus oligodon</i> | | | | | | | 1 | 6.0 | | | | | | | |
| | <i>Saurida elongata</i> | | | | | | | | | | | | | | | |
| | <i>Scatophagus argus</i> | | | | | | | | | | | | | | | |
| | <i>Sebastiscus marmoratus</i> | | | | | | | | | | | | | | | |
| | <i>Siganus fuscescens</i> | | | | | | | | | | | | | | | |
| | <i>Sillago sihama</i> | | | | | | | | | | | | | | | |
| | <i>Solea ovata</i> | 1 | 5.0 | | | | | | | | | | | | | |
| | <i>Stephanolepis cirrifer</i> | | | | | | | | | | | | | | | |

Trawling Data in August 2021

| Group | Species Name | TNA | | | | | | | | | | | | | | |
|-----------------|-----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | TNA-1 | | | TNA-2 | | | TNA-3 | | | TNA-4 | | | TNA-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Takifugu niphobles</i> | | | | | | | | | | | | | | | |
| | <i>Takifugu oblongus</i> | | | | | | | | | | | | | | | |
| | <i>Terapon jarbua</i> | | | | | | | | | | | | | | | |
| | <i>Terapon theraps</i> | | | | | | | | | | | | | | | |
| | <i>Thryssa kammalensis</i> | | | | | | | | | | | | | | | |
| | <i>Trachicephalus uranoscopus</i> | | | | 2 | 40.0 | | | | | | | | | | |
| | <i>Trypauchen vagina</i> | 3 | 20.0 | | 6 | 40.0 | | 5 | 60.0 | | 2 | 20.0 | | 1 | 15.0 | |
| Prawn or shrimp | <i>Alpheus brevirostratus</i> | | | | | | | | | | | | | | | |
| | <i>Alpheus distinguendus</i> | | | | 2 | 10.0 | | | | | | | | | | |
| | <i>Alpheus japonicus</i> | 1 | 15.0 | | | | | 1 | 6.0 | | | | | | | |
| | <i>Metapenaeopsis barbata</i> | 2 | 8.0 | | 1 | 4.0 | | | | | | | | | | |
| | <i>Metapenaeus affinis</i> | 102 | 1070.0 | | 60 | 370.0 | | 31 | 230.0 | | 62 | 620.0 | | 50 | 480.0 | |
| | <i>Metapenaeus ensis</i> | | | | | | | | | | 1 | 5.0 | | | | |
| | <i>Metapenaeus jowneri</i> | 1 | 10.0 | | | | | | | | 1 | 8.0 | | 1 | 10.0 | |
| | <i>Parapeneopsis hardwickii</i> | | | | 2 | 23.0 | | | | | 3 | 35.0 | | 7 | 80.0 | |
| | <i>Parapeneopsis hungerfordii</i> | 10 | 70.0 | | 10 | 60.0 | | 4 | 22.0 | | 20 | 120.0 | | 21 | 110.0 | |
| | <i>Parapeneopsis tenella</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus japonicus</i> | | | | 1 | 25.0 | | | | | | | | | | |
| | <i>Penaeus merguensis</i> | 7 | 102.0 | | 2 | 62.0 | | | | | 2 | 30.0 | | 9 | 110.0 | |
| | <i>Penaeus penicillatus</i> | | | | | | | | | | | | | | | |
| | <i>Solenocera crassicornis</i> | 2 | 7.0 | | 2 | 10.0 | | | | | 1 | 6.0 | | | | |
| | <i>c.f. Thenus sp.</i> | | | | | | | | | | | | | | | |
| Crab | <i>Calappa philargius</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis affinis</i> | 46 | 550.0 | | 40 | 570.0 | | 18 | 230.0 | | 60 | 770.0 | | 28 | 320.0 | |
| | <i>Charybdis annulata</i> | | | | | | | | | | 2 | 23.0 | | | | |
| | <i>Charybdis cruciata</i> | | | | 1 | 7.0 | | 1 | 28.0 | | 1 | 110.0 | | | | |
| | <i>Charybdis hellerii</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis japonica</i> | | | | | | | | | | 1 | 40.0 | | | | |
| | <i>Charybdis truncata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis variegata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis spp.</i> | 30 | 320.0 | * | 30 | 320.0 | * | 40 | 410.0 | * | 30 | 320.0 | * | 40 | 420.0 | * |
| | <i>Doclea spp.</i> | | | | | | | | | | | | | | | |
| | <i>Dorippe dorsipes</i> | | | | | | | | | | | | | | | |
| | <i>Dorippe polita</i> | | | | 3 | 31.6 | | | | | 6 | 53.0 | | 4 | 37.0 | |
| | <i>Eucreate crenata</i> | | | | | | | | | | | | | 1 | 7.6 | |
| | <i>Eucreate solaris</i> | | | | | | | | | | | | | | | |
| | <i>Galene hispinosa</i> | 11 | 390.0 | | | | | 20 | 890.0 | * | 30 | 1340.0 | * | 2 | 53.0 | |
| | <i>Maorophthalmus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Parapanope euagora</i> | | | | | | | | | | | | | | | |
| | <i>Parthenope validus</i> | 6 | 66.0 | | | | | | | | | | | | | |
| | <i>Porcellanella picta</i> | | | | | | | | | | | | | | | |
| | <i>Portunus hastatooides</i> | 2 | 8.0 | | | | | | | | | | | | | |
| | <i>Portunus pelagicus</i> | | | | | | | | | | | | | 1 | 190.0 | |
| | <i>Portunus trituberculatus</i> | | | | | | | | | | | | | | | |
| | <i>Scylla paramamosain</i> | | | | | | | | | | | | | | | |
| | <i>Typhlocarcinus sp.</i> | | | | | | | | | | | | | | | |
| Mantis shrimp | <i>Dictyosquilla foveolata</i> | 2 | 40.0 | | | | | 2 | 30.0 | | 3 | 60.0 | | | | |
| | <i>Harpiosquilla harpax</i> | | | | 2 | 230.0 | | | | | 2 | 190.0 | | | | |
| | <i>Oratosquilla interrupta</i> | 51 | 540.0 | | 28 | 430.0 | | 10 | 160.0 | | 34 | 530.0 | | 24 | 236.0 | |
| | <i>Oratosquilla kempii</i> | | | | | | | | | | | | | | | |
| | <i>Oratosquilla oratoria</i> | 1 | 30.0 | | 2 | 30.0 | | 1 | 20.0 | | 5 | 80.0 | | 2 | 20.0 | |
| Gastropod | <i>Calyptreaea sp.</i> | | | | | | | | | | | | | | | |

Trawling Data in August 2021

| Group | Species Name | TNA | | | | | | | | | | | | | | |
|------------|----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | TNA-1 | | | TNA-2 | | | TNA-3 | | | TNA-4 | | | TNA-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Drupa musiva</i> | | | | | | | | | | | | | | | |
| | <i>Lophiotoma leucotropis</i> | | | | | | | | | | | | | | | |
| | <i>Murex trapa</i> | | | | | | | | | | 3 | 21.0 | | 6 | 48.0 | |
| | <i>Nassarius hepaticus</i> | | | | | | | | | | | | | | | |
| | <i>Nassarius succinctus</i> | | | | | | | | | | | | | | | |
| | <i>Siphonotus walshi</i> | | | | | | | | | | | | | | | |
| | <i>Tonna sulcosa</i> | | | | | | | | | | | | | | | |
| | <i>Turritella terebra</i> | 30 | 280.0 | * | 40 | 390.0 | * | 40 | 390.0 | * | 40 | 390.0 | * | 50 | 480.0 | * |
| | <i>UNID Nudibranch</i> | | | | | | | | | | | | | | | |
| Bivalve | <i>Anomia chinensis</i> | | | | | | | | | | | | | | | |
| | <i>Anomalocardia squamosa</i> | | | | | | | | | | | | | | | |
| | <i>Chlamys spp.</i> | | | | | | | | | | | | | | | |
| | <i>Paphia undulata</i> | | | | | | | | | | | | | | | |
| | <i>Perna viridis</i> | | | | | | | | | | | | | | | |
| | <i>Pteria sp.</i> | | | | | | | | | | | | | | | |
| | <i>Ruditapes philippinarum</i> | | | | | | | | | | | | | | | |
| | <i>Tegillarca nodifera</i> | | | | | | | | | | | | | | | |
| Cephalopod | <i>Lolingid squid</i> | | | | | | | | | | | | | | | |
| | <i>Sepiid cuttlefish</i> | | | | | | | | | | | | | | | |
| | <i>Octopus</i> | | | | | | | | | | | | | | | |
| Echinoderm | <i>Acaudina molpadioides</i> | | | | | | | | | | | | | | | |
| | <i>Comanthus sp.</i> | | | | | | | | | | | | | | | |
| | <i>Styela plicata</i> | | | | | | | | | | | | | | | |
| | <i>Temnopleurus toreumaticus</i> | 520 | 15120.0 | * | 1230 | 11890.0 | * | 20 | 210.0 | * | 20 | 210.0 | * | 17 | 160.0 | |
| Cnidaria | <i>Carcinactis ichikawai</i> | | | | | | | | | | | | | | | |
| | <i>Cavernularia habereri</i> | 11 | 96.0 | | 24 | 210.0 | | 5 | 26.0 | | 4 | 22.0 | | 6 | 25.0 | |
| | <i>Ellisella laevis</i> | | | | | | | | | | | | | | | |
| | <i>Pteroeides chinense</i> | 18 | 370.0 | | 22 | 460.0 | | 16 | 340.0 | | 20 | 480.0 | | 17 | 370.0 | |
| | <i>Gorgonia spp.</i> | | | | | | | | | | | | | | | |
| | Jellyfish | 3 | 1210.0 | | 4 | 1660.0 | | 7 | 2160.0 | | 2 | 726.0 | | 3 | 940.0 | |
| | <i>Virgularia gustaviana</i> | | | | | | | 1 | 12.0 | | | | | | | |
| Cirripedia | <i>c.f. Balanus spp.</i> | | | | | | | | | | | | | | | |

Trawling Data in August 2021

| Group | Species Name | TNB-1 | | | TNB-2 | | | TNB | | | TNB-4 | | | TNB-5 | | |
|-------|----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| Fish | <i>Arnoglossus waitei</i> | | | | | | | | | | | | | | | |
| | <i>Anchoviella commersonii</i> | | | | | | | | | | | | | | | |
| | <i>Apogon kiensis</i> | 14 | 82.0 | | 10 | 52.0 | | 9 | 53.0 | | 4 | 22.0 | | 3 | 18.0 | |
| | <i>Apogon lineatus</i> | | | | | | | | | | | | | | | |
| | <i>Arothron hispidus</i> | | | | | | | | | | | | | | | |
| | <i>Arius chinensis</i> | | | | 1 | 28.0 | | | | | | | | | | |
| | <i>Butis koilomatodon</i> | | | | | | | | | | | | | | | |
| | <i>Callionymus curvicornis</i> | | | | | | | | | | | | | | | |
| | <i>Caranx kalla</i> | | | | | | | | | | | | | | | |
| | <i>Chrysochir aureus</i> | | | | 1 | 28.0 | | | | | | | | | | |
| | <i>Clupanodon thrissa</i> | | | | | | | | | | | | | | | |
| | <i>Coilia mystus</i> | | | | | | | | | | | | | | | |
| | <i>Collichthys lucidus</i> | | | | | | | | | | | | | | | |
| | <i>Cryptocentrus filifer</i> | 3 | 12.0 | | 1 | 11.0 | | | | | | | | 1 | 8.0 | |
| | <i>Cynoglossus arel</i> | 2 | 11.0 | | 10 | 240.0 | | 6 | 140.0 | | 7 | 80.0 | | 11 | 565.0 | |
| | <i>Cynoglossus puncticeps</i> | | | | | | | | | | | | | | | |
| | <i>Cynoglossus roulei</i> | | | | | | | | | | | | | | | |
| | <i>Dasvatis akajei</i> | | | | | | | | | | | | | | | |
| | <i>Dasvatis zugei</i> | 11 | 1890.0 | | 6 | 2060.0 | | 1 | 20.0 | | 11 | 1340.0 | | 8 | 870.0 | |
| | <i>Dendrophysa russelli</i> | 23 | 308.0 | | 15 | 120.0 | | 8 | 103.0 | | 10 | 180.0 | | 8 | 138.0 | |
| | <i>Drepane punctata</i> | | | | | | | | | | | | | | | |
| | <i>Epinephelus awoara</i> | 4 | 21.0 | | 5 | 33.0 | | 2 | 22.0 | | 4 | 21.0 | | | | |
| | <i>Gerres abbreviatus</i> | | | | | | | | | | | | | | | |
| | <i>Gerres lucidus</i> | | | | | | | | | | | | | | | |
| | <i>Grammolites scaber</i> | | | | 1 | 80.0 | | 3 | 34.0 | | | | | 3 | 44.0 | |
| | <i>Gymnura japonica</i> | | | | | | | | | | | | | | | |
| | <i>Harpadon nehereus</i> | | | | | | | | | | | | | | | |
| | <i>Ilisha elongata</i> | | | | | | | | | | | | | | | |
| | <i>Inimicus japonicus</i> | 1 | 190.0 | | | | | | | | | | | | | |
| | <i>Johnius belangerii</i> | 4 | 20.0 | | 2 | 10.0 | | 1 | 6.0 | | | | | 2 | 50.0 | |
| | <i>Konosirus punctatus</i> | | | | | | | | | | | | | | | |
| | <i>Lagocephalus inermis</i> | | | | | | | | | | | | | | | |
| | <i>Lagocephalus sceleratus</i> | | | | | | | | | | | | | | | |
| | <i>Larimichthys polyactis</i> | | | | | | | | | | | | | | | |
| | <i>Leiognathus nuchalis</i> | 2 | 30.0 | | 1 | 12.0 | | | | | | | | | | |
| | <i>Lisa macrolepis</i> | | | | | | | | | | | | | | | |
| | <i>Lutianus sp.</i> | | | | | | | | | | | | | | | |
| | <i>Muraenesox cinereus</i> | | | | | | | | | | 1 | 14.0 | | | | |
| | <i>Nemipterus japonicus</i> | 6 | 90.0 | | 8 | 130.0 | | 3 | 44.0 | | 1 | 13.0 | | 1 | 22.0 | |
| | <i>Oxyurichthys tentacularis</i> | | | | | | | | | | | | | | | |
| | <i>Paramonacanthus polynema</i> | | | | 1 | 15.0 | | | | | 1 | 15.0 | | | | |
| | <i>Pennahia anea</i> | 3 | 20.0 | | | | | 1 | 24.0 | | | | | 2 | 20.0 | |
| | <i>Pisodonophis boro</i> | | | | | | | | | | | | | | | |
| | <i>Platycephalus indicus</i> | 6 | 205.0 | | 1 | 30.0 | | 2 | 22.0 | | 1 | 240.0 | | 1 | 220.0 | |
| | <i>Pseudorhombus cinnamomeus</i> | | | | | | | | | | | | | | | |
| | <i>Pseudorhombus oligodon</i> | 1 | 40.0 | | | | | | | | | | | | | |
| | <i>Saurida elongata</i> | | | | | | | | | | | | | | | |
| | <i>Scatophagus argus</i> | | | | | | | | | | | | | | | |
| | <i>Sebastiscus marmoratus</i> | | | | | | | | | | | | | | | |
| | <i>Siganus fuscescens</i> | | | | | | | | | | | | | | | |
| | <i>Sillago sihama</i> | | | | | | | | | | | | | | | |
| | <i>Solea ovata</i> | 1 | 10.0 | | | | | 1 | 20.0 | | | | | | | |
| | <i>Stephanolepis cirrhifer</i> | 1 | 28.0 | | | | | | | | | | | | | |

Trawling Data in August 2021

| Group | Species Name | TNB-1 | | | TNB-2 | | | TNB | | | TNB-4 | | | TNB-5 | | |
|-----------------|-----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Takifugu niphobles</i> | | | | | | | | | | | | | | | |
| | <i>Takifugu oblongus</i> | | | | | | | | | | | | | | | |
| | <i>Terapon jarbua</i> | | | | | | | | | | | | | | | |
| | <i>Terapon theraps</i> | | | | | | | | | | | | | | | |
| | <i>Thryssa kammalensis</i> | | | | | | | | | | | | | | | |
| | <i>Trachicephalus uranoscopus</i> | 5 | 84.0 | | 2 | 32.0 | | | | | | | | | | |
| | <i>Trypauchen vagina</i> | 12 | 102.0 | | 13 | 130.0 | | 12 | 160.0 | | 11 | 130.0 | | 16 | 150.0 | |
| Prawn or shrimp | <i>Alpheus brevicristatus</i> | 2 | 8.0 | | 1 | 18.0 | | | | | | | | | | |
| | <i>Alpheus distinguendus</i> | | | | | | | | | | | | | | | |
| | <i>Alpheus japonicus</i> | | | | | | | 1 | 12.0 | | | | | | | |
| | <i>Metapenaeopsis barbata</i> | | | | | | | 1 | 4.0 | | | | | | | |
| | <i>Metapenaeus affinis</i> | | | | 19 | 130.0 | | 14 | 80.0 | | 4 | 30.0 | | 6 | 30.0 | |
| | <i>Metapenaeus ensis</i> | | | | | | | | | | | | | | | |
| | <i>Metapenaeus joyneri</i> | | | | 2 | 12.0 | | | | | 3 | 15.0 | | 4 | 22.0 | |
| | <i>Parapeneopsis hardwickii</i> | | | | | | | | | | | | | | | |
| | <i>Parapeneopsis hungerfordii</i> | 1 | 5.0 | | 4 | 33.0 | | 2 | 15.0 | | 5 | 36.0 | | 8 | 63.0 | |
| | <i>Parapeneopsis tenella</i> | 1 | 12.0 | | 4 | 42.0 | | | | | 1 | 14.0 | | | | |
| | <i>Penaeus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus merguensis</i> | 28 | 210.0 | | 12 | 150.0 | | 5 | 60.0 | | 2 | 25.0 | | 4 | 44.0 | |
| | <i>Penaeus penicillatus</i> | | | | 3 | 21.0 | | 10 | 90.0 | | | | | | | |
| | <i>Solenocera crassicornis</i> | 1 | 6.0 | | | | | 3 | 16.0 | | 1 | 12.0 | | | | |
| | <i>c.f. Thenus sp.</i> | | | | | | | | | | | | | | | |
| Crab | <i>Calappa philargius</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis affinis</i> | 103 | 1010.0 | | 90 | 1530.0 | | 110 | 1830.0 | | 35 | 480.0 | | 54 | 500.0 | |
| | <i>Charybdis annulata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis cruciata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis hellerii</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis japonica</i> | | | | | | | | | | | | | 2 | 123.0 | |
| | <i>Charybdis truncata</i> | | | | | | | 1 | 14.0 | | | | | | | |
| | <i>Charybdis variegata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis spp.</i> | 50 | 460.0 | * | 40 | 410.0 | * | 50 | 460.0 | * | 50 | 460.0 | * | 60 | 590.0 | * |
| | <i>Doclea spp.</i> | | | | | | | | | | | | | | | |
| | <i>Dorippe dorsipes</i> | | | | | | | | | | | | | | | |
| | <i>Dorippe polita</i> | 11 | 96.0 | | 7 | 62.0 | | 30 | 280.0 | * | 30 | 280.0 | * | 20 | 220.0 | * |
| | <i>Eucrater crenata</i> | | | | | | | | | | | | | | | |
| | <i>Eucrater solaris</i> | | | | | | | 1 | 6.0 | | 1 | 7.6 | | | | |
| | <i>Galene bispinosa</i> | 40 | 2310.0 | * | 30 | 1270.0 | * | 7 | 260.0 | | 30 | 1360.0 | * | 20 | 860.0 | * |
| | <i>Maorophthalmus japonicus</i> | | | | | | | | | | 1 | 5.2 | | | | |
| | <i>Parapanope euagora</i> | | | | | | | | | | | | | | | |
| | <i>Parthenope validus</i> | | | | | | | | | | 11 | 130.0 | | | | |
| | <i>Porcellanella picta</i> | | | | | | | | | | | | | | | |
| | <i>Portunus hastatoides</i> | | | | | | | | | | | | | | | |
| | <i>Portunus pelagicus</i> | | | | | | | 1 | 60.0 | | | | | 1 | 80.0 | |
| | <i>Portunus trituberculatus</i> | | | | | | | | | | | | | | | |
| | <i>Scylla paramamosain</i> | | | | | | | | | | 1 | 72.0 | | | | |
| | <i>Typhlocarcinus sp.</i> | | | | | | | | | | | | | | | |
| Mantis shrimp | <i>Dictyosquilla foveolata</i> | 2 | 33.0 | | 1 | 9.0 | | 4 | 55.0 | | | | | 1 | 12.0 | |
| | <i>Harpisquilla harpax</i> | 2 | 85.0 | | 5 | 290.0 | | 4 | 73.0 | | 2 | 101.0 | | 1 | 52.0 | |
| | <i>Oratosquilla interrupta</i> | 53 | 580.0 | | 52 | 703.0 | | 58 | 610.0 | | 23 | 380.0 | | 31 | 504.0 | |
| | <i>Oratosquilla kempii</i> | | | | | | | | | | | | | | | |
| | <i>Oratosquilla oratoria</i> | 4 | 45.0 | | 3 | 40.0 | | 5 | 60.0 | | 2 | 22.0 | | 4 | 60.0 | |
| Gastropod | <i>Calyptreaea sp.</i> | | | | | | | | | | | | | | | |

Trawling Data in August 2021

| Group | Species Name | TNB-1 | | | TNB-2 | | | TNB | | | TNB-4 | | | TNB-5 | | |
|------------|----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Drupa musiva</i> | | | | | | | | | | | | | | | |
| | <i>Lophiotoma leucotropis</i> | | | | | | | | | | | | | | | |
| | <i>Murex trapa</i> | 20 | 160.0 | * | 20 | 160.0 | * | 20 | 160.0 | * | 20 | 180.0 | * | 7 | 68.0 | |
| | <i>Nassarius hepaticus</i> | | | | | | | | | | | | | | | |
| | <i>Nassarius succinctus</i> | | | | | | | | | | | | | | | |
| | <i>Siphonotella walshi</i> | | | | | | | | | | | | | | | |
| | <i>Tomia sulcosa</i> | | | | 1 | 147.0 | | 2 | 187.0 | | | | | | | |
| | <i>Turritella terebra</i> | 190 | 1880.0 | * | 180 | 1780.0 | * | 140 | 1370.0 | * | 380 | 3270.0 | * | 110 | 1070.0 | * |
| | <i>UNID Nudibranch</i> | | | | | | | | | | | | | | | |
| Bivalve | <i>Anomia chinensis</i> | | | | | | | | | | | | | | | |
| | <i>Anomalocardia squamosa</i> | | | | | | | | | | | | | | | |
| | <i>Chlamys spp.</i> | | | | | | | | | | | | | | | |
| | <i>Paphia undulata</i> | | | | | | | | | | | | | | | |
| | <i>Perna viridis</i> | | | | | | | | | | | | | | | |
| | <i>Pteria sp.</i> | | | | | | | | | | | | | | | |
| | <i>Ruditapes philippinarum</i> | | | | | | | | | | | | | | | |
| | <i>Tegillarca nodifera</i> | | | | | | | | | | | | | | | |
| Cephalopod | <i>Lolingid squid</i> | 1 | 30.0 | | | | | | | | | | | | | |
| | <i>Sepiid cuttlefish</i> | | | | | | | | | | | | | | | |
| | <i>Octopus</i> | | | | | | | | | | | | | | | |
| Echinoderm | <i>Acaudina molpadioides</i> | 3 | 68.0 | | 2 | 51.0 | | 3 | 72.0 | | 6 | 140.0 | | 2 | 54.0 | |
| | <i>Comanthus sp.</i> | | | | | | | | | | | | | | | |
| | <i>Styela plicata</i> | | | | | | | | | | | | | | | |
| | <i>Temnopleurus toreumaticus</i> | | | | | | | | | | | | | | | |
| Cnidaria | <i>Carcinacis ichikawai</i> | | | | | | | | | | | | | | | |
| | <i>Cavernularia habereri</i> | 20 | 120.0 | * | 20 | 120.0 | * | 6 | 25.0 | | 20 | 130.0 | * | 20 | 130.0 | * |
| | <i>Ellisella laevis</i> | | | | | | | | | | | | | | | |
| | <i>Pteroeides chinense</i> | 60 | 1410.0 | * | 74 | 1308.0 | * | 80 | 2600.0 | * | 60 | 1340.0 | * | 50 | 1190.0 | * |
| | <i>Gorgonia spp.</i> | | | | | | | | | | | | | | | |
| | Jellyfish | | | | | | | | | | 5 | 1780.0 | | 1 | 240.0 | |
| | <i>Virgularia gustaviana</i> | 2 | 9.6 | | | | | | | | | | | 2 | 19.4 | |
| Cirripedia | <i>c.f. Balanus spp.</i> | | | | | | | | | | | | | | | |

Trawling Data in August 2021

| Group | Species Name | TSA-1 | | | TSA-2 | | | TSA-3 | | | TSA-4 | | | TSA-5 | | | |
|-------|----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------|
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | |
| Fish | <i>Arnoglossus waitei</i> | | | | | | | | | | | | | | | | |
| | <i>Anchoviella commersonii</i> | | | | | | | 1 | 9.0 | | | 1 | 5.0 | | | | |
| | <i>Apogon kiensis</i> | 12 | 60.0 | | 5 | 28.0 | | 21 | 141.0 | | | 6 | 21.0 | | | 17 | 92.0 |
| | <i>Apogon lineatus</i> | | | | | | | | | | | | | | | | |
| | <i>Arothron hispidus</i> | | | | | | | | | | | | | | | | |
| | <i>Arius chinensis</i> | 6 | 960.0 | | 6 | 1460.0 | | 7 | 1320.0 | | | | | | | 1 | 140.0 |
| | <i>Butis koilomatodon</i> | | | | | | | | | | | | | | | | |
| | <i>Callionymus curvicornis</i> | | | | | | | 1 | 3.0 | | | 1 | 6.0 | | | | |
| | <i>Caranx kalla</i> | | | | | | | | | | | | | | | | |
| | <i>Chrysochir aureus</i> | 1 | 60.0 | | | | | | | | | 2 | 240.0 | | | 2 | 470.0 |
| | <i>Clupanodon thrissa</i> | | | | | | | | | | | 3 | 290.0 | | | 3 | 215.0 |
| | <i>Coilia mystus</i> | | | | | | | | | | | | | | | | |
| | <i>Collichthys lucidus</i> | | | | | | | | | | | | | | | | |
| | <i>Cryptocentrus filifer</i> | 6 | 33.0 | | 4 | 22.0 | | 8 | 41.0 | | | 4 | 24.0 | | | 2 | 12.0 |
| | <i>Cynoglossus arel</i> | 1 | 50.0 | | 2 | 36.0 | | 12 | 190.0 | | | 13 | 410.0 | | | 11 | 260.0 |
| | <i>Cynoglossus puncticeps</i> | | | | | | | | | | | | | | | | |
| | <i>Cynoglossus roulei</i> | | | | | | | | | | | 1 | 870.0 | | | | |
| | <i>Dasvatis akajei</i> | | | | | | | | | | | | | | | | |
| | <i>Dasvatis zugei</i> | 9 | 890.0 | | 11 | 1110.0 | | 5 | 2740.0 | | | | | | | 2 | 62.0 |
| | <i>Dendrophysa russelli</i> | 1 | 8.0 | | 1 | 8.0 | | 3 | 28.0 | | | 1 | 20.0 | | | 4 | 80.0 |
| | <i>Drepane punctata</i> | | | | | | | | | | | | | | | | |
| | <i>Epinephelus awoara</i> | 8 | 52.0 | | 7 | 35.0 | | 13 | 70.0 | | | 3 | 20.0 | | | 9 | 41.0 |
| | <i>Gerres abbreviatus</i> | 2 | 230.0 | | 1 | 80.0 | | | | | | | | | | | |
| | <i>Gerres lucidus</i> | 3 | 60.0 | | 4 | 70.0 | | 6 | 135.0 | | | 1 | 30.0 | | | 1 | 20.0 |
| | <i>Grammolites scaber</i> | | | | | | | | | | | | | | | | |
| | <i>Gymnura japonica</i> | | | | | | | | | | | | | | | | |
| | <i>Harpadon nehereus</i> | | | | | | | | | | | | | | | | |
| | <i>Ilisha elongata</i> | | | | | | | 2 | 11.0 | | | 2 | 11.0 | | | 1 | 6.0 |
| | <i>Inimicus japonicus</i> | | | | | | | | | | | | | | | | |
| | <i>Johnius belangerii</i> | | | | | | | | | | | 1 | 60.0 | | | | |
| | <i>Konosirus punctatus</i> | 27 | 1010.0 | | 25 | 850.0 | | 21 | 820.0 | | | 4 | 230.0 | | | 8 | 480.0 |
| | <i>Lagocephalus inermis</i> | | | | | | | | | | | | | | | | |
| | <i>Lagocephalus sceleratus</i> | | | | | | | | | | | | | | | | |
| | <i>Larimichthys polyactis</i> | 1 | 410.0 | | | | | | | | | | | | | | |
| | <i>Leiognathus nuchalis</i> | 187 | 990.0 | | 122 | 950.0 | | 120 | 1420.0 | | | 83 | 1050.0 | | | 73 | 890.0 |
| | <i>Lisa macrolepis</i> | | | | | | | | | | | 1 | 30.0 | | | | |
| | <i>Lutjanus sp.</i> | | | | | | | | | | | | | | | 1 | 77.0 |
| | <i>Muraenesox cinereus</i> | 2 | 177.0 | | 1 | 90.0 | | | | | | 1 | 180.0 | | | | |
| | <i>Nemipterus japonicus</i> | 1 | 13.0 | | 2 | 22.0 | | 3 | 31.0 | | | | | | | | |
| | <i>Oxyurichthys tentacularis</i> | | | | | | | | | | | | | | | | |
| | <i>Paramonacanthus polynema</i> | | | | | | | 3 | 40.0 | | | 3 | 48.0 | | | 2 | 40.0 |
| | <i>Pennahia anea</i> | 19 | 320.0 | | 23 | 370.0 | | 42 | 640.0 | | | 79 | 1620.0 | | | 60 | 1210.0 |
| | <i>Pisodonophis boro</i> | | | | | | | | | | | 1 | 150.0 | | | | |
| | <i>Platycephalus indicus</i> | 12 | 513.0 | | 8 | 455.0 | | 19 | 1550.0 | | | 11 | 1350.0 | | | 14 | 1980.0 |
| | <i>Pseudorhombus cinnamomeus</i> | | | | | | | | | | | | | | | | |
| | <i>Pseudorhombus oligodon</i> | | | | 1 | 60.0 | | | | | | | | | | 1 | 30.0 |
| | <i>Saurida elongata</i> | | | | 2 | 94.0 | | 1 | 70.0 | | | | | | | | |
| | <i>Scatophagus argus</i> | 2 | 670.0 | | 1 | 450.0 | | | | | | 1 | 220.0 | | | | |
| | <i>Sebastiscus marmoratus</i> | 1 | 150.0 | | 1 | 110.0 | | 2 | 76.0 | | | | | | | | |
| | <i>Siganus fuscescens</i> | | | | | | | | | | | | | | | | |
| | <i>Sillago sihama</i> | | | | 1 | 50.0 | | | | | | | | | | | |
| | <i>Solea ovata</i> | 1 | 20.0 | | | | | | | | | | | | | 1 | 5.0 |
| | <i>Stephanolepis cirrhifer</i> | | | | | | | | | | | | | | | | |

Trawling Data in August 2021

| Group | Species Name | TSA-1 | | | TSA-2 | | | TSA | | | TSA-4 | | | TSA-5 | | |
|-----------------|------------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Takifugu niphobles</i> | 2 | 1070.0 | | | | | | | | | | | | | |
| | <i>Takifugu oblongus</i> | | | | | | | | | | | | | | | |
| | <i>Terapon jarbua</i> | | | | | | | | | | | | | | | |
| | <i>Terapon theraps</i> | | | | | | | | | | | | | | | |
| | <i>Thryssa kammalensis</i> | | | | | | | | | | 23 | 72.0 | | 22 | 83.0 | |
| | <i>Trachicephalus uranoscopus</i> | 1 | 13.0 | | | | | 7 | 90.0 | | 5 | 90.0 | | 3 | 50.0 | |
| | <i>Trypauchen vagina</i> | 2 | 16.0 | | | | | 2 | 12.0 | | 2 | 30.0 | | 4 | 50.0 | |
| Prawn or shrimp | <i>Alpheus brevicristatus</i> | | | | | | | | | | | | | | | |
| | <i>Alpheus distinguendus</i> | | | | | | | | | | | | | | | |
| | <i>Alpheus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Metapenaeopsis barbata</i> | | | | 1 | 7.0 | | | | | | | | | | |
| | <i>Metapenaeus affinis</i> | 17 | 80.0 | | 10 | 40.0 | | 13 | 60.0 | | 11 | 60.0 | | 9 | 30.0 | |
| | <i>Metapenaeus ensis</i> | | | | | | | | | | | | | | | |
| | <i>Metapenaeus joyneri</i> | | | | | | | | | | 1 | 12.0 | | | | |
| | <i>Parapenaeopsis hardwickii</i> | | | | | | | | | | | | | | | |
| | <i>Parapenaeopsis hungerfordii</i> | 3 | 12.0 | | 3 | 33.0 | | 7 | 34.0 | | 7 | 41.0 | | 4 | 22.0 | |
| | <i>Parapenaeopsis tenella</i> | 5 | 33.0 | | | | | 4 | 33.0 | | 3 | 22.0 | | 10 | 85.0 | |
| | <i>Penaeus japonicus</i> | | | | | | | | | | | | | 1 | 25.0 | |
| | <i>Penaeus merguensis</i> | 2 | 40.0 | | 3 | 35.0 | | 5 | 90.0 | | 10 | 140.0 | | 13 | 220.0 | |
| | <i>Penaeus penicillatus</i> | | | | | | | | | | | | | | | |
| | <i>Solenocera crassicornis</i> | | | | | | | | | | | | | | | |
| | <i>c.f. Thenus sp.</i> | | | | | | | | | | | | | | | |
| Crab | <i>Calappa philargius</i> | | | | | | | | | | | | | | | |
| | <i>Charabydis affinis</i> | 84 | 910.0 | | 35 | 470.0 | | 156 | 1530.0 | | 120 | 1180.0 | | 110 | 1160.0 | |
| | <i>Charabydis annulata</i> | 11 | 26.0 | | | | | | | | 1 | 8.0 | | | | |
| | <i>Charabydis cruciata</i> | | | | | | | 1 | 15.0 | | 4 | 96.0 | | 5 | 196.0 | |
| | <i>Charabydis hellerii</i> | 2 | 10.0 | | | | | | | | | | | | | |
| | <i>Charabydis japonica</i> | | | | | | | 4 | 18.0 | | | | | | | |
| | <i>Charabydis truncata</i> | | | | | | | 1 | 10.0 | | | | | | | |
| | <i>Charabydis variegata</i> | | | | | | | | | | | | | 3 | 21.0 | |
| | <i>Charabydis spp.</i> | 30 | 310.0 | * | 40 | 420.0 | * | 30 | 310.0 | * | 50 | 520.0 | * | 40 | 410.0 | * |
| | <i>Doclea spp.</i> | | | | | | | | | | | | | | | |
| | <i>Dorippe dorsipes</i> | | | | | | | | | | | | | | | |
| | <i>Dorippe polita</i> | 20 | 120.0 | | 30 | 164.0 | * | 16 | 104.0 | | 14 | 96.0 | | 8 | 76.0 | |
| | <i>Eucrate crenata</i> | 1 | 7.2 | | | | | | | | | | | | | |
| | <i>Eucrate solaris</i> | | | | | | | | | | | | | | | |
| | <i>Galene hispinosa</i> | 13 | 558.0 | | 14 | 570.0 | | 9 | 410.0 | | 12 | 560.0 | | 18 | 630.0 | |
| | <i>Maorophthalmus japonicus</i> | | | | | | | 5 | 36.0 | | | | | 3 | 13.7 | |
| | <i>Parapanope euagora</i> | | | | | | | | | | | | | | | |
| | <i>Parthenope validus</i> | | | | | | | | | | | | | | | |
| | <i>Porcellanella picta</i> | 6 | 2.1 | | | | | 2 | 0.7 | | 4 | 1.8 | | | | |
| | <i>Portunus hastatoides</i> | 9 | 19.0 | | 3 | 10.0 | | 11 | 28.0 | | 15 | 25.0 | | 4 | 13.0 | |
| | <i>Portunus pelagicus</i> | 2 | 160.0 | | 2 | 190.0 | | 3 | 160.0 | | 3 | 210.0 | | 3 | 310.0 | |
| | <i>Portunus trituberculatus</i> | 1 | 190.0 | | 1 | 260.0 | | | | | 2 | 350.0 | | | | |
| | <i>Scylla paramamosain</i> | | | | | | | | | | | | | | | |
| | <i>Typhlocarcinus sp.</i> | | | | | | | 1 | 2.2 | | | | | | | |
| Mantis shrimp | <i>Dictyosquilla foveolata</i> | | | | | | | | | | | | | 1 | 14.0 | |
| | <i>Harpiosquilla harpax</i> | 2 | 18.0 | | | | | 6 | 280.0 | | 9 | 210.0 | | 5 | 130.0 | |
| | <i>Oratosquilla interrupta</i> | | | | 1 | 23.0 | | 1 | 20.0 | | 1 | 20.0 | | 1 | 20.0 | |
| | <i>Oratosquilla kempii</i> | | | | | | | | | | | | | 1 | 11.0 | |
| | <i>Oratosquilla oratoria</i> | | | | | | | | | | 1 | 10.0 | | | | |
| Gastropod | <i>Calyptraea sp.</i> | | | | | | | | | | | | | | | |

Trawling Data in August 2021

| Group | Species Name | TSA-1 | | | TSA-2 | | | TSA | | | TSA-4 | | | TSA-5 | | |
|------------|----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Drupa musiva</i> | 3 | 18.0 | | | | | | | | | | | | | |
| | <i>Lophiotoma leucotropis</i> | | | | | | | | | | | | | | | |
| | <i>Murex trapa</i> | 18 | 166.0 | | 10 | 97.0 | | 7 | 69.0 | | | | | 8 | 22.0 | |
| | <i>Nassarius hepaticus</i> | | | | | | | | | | | | | | | |
| | <i>Nassarius succinctus</i> | 2 | 5.1 | | 6 | 143.0 | | | | | | | | | | |
| | <i>Siphonotus walshi</i> | | | | | | | | | | | | | | | |
| | <i>Tonna sulcosa</i> | | | | 1 | 114.0 | | | | | | | | 1 | 152.0 | |
| | <i>Turritella terebra</i> | 40 | 390.0 | * | 30 | 280.0 | * | 30 | 280.0 | * | 30 | 270.0 | * | 30 | 260.0 | * |
| | <i>UNID Nudibranch</i> | | | | | | | | | | | | | | | |
| Bivalve | <i>Anomia chinensis</i> | | | | | | | | | | | | | | | |
| | <i>Anomalocardia squamosa</i> | | | | | | | | | | | | | | | |
| | <i>Chlamys spp.</i> | | | | | | | | | | | | | | | |
| | <i>Paphia undulata</i> | | | | | | | | | | | | | 5 | 31.0 | |
| | <i>Perna viridis</i> | | | | | | | | | | | | | | | |
| | <i>Pteria sp.</i> | | | | | | | | | | | | | | | |
| | <i>Ruditapes philippinarum</i> | | | | | | | | | | | | | | | |
| | <i>Tegillarca nodifera</i> | 4 | 26.0 | | 2 | 79.0 | | | | | | | | | | |
| Cephalopod | <i>Lolingid squid</i> | | | | | | | | | | | | | | | |
| | <i>Sepiid cuttlefish</i> | 8 | 810.0 | | 7 | 830.0 | | 10 | 1050.0 | | 3 | 303.0 | | 3 | 310.0 | |
| | <i>Octopus</i> | | | | | | | 1 | 70.0 | | | | | | | |
| Echinoderm | <i>Acaudina molpadioides</i> | 12 | 280.0 | | 16 | 390.0 | | 20 | 610.0 | * | 6 | 160.0 | | 5 | 110.0 | |
| | <i>Comanthus sp.</i> | | | | | | | | | | | | | | | |
| | <i>Styela plicata</i> | | | | | | | | | | | | | | | |
| | <i>Temnopleurus toreumaticus</i> | | | | | | | | | | | | | | | |
| Cnidaria | <i>Carcinactis ichikawai</i> | | | | 3 | 1.3 | | | | | | | | | | |
| | <i>Cavernularia habereri</i> | 16 | 70.0 | | 22 | 110.0 | | 20 | 106.0 | * | 16 | 92.0 | | 9 | 72.0 | |
| | <i>Ellisella laevis</i> | | | | | | | | | | | | | | | |
| | <i>Pteroeides chinense</i> | 80 | 2410.0 | * | 80 | 2460.0 | * | 90 | 2160.0 | * | 90 | 3210.0 | * | 80 | 2640.0 | * |
| | <i>Gorgonia spp.</i> | | | | | | | | | | | | | | | |
| | Jellyfish | | | | | | | | | | | | | | | |
| | <i>Virgularia gustaviana</i> | | | | | | | 2 | 10.0 | | | | | 2 | 11.6 | |
| Cirripedia | <i>c.f. Balanus spp.</i> | | | | | | | | | | | | | | | |

Trawling Data in August 2021

| Group | Species Name | TSB | | | | | | | | | | | | | | |
|-------|----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | TSB-1 | | | TSB-2 | | | TSB-3 | | | TSB-4 | | | TSB-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| Fish | <i>Arnoglossus waitei</i> | | | | 2 | 20.0 | | 1 | 13.0 | | | | | | | |
| | <i>Anchoviella commersonii</i> | 4 | 25.0 | | 1 | 9.0 | | 8 | 42.0 | | | | | | | |
| | <i>Apogon kiensis</i> | 5 | 26.0 | | 6 | 21.0 | | 7 | 22.0 | | | | | 1 | 13.0 | |
| | <i>Apogon lineatus</i> | | | | | | | 1 | 6.0 | | | | | | | |
| | <i>Arothron hispidus</i> | | | | 1 | 798.0 | | | | | | | | | | |
| | <i>Arius chinensis</i> | 1 | 130.0 | | 4 | 510.0 | | 3 | 320.0 | | 5 | 1050.0 | | 2 | 360.0 | |
| | <i>Butis koilomatodon</i> | | | | | | | | | | | | | | | |
| | <i>Callionymus curvicornis</i> | | | | | | | | | | | | | 1 | 4.0 | |
| | <i>Caranx kalla</i> | | | | | | | | | | | | | | | |
| | <i>Chrysochir aureus</i> | | | | | | | | | | 1 | 30.0 | | | | |
| | <i>Clupanodon thrissa</i> | | | | | | | | | | | | | | | |
| | <i>Coilia mystus</i> | | | | | | | | | | | | | | | |
| | <i>Collichthys lucidus</i> | | | | 2 | 50.0 | | | | | | | | | | |
| | <i>Cryptocentrus filifer</i> | | | | 2 | 11.0 | | 3 | 11.0 | | | | | 1 | 11.0 | |
| | <i>Cynoglossus arel</i> | 1 | 10.0 | | 2 | 40.0 | | 6 | 110.0 | | 2 | 40.0 | | 7 | 150.0 | |
| | <i>Cynoglossus puncticeps</i> | | | | | | | | | | | | | | | |
| | <i>Cynoglossus roulei</i> | | | | | | | | | | | | | | | |
| | <i>Dasvatis akajei</i> | | | | | | | | | | | | | | | |
| | <i>Dasvatis zugei</i> | | | | 6 | 778.0 | | 4 | 960.0 | | | | | 2 | 620.0 | |
| | <i>Dendrophysa russelli</i> | 2 | 23.0 | | 5 | 55.0 | | 8 | 70.0 | | 3 | 60.0 | | 3 | 90.0 | |
| | <i>Drepane punctata</i> | | | | | | | | | | | | | | | |
| | <i>Epinephelus awoara</i> | 19 | 101.0 | | 16 | 93.0 | | 57 | 215.0 | | 5 | 23.0 | | 4 | 21.0 | |
| | <i>Gerres abbreviatus</i> | | | | | | | | | | | | | | | |
| | <i>Gerres lucidus</i> | 2 | 50.0 | | | | | | | | | | | 1 | 25.0 | |
| | <i>Grammolites scaber</i> | | | | 1 | 58.0 | | 1 | 70.0 | | | | | | | |
| | <i>Gymnura japonica</i> | 1 | 806.0 | | | | | 1 | 470.0 | | 1 | 220.0 | | | | |
| | <i>Harpadon nehereus</i> | | | | | | | | | | | | | | | |
| | <i>Ilisha elongata</i> | 2 | 18.0 | | 2 | 10.0 | | 6 | 25.0 | | | | | | | |
| | <i>Inimicus japonicus</i> | | | | | | | 1 | 120.0 | | | | | | | |
| | <i>Johnius belangerii</i> | 3 | 40.0 | | 1 | 10.0 | | 1 | 50.0 | | | | | 1 | 20.0 | |
| | <i>Konosirus punctatus</i> | 2 | 80.0 | | 9 | 430.0 | | 5 | 220.0 | | 3 | 210.0 | | | | |
| | <i>Lagocephalus inermis</i> | | | | | | | | | | | | | | | |
| | <i>Lagocephalus sceleratus</i> | | | | | | | 4 | 200.0 | | | | | | | |
| | <i>Larimichthys polyactis</i> | | | | | | | | | | | | | | | |
| | <i>Leiognathus nuchalis</i> | 110 | 1370.0 | | 150 | 2090.0 | | 180 | 2720.0 | | 95 | 1990.0 | | 80 | 1480.0 | |
| | <i>Lisa macrolepis</i> | | | | | | | | | | | | | | | |
| | <i>Lutjanus sp.</i> | | | | | | | | | | | | | | | |
| | <i>Muraenesox cinereus</i> | 1 | 102.0 | | | | | | | | | | | | | |
| | <i>Nemipterus japonicus</i> | | | | | | | | | | 1 | 18.0 | | | | |
| | <i>Oxyurichthys tentacularis</i> | | | | | | | | | | | | | 1 | 14.0 | |
| | <i>Paramonacanthus polynema</i> | 1 | 15.0 | | 1 | 20.0 | | 1 | 18.0 | | | | | | | |
| | <i>Pennahia anea</i> | 21 | 450.0 | | 25 | 570.0 | | 29 | 540.0 | | 20 | 404.0 | | 22 | 450.0 | |
| | <i>Pisodonophis boro</i> | | | | | | | | | | | | | | | |
| | <i>Platycephalus indicus</i> | | | | 7 | 790.0 | | 4 | 560.0 | | 1 | 20.0 | | 1 | 420.0 | |
| | <i>Pseudorhombus cinnamomeus</i> | | | | 1 | 40.0 | | | | | | | | | | |
| | <i>Pseudorhombus oligodon</i> | | | | | | | | | | | | | 2 | 56.0 | |
| | <i>Saurida elongata</i> | | | | | | | | | | | | | | | |
| | <i>Scatophagus argus</i> | | | | 1 | 130.0 | | | | | | | | | | |
| | <i>Sebastes marmoratus</i> | 1 | 24.0 | | | | | | | | 1 | 15.0 | | | | |
| | <i>Siganus fuscescens</i> | | | | | | | 5 | 120.0 | | | | | 1 | 38.0 | |
| | <i>Sillago sihama</i> | | | | | | | | | | | | | | | |
| | <i>Solea ovata</i> | | | | 1 | 4.0 | | | | | | | | | | |
| | <i>Stephanolepis cirrhifer</i> | | | | | | | | | | | | | | | |

Trawling Data in August 2021

| Group | Species Name | TSB | | | | | | | | | | | | | | |
|-----------------|-----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | TSB-1 | | | TSB-2 | | | TSB-3 | | | TSB-4 | | | TSB-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Takifugu niphobles</i> | | | | | | | | | | | | | | | |
| | <i>Takifugu oblongus</i> | | | | | | | | | | | | | | | |
| | <i>Terapon jarbua</i> | | | | | | | | | | | | | 1 | 17.0 | |
| | <i>Terapon theraps</i> | | | | | | | | | | | | | | | |
| | <i>Thryssa kammalensis</i> | 2 | 8.0 | | 1 | 6.0 | | 38 | 130.0 | | 1 | 12.0 | | 5 | 33.0 | |
| | <i>Trachicephalus uranoscopus</i> | 1 | 14.0 | | 1 | 19.0 | | 3 | 44.0 | | | | | 3 | 36.0 | |
| | <i>Trypauchen vagina</i> | | | | 1 | 5.0 | | 3 | 60.0 | | 1 | 20.0 | | 3 | 40.0 | |
| Prawn or shrimp | <i>Alpheus brevicristatus</i> | | | | | | | | | | 1 | 8.0 | | | | |
| | <i>Alpheus distinguendus</i> | | | | | | | | | | | | | | | |
| | <i>Alpheus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Metapenaeopsis barbata</i> | | | | | | | | | | | | | | | |
| | <i>Metapenaeus affinis</i> | 10 | 40.0 | | 11 | 80.0 | | 19 | 148.0 | | 12 | 96.0 | | 4 | 32.0 | |
| | <i>Metapenaeus ensis</i> | | | | | | | | | | | | | | | |
| | <i>Metapenaeus joyneri</i> | 1 | 12.0 | | | | | 3 | 20.0 | | | | | | | |
| | <i>Parapeneopsis hardwickii</i> | | | | | | | | | | | | | | | |
| | <i>Parapeneopsis hungerfordii</i> | 3 | 31.0 | | 4 | 22.0 | | 7 | 41.0 | | 4 | 23.0 | | 3 | 22.0 | |
| | <i>Parapeneopsis tenella</i> | 2 | 22.0 | | 7 | 76.0 | | 2 | 30.0 | | | | | | | |
| | <i>Penaeus japonicus</i> | | | | 1 | 13.0 | | | | | | | | | | |
| | <i>Penaeus merguensis</i> | 7 | 110.0 | | 11 | 170.0 | | 13 | 250.0 | | 18 | 310.0 | | 34 | 620.0 | |
| | <i>Penaeus penicillatus</i> | | | | | | | | | | 4 | 70.0 | | 3 | 28.0 | |
| | <i>Solenocera crassicornis</i> | | | | | | | | | | | | | | | |
| | <i>c.f. Themis sp.</i> | | | | | | | | | | | | | | | |
| Crab | <i>Calappa philargius</i> | | | | | | | | | | 1 | 18.3 | | | | |
| | <i>Charybdis affinis</i> | 65 | 490.0 | | 63 | 700.0 | | 105 | 1620.0 | | 22 | 210.0 | | 18 | 260.0 | |
| | <i>Charybdis annulata</i> | | | | 2 | 12.0 | | | | | | | | | | |
| | <i>Charybdis cruciata</i> | | | | 3 | 130.0 | | | | | | | | | | |
| | <i>Charybdis hellerii</i> | 2 | 9.0 | | 4 | 16.0 | | | | | | | | | | |
| | <i>Charybdis japonica</i> | 2 | 80.0 | | 5 | 35.0 | | 2 | 14.0 | | | | 2 | 9.0 | | |
| | <i>Charybdis truncata</i> | | | | 4 | 22.0 | | 3 | 13.0 | | | | | | | |
| | <i>Charybdis variegata</i> | | | | 2 | 9.0 | | 1 | 3.0 | | | | | | | |
| | <i>Charybdis spp.</i> | 40 | 420.0 | * | 30 | 320.0 | * | 30 | 320.0 | * | 40 | 380.0 | * | 30 | 330.0 | * |
| | <i>Doclea spp.</i> | | | | | | | | | | | | | | | |
| | <i>Dorippe dorsipes</i> | | | | | | | | | | | | | | | |
| | <i>Dorippe polita</i> | 9 | 84.0 | | 7 | 68.0 | | 30 | 209.0 | * | 6 | 49.0 | | 3 | 27.0 | * |
| | <i>Eucrate crenata</i> | | | | 2 | 19.0 | | | | | | | | | | |
| | <i>Eucrate solaris</i> | | | | | | | | | | | | | | | |
| | <i>Galene bispinosa</i> | | | | 2 | 66.0 | | 3 | 112.0 | | | | | 1 | 70.0 | |
| | <i>Maorophthalmus japonicus</i> | 5 | 38.4 | | 2 | 8.2 | | 4 | 32.4 | | | | | | | |
| | <i>Parapanope euagora</i> | | | | | | | | | | | | | | | |
| | <i>Parthenope validus</i> | | | | | | | | | | 5 | 51.8 | | 7 | 69.4 | |
| | <i>Porcellanella picta</i> | 2 | 0.9 | | | | | | | | | | | | | |
| | <i>Portunus hastatoides</i> | 15 | 33.0 | | 25 | 38.0 | | 12 | 25.0 | | 2 | 7.0 | | 8 | 11.0 | |
| | <i>Portunus pelagicus</i> | | | | 1 | 80.0 | | 3 | 180.0 | | 1 | 80.0 | | | | |
| | <i>Portunus trituberculatus</i> | | | | | | | 1 | 100.0 | | 1 | 110.0 | | | | |
| | <i>Scylla paramamosain</i> | | | | | | | | | | | | | | | |
| | <i>Typhlocarcinus sp.</i> | | | | | | | 1 | 1.6 | | | | | | | |
| Mantis shrimp | <i>Dictyosquilla foveolata</i> | 1 | 8.0 | | | | | 3 | 28.0 | | 1 | 12.0 | | 1 | 16.0 | |
| | <i>Harpiosquilla harpax</i> | 4 | 44.0 | | 6 | 28.0 | | 14 | 202.0 | | 4 | 106.0 | | 4 | 150.0 | |
| | <i>Oratosquilla interrupta</i> | 1 | 36.0 | | | | | 1 | 30.0 | | 9 | 230.0 | | 3 | 100.0 | |
| | <i>Oratosquilla kempii</i> | 1 | 5.0 | | 4 | 22.0 | | 2 | 16.0 | | 3 | 14.0 | | 2 | 7.0 | |
| | <i>Oratosquilla oratoria</i> | | | | | | | 1 | 10.0 | | 1 | 15.0 | | | | |
| Gastropod | <i>Calyptrea sp.</i> | | | | | | | | | | | | | | | |

Trawling Data in August 2021

| Group | Species Name | TSB-1 | | | TSB-2 | | | TSB | | | TSB-4 | | | TSB-5 | | |
|------------|----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Drupa musiva</i> | | | | | | | | | | | | | | | |
| | <i>Lophiotoma leucotropis</i> | | | | | | | | | | | | | | | |
| | <i>Murex trapa</i> | 13 | 121.0 | | 19 | 152.0 | | 18 | 148.0 | | 6 | 52.0 | | | | |
| | <i>Nassarius hepaticus</i> | 1 | 2.3 | | | | | | | | | | | | | |
| | <i>Nassarius succinctus</i> | 12 | 25.4 | | 30 | 74.0 | * | 30 | 74.0 | * | 5 | 10.4 | | | | |
| | <i>Siphonotella walshi</i> | | | | | | | | | | | | | | | |
| | <i>Tonna sulcosa</i> | | | | | | | | | | | | | | | |
| | <i>Turritella terebra</i> | 30 | 270.0 | * | 30 | 270.0 | * | 40 | 380.0 | * | 30 | 260.0 | * | 30 | 260.0 | * |
| | <i>UNID Nudibranch</i> | | | | 2 | 19.0 | | | | | | | | | | |
| Bivalve | <i>Anomia chinensis</i> | | | | | | | | | | | | | | | |
| | <i>Anomalocardia squamosa</i> | 2 | 4.2 | | | | | 4 | 9.2 | | | | | | | |
| | <i>Chlamys spp.</i> | | | | | | | | | | | | | | | |
| | <i>Paphia undulata</i> | | | | | | | | | | | | | | | |
| | <i>Perna viridis</i> | | | | | | | | | | | | | | | |
| | <i>Pteria sp.</i> | | | | | | | | | | | | | | | |
| | <i>Ruditapes philippinarum</i> | | | | | | | | | | | | | | | |
| | <i>Tegillarca nodifera</i> | | | | | | | | | | | | | | | |
| Cephalopod | <i>Lolingid squid</i> | | | | | | | | | | | | | | | |
| | <i>Sepiid cuttlefish</i> | 1 | 10.0 | | 2 | 220.0 | | | | | | | | | | |
| | <i>Octopus</i> | | | | | | | 1 | 60.0 | | 1 | 100.0 | | 1 | 77.0 | |
| Echinoderm | <i>Acaudina molpadioides</i> | 12 | 260.0 | | 20 | 590.0 | * | 16 | 310.0 | | 8 | 170.0 | | 13 | 240.0 | |
| | <i>Comanthus sp.</i> | | | | | | | | | | | | | | | |
| | <i>Styela plicata</i> | | | | | | | | | | | | | | | |
| | <i>Temnopleurus toreumaticus</i> | | | | | | | | | | 210 | 2040.0 | * | 220 | 2130.0 | * |
| Cnidaria | <i>Carcinactis ichikawai</i> | 2 | 1.2 | | 3 | 2.1 | | | | | | | | | | |
| | <i>Cavernularia habereri</i> | 10 | 52.0 | | 11 | 78.0 | | 20 | 130.0 | * | 11 | 56.0 | | 13 | 62.0 | |
| | <i>Ellisella laevis</i> | | | | | | | | | | | | | | | |
| | <i>Pteroeides chinense</i> | 50 | 1600.0 | * | 40 | 1300.0 | * | 40 | 1300.0 | * | 30 | 620.0 | * | 36 | 620.0 | * |
| | <i>Gorgonia spp.</i> | | | | | | | | | | | | | | | |
| | Jellyfish | | | | | | | | | | | | | | | |
| | <i>Virgularia gustaviana</i> | | | | 1 | 0.2 | | | | | 3 | 21.0 | | 2 | 18.0 | |
| Cirripedia | <i>c.f. Balanus spp.</i> | | | | | | | | | | | | | | | |