

Trawling Data in July 2019

| Group | Species Name | INA | | | | | | | | | | | | | | |
|-------|--------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | INA-1 | | | INA-2 | | | INA-3 | | | INA-4 | | | INA-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| Fish | <i>Arnoglossus waitei</i> | | | | | | | | | | | | | | | |
| | <i>Acentrogobius caninus</i> | | | | | | | | | | | | | | | |
| | <i>Anchoviella commersonii</i> | | | | | | | | | | | | | | | |
| | <i>Apogon kiensis</i> | 1 | 3 | | | | | 2 | 10 | | | | | | | |
| | <i>Apogon lineatus</i> | | | | 2 | 10 | | | | | | | | 2 | 10 | |
| | <i>Arius arius</i> | | | | 2 | 70 | | | | | | | | | | |
| | <i>Callionymus richardsoni</i> | | | | | | | | | | | | | | | |
| | <i>Callionymus curvicornis</i> | | | | 1 | 12 | | | | | 1 | 10 | | | | |
| | <i>Caranx kalla</i> | | | | | | | | | | | | | | | |
| | <i>Clupanodon thrissa</i> | | | | | | | | | | | | | | | |
| | <i>Collichthys lucidus</i> | | | | | | | 1 | 5 | | | | | | | |
| | <i>Coilia mystus</i> | | | | | | | | | | | | | | | |
| | <i>Crossorhombus kanekonis</i> | | | | | | | | | | | | | | | |
| | <i>Cryptocentrus filifer</i> | | | | | | | | | | | | | | | |
| | <i>Cynoglossus abbreviatus</i> | | | | | | | | | | | | | | | |
| | <i>Cynoglossus arel</i> | | | | 1 | 8 | | | | | | | | 1 | 30 | |
| | <i>Cynoglossus puncticeps</i> | | | | | | | | | | | | | 3 | 40 | |
| | <i>Cynoglossus roulei</i> | | | | | | | | | | | | | | | |
| | <i>Dasyatis akajei</i> | 2 | 610 | | | | | | | | | | | | | |
| | <i>Dasyatis zugei</i> | | | | | | | | | | 1 | 230 | | 3 | 520 | |
| | <i>Drepane punctata</i> | | | | | | | | | | 1 | 170 | | | | |
| | <i>Dendrophysa russelii</i> | | | | | | | | | | | | | | | |
| | <i>Epinephelus awoara</i> | | | | 1 | 5 | | | | | | | | | | |
| | <i>Epinephelus bruneus</i> | | | | | | | | | | | | | | | |
| | <i>Erisphex pottii</i> | 1 | 8 | | | | | 1 | 10 | | | | | | | |
| | <i>Gerres lucidus</i> | | | | 1 | 50 | | | | | | | | | | |
| | <i>Grammoplites scaber</i> | 1 | 40 | | 1 | 110 | | 2 | 90 | | | | | 2 | 80 | |
| | <i>Gymnura japonica</i> | | | | | | | | | | | | | | | |
| | <i>Ilisha elongata</i> | | | | | | | | | | | | | | | |
| | <i>Inegocia japonicus</i> | 1 | 20 | | 1 | 40 | | 2 | 50 | | | | | | | |
| | <i>Inimicus didactylus</i> | | | | 1 | 280 | | | | | | | | | | |
| | <i>Inimicus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Johnius belangerii</i> | | | | 3 | 130 | | | | | | | | | | |
| | <i>Johnius carona</i> | | | | | | | | | | | | | | | |
| | <i>Johnius poecilonotus</i> | | | | 1 | 260 | | | | | | | | | | |
| | <i>Johnius taiwanense</i> | | | | 1 | 60 | | 2 | 90 | | | | | | | |
| | <i>Konosirus punctatus</i> | | | | | | | | | | | | | | | |
| | <i>Lagocephalus wheeleri</i> | | | | | | | | | | | | | | | |
| | <i>Lepidotrigla japonice</i> | | | | | | | | | | | | | | | |
| | <i>Leiognathus nuchalis</i> | | | | 1 | 20 | | 2 | 20 | | 2 | 20 | | 2 | 30 | |
| | <i>Lisa macrolepis</i> | | | | | | | | | | | | | | | |
| | <i>Monacanthus chinensis</i> | 1 | 60 | | 1 | 70 | | | | | | | | | | |
| | <i>Muraenesox cinereus</i> | | | | | | | | | | | | | | | |
| | <i>Minous monodactylus</i> | | | | | | | | | | 1 | 10 | | | | |
| | <i>Navodon tessellates</i> | | | | | | | | | | | | | | | |
| | <i>Nemipterus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Nibea acuta</i> | | | | | | | | | | | | | | | |
| | <i>Nibea albiflora</i> | | | | | | | | | | | | | | | |

Trawling Data in July 2019

| Group | Species Name | INA | | | | | | | | | | | | | | |
|-----------------|------------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | INA-1 | | | INA-2 | | | INA-3 | | | INA-4 | | | INA-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Oxyurichthys papuensis</i> | | | | | | | | | | | | | | | |
| | <i>Parachaeturichthys polynema</i> | | | | | | | | | | | | | | | |
| | <i>Paraplagusia japonica</i> | | | | | | | | | | | | | | | |
| | <i>Pennahia anea</i> | | | | 3 | 16 | | 21 | 130 | | | | | | | |
| | <i>Pennahia argentata</i> | | | | | | | | | | | | | | | |
| | <i>Platycephalus indicus</i> | | | | | | | | | | | | 2 | 50 | | |
| | <i>Plectorhinchus pictus</i> | | | | 1 | 50 | | | | | | | | | | |
| | <i>Plotosus lineatus</i> | 1 | 670 | | | | | | | | | | | | | |
| | <i>Polydactylus sextarius</i> | | | | | | | | | | | | | | | |
| | <i>Pseudorhombus arsius</i> | | | | 1 | 110 | | | | | | | | | | |
| | <i>Pseudorhombus cinnamomeus</i> | | | | | | | | | | 1 | 10 | | | | |
| | <i>Pseudorhombus oligodon</i> | | | | | | | | | | | | 1 | 20 | | |
| | <i>Saurida elongata</i> | | | | | | | | | | | | | | | |
| | Sciaenidae | | | | | | | | | | | | | | | |
| | <i>Scoliodon sorrakowah</i> | | | | | | | | | | | | | | | |
| | <i>Scorpaenopsis longispina</i> | | | | | | | | | | | | | | | |
| | <i>Secutor ruconius</i> | | | | | | | | | | | | | | | |
| | <i>Sebastes marmoratus</i> | 2 | 80 | | 12 | 460 | | 1 | 6 | | | | 1 | 70 | | |
| | <i>Siganus fuscescens</i> | | | | 1 | 50 | | | | | | | 1 | 60 | | |
| | <i>Sillago sihama</i> | | | | | | | | | | | | | | | |
| | <i>Solea ovata</i> | | | | 2 | 20 | | 2 | 10 | | | | 3 | 30 | | |
| | <i>Takifugu niphobles</i> | | | | | | | | | | | | | | | |
| | <i>Thryssa hamiltonii</i> | | | | | | | | | | | | | | | |
| | <i>Thryssa kammalensis</i> | | | | | | | | | | | | | | | |
| | <i>Trypauchen vagina</i> | | | | | | | | | | | | | | | |
| Prawn or shrimp | <i>Alpheus brevicristatus</i> | | | | | | | 1 | 4 | | | | | | | |
| | <i>Alpheus distinguendus</i> | | | | | | | | | | | | | | | |
| | <i>Exopalaemon carimicauda</i> | | | | | | | | | | | | | | | |
| | <i>Metapenaeopsis barbata</i> | | | | | | | | | | | | | | | |
| | <i>Metapenaeus affinis</i> | | | | | | | | | | | | 1 | 10 | | |
| | <i>Metapenaeus joyneri</i> | | | | | | | | | | | | 1 | 10 | | |
| | <i>Metapenaeus intermedius</i> | | | | | | | | | | | | | | | |
| | <i>Parapenaeopsis cultirostris</i> | | | | | | | | | | | | | | | |
| | <i>Parapenaeopsis cornutus</i> | | | | | | | | | | | | | | | |
| | <i>Parapenaeopsis hardwickii</i> | | | | | | | | | | | | | | | |
| | <i>Parapenaeopsis hungerfordii</i> | | | | | | | | | | | | | | | |
| | <i>Parapenaeopsis tenella</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus merguensis</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus penicillatus</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus semisulcatus</i> | | | | | | | | | | 1 | 20 | | | | |
| | <i>Solenocera crassicornis</i> | | | | | | | | | | | | | | | |
| | <i>Trachypenaeus curvirostris</i> | | | | | | | | | | | | | | | |
| Crab | <i>Charybdis affinis</i> | 4 | 90 | | 17 | 400 | | 16 | 270 | | 1 | 25 | | 10 | 260 | |
| | <i>Charybdis annulata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis bimaculata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis cruciata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis japonica</i> | | | | | | | 2 | 140 | | | | | | | |

Trawling Data in July 2019

| Group | Species Name | INA | | | | | | | | | | | | | | |
|---------------|-----------------------------------|-----------------|--------------------|--------------------|-----------------|--------------------|--------------------|-----------------|--------------------|--------------------|-----------------|--------------------|--------------------|-----------------|--------------------|--------------------|
| | | INA-1 | | | INA-2 | | | INA-3 | | | INA-4 | | | INA-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Charybdis obtusifrons</i> | 6 | 280 | | 7 | 260 | | 6 | 250 | | 1 | 50 | | 4 | 210 | |
| | <i>Charybdis truncata</i> | | | | | | | | | | | | | 1 | 8 | |
| | <i>Charybdis variegata</i> | 2 | 10 | | | | | | | | | | | | | |
| | <i>Charybdis spp.</i> | 20 | 190 | * | 20 | 180 | * | 30 | 210 | * | 20 | 180 | * | 20 | 180 | * |
| | <i>Doclea sp.</i> | | | | | | | | | | | | | | | |
| | <i>Doclea ovis</i> | | | | | | | | | | 1 | 52 | | | | |
| | <i>Dorippe facchino</i> | | | | 30 | 490 | * | 20 | 280 | * | | | | 50 | 790 | * |
| | <i>Dorippe polita</i> | | | | 6 | 63 | | | | | | | | | | |
| | <i>Eucrate crenata</i> | 1 | 10 | | | | | | | | | | | | | |
| | <i>Galene bispinosa</i> | | | | | | | | | | | | | | | |
| | Hermit crab | 40 | 130 | * | 40 | 120 | * | 50 | 140 | * | 20 | 76 | * | 50 | 130 | * |
| | <i>Leucosia craniolearis</i> | | | | | | | | | | | | | | | |
| | <i>Macromedaeus distinguendus</i> | | | | 1 | 7.2 | | | | | | | | | | |
| | <i>Maorophthalmus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Matuta planipes</i> | | | | | | | | | | | | | | | |
| | <i>Ocypode ceratophthalmus</i> | | | | | | | | | | | | | | | |
| | <i>Parapanope euagora</i> | | | | | | | 1 | 9.9 | | 2 | 22.4 | | 2 | 21.8 | |
| | <i>Parthenope validus</i> | 20 | 260 | * | 40 | 570 | * | 20 | 270 | * | 20 | 240 | * | 30 | 360 | * |
| | <i>Plummus hirtellus</i> | | | | 1 | 6.8 | | | | | | | | | | |
| | <i>Porcellanella picta</i> | | | | | | | 2 | 0.7 | | | | | | | |
| | <i>Portunus hastatooides</i> | | | | | | | | | | | | | 1 | 2 | |
| | <i>Portunus pelagicus</i> | | | | 4 | 330 | | 2 | 50 | | | | | 2 | 210 | |
| | <i>Portunus sanguinolentus</i> | | | | | | | | | | | | | | | |
| | <i>Portunus trituberculatus</i> | | | | | | | | | | | | | | | |
| | <i>Thalamita sima</i> | | | | 1 | 10 | | 2 | 10 | | | | | 2 | 20 | |
| | <i>Typhlocarcinus sp.</i> | | | | 1 | 5.8 | | | | | | | | | | |
| Mantis shrimp | <i>Anchisquilla fasciata</i> | | | | | | | | | | | | | | | |
| | <i>Dictyosquilla foveolata</i> | | | | | | | | | | | | | | | |
| | <i>Harpisquilla harpax</i> | | | | 1 | 40 | | | | | 1 | 10 | | | | |
| | <i>Oratosquilla interrupta</i> | | | | 1 | 9 | | | | | | | | | | |
| | <i>Oratosquilla oratoria</i> | | | | | | | | | | | | | | | |
| | <i>Oratosquilla kempfi</i> | | | | | | | | | | | | | | | |
| Gastropod | <i>Bursa rana</i> | 2 | 14.6 | | | | | | | | | | | | | |
| | <i>Calyptrea sp.</i> | 6 | 2.1 | | 20 | 11 | * | | | | | | | 20 | 12 | * |
| | <i>Desmaulus extinctorius</i> | | | | | | | | | | 4 | 1.2 | | 20 | 16 | * |
| | <i>Drupa musiva</i> | | | | | | | | | | | | | | | |
| | <i>Epitonium scalare</i> | | | | | | | | | | | | | | | |
| | <i>Inquistor flavidula</i> | | | | | | | | | | | | | | | |
| | <i>Gemmula deshavesii</i> | | | | | | | | | | | | | | | |
| | <i>Lophiotoma leucotropis</i> | | | | 1 | 6.2 | | | | | | | | | | |
| | <i>Murex trapa</i> | | | | | | | | | | | | | | | |
| | <i>Nassarius hepaticus</i> | | | | | | | | | | | | | | | |
| | <i>Nassarius succinctus</i> | | | | | | | | | | 2 | 4.2 | | | | |
| | <i>Philine sp.</i> | | | | | | | | | | | | | | | |
| | <i>Pilosabia pilosa</i> | | | | | | | | | | | | | | | |
| | <i>Siphonotella walshi</i> | 3 | 1.1 | | | | | 20 | 10 | * | 6 | 2.1 | | 1 | 2.3 | |
| | <i>Tonna sulcosa</i> | | | | | | | | | | | | | | | |
| | <i>Turritella terebra</i> | 70 | 610 | * | 70 | 610 | * | 90 | 810 | * | 110 | 1030 | * | 80 | 720 | * |

Trawling Data in July 2019

| Group | Species Name | INA | | | | | | | | | | | | | | |
|------------|------------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | INA-1 | | | INA-2 | | | INA-3 | | | INA-4 | | | INA-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>UNID Nudibranch</i> | | | | | | | | | | | | | | | |
| Bivalve | <i>Anomia chinensis</i> | | | | | | | | | | | | | | | |
| | <i>Anomalodiscus squamosus</i> | | | | | | | | | | | | | | | |
| | <i>Chione isabellina</i> | | | | | | | | | | | | | | | |
| | <i>Chlamys nobilis</i> | | | | | | | | | | | | | | | |
| | <i>Macoma candida</i> | | | | | | | | | | | | | | | |
| | <i>Ostrea sp.</i> | | | | 20 | 44 | * | 80 | 120 | * | | | | | | |
| | <i>Paphia undulata</i> | | | | | | | | | | | | | | | |
| | <i>Parahyatissa sp.</i> | | | | 2 | 58 | | | | | | | | | | |
| | <i>Pinctada martensii</i> | | | | 3 | 160 | | 1 | 54 | | | | | 2 | 140 | |
| | <i>Placuna placenta</i> | | | | | | | | | | | | | | | |
| | <i>Ruditapes philippinarum</i> | 20 | 85 | * | 2 | 8.2 | | 20 | 190 | * | | | | | | |
| | <i>Samacar sp.</i> | | | | 1 | 19.6 | | | | | | | | | | |
| | <i>Scapharca subcrenata</i> | | | | | | | | | | | | | | | |
| | <i>Tegillarca nodifera</i> | | | | | | | | | | | | | | | |
| | <i>Tolonostrea talonata</i> | | | | | | | | | | | | | | | |
| | <i>Vepricardium asiaticum</i> | | | | | | | | | | | | | | | |
| Cephalopod | <i>Lolingid squid</i> | | | | | | | | | | | | | | | |
| | <i>Amphioctopus fangsiao</i> | | | | | | | | | | | | | | | |
| | <i>Sepiid cuttlefish</i> | | | | | | | | | | | | | | | |
| Echinoderm | <i>Acaudina molpadioides</i> | | | | | | | | | | | | | | | |
| | <i>Colochirus quadrangularis</i> | | | | 2 | 46 | | 1 | 18.4 | | | | | | | |
| | <i>Protankyra bidentata (c.f.)</i> | | | | 1 | 33 | | | | | | | | | | |
| | <i>Styela plicata</i> | | | | 3 | 11 | | 20 | 170 | * | 2 | 8.1 | | 20 | 230 | * |
| | <i>Temnopleurus toreumaticus</i> | 60 | 770 | * | 20 | 270 | * | 50 | 640 | * | 40 | 460 | * | 30 | 360 | * |
| Cnidaria | <i>Carcinactis ichikawai</i> | | | | | | | | | | | | | | | |
| | <i>Cavernularia habereri</i> | | | | | | | 2 | 7.4 | | | | | 1 | 8.4 | |
| | <i>Comanthus japonicus</i> | | | | 1 | 37 | | | | | | | | | | |
| | <i>Edwardsia japonica (c.f.)</i> | | | | | | | | | | | | | | | |
| | <i>Ellisella laevis</i> | 2 | 16.3 | | 1 | 16 | | 3 | 21 | | | | | 1 | 13 | |
| | Jelly fish | | | | | | | | | | | | | | | |
| | <i>Pteroeides chinense</i> | | | | 3 | 56 | | | | | 6 | 130 | | | | |
| | <i>Gorgonia spp.</i> | 2 | 38 | | 2 | 41 | | 3 | 238 | | 1 | 62 | | | | |
| | <i>Sea anomea</i> | | | | 2 | 18.1 | | | | | 3 | 21 | | | 1 | 6.1 |
| | <i>Virgularia gustaviana</i> | | | | 3 | 21.6 | | | | | | | | 1 | 12 | |
| Porifera | <i>Spongiatia</i> | 1 | 41 | | 1 | 32 | | | | | 1 | 13 | | | | |
| Cirripedia | <i>c.f. Bananus spp.</i> | 80 | 71 | * | 190 | 230 | * | 140 | 110 | * | 190 | 210 | * | 110 | 82 | * |

Trawling Data in July 2019

| Group | Species Name | INB-1 | | | INB-2 | | | INB | | | INB-4 | | | INB-5 | | |
|-------|--------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| Fish | <i>Arnoglossus waitei</i> | | | | | | | | | | | | | | | |
| | <i>Acentrogobius caninus</i> | | | | | | | | | | | | | | 1 | 20 |
| | <i>Anchoviella commersonii</i> | | | | | | | | | | | | | | | |
| | <i>Apogon kiensis</i> | | | | 1 | 3 | | 5 | 20 | | 1 | 5 | | 7 | 20 | |
| | <i>Apogon lineatus</i> | | | | | | | | | | | | | | | |
| | <i>Arius arius</i> | | | | | | | | | | | | | | | |
| | <i>Callionymus richardsoni</i> | | | | | | | | | | | | | | | |
| | <i>Callionymus curvicornis</i> | | | | | | | | | | | | | 1 | 10 | |
| | <i>Caranx kalla</i> | | | | | | | 1 | 4 | | | | | | | |
| | <i>Clupanodon thrissa</i> | | | | | | | | | | | | | | | |
| | <i>Collichthys lucidus</i> | | | | | | | | | | | | | | | |
| | <i>Coilia mystus</i> | | | | | | | | | | | | | | | |
| | <i>Crossorhombus kanekonis</i> | | | | | | | | | | | | | | | |
| | <i>Cryptocentrus filifer</i> | | | | 1 | 10 | | | | | | | | | | |
| | <i>Cynoglossus abbreviatus</i> | | | | | | | | | | | | | | | |
| | <i>Cynoglossus arel</i> | | | | | | | | | | 1 | 30 | | 1 | 40 | |
| | <i>Cynoglossus puncticeps</i> | | | | | | | | | | 1 | 10 | | 2 | 30 | |
| | <i>Cynoglossus roulei</i> | | | | | | | | | | | | | | | |
| | <i>Dasvatis akajei</i> | | | | | | | | | | | | | | | |
| | <i>Dasvatis zugei</i> | 3 | 260 | | | | | 2 | 160 | | 4 | 820 | | | | |
| | <i>Drepane punctata</i> | | | | | | | | | | | | | | | |
| | <i>Dendrophysa russelii</i> | 1 | 10 | | | | | | | | | | | | | |
| | <i>Epinephelus awoara</i> | | | | | | | 1 | 10 | | | | | | | |
| | <i>Epinephelus bruneus</i> | | | | | | | | | | | | | | | |
| | <i>Erisphex pottii</i> | 1 | 3 | | | | | | | | | | | 1 | 8 | |
| | <i>Gerres lucidus</i> | | | | | | | | | | | | | | | |
| | <i>Grammolites scaber</i> | | | | | | | | | | | | | | | |
| | <i>Gymnura japonica</i> | | | | | | | | | | | | | | | |
| | <i>Ilisha elongata</i> | | | | | | | | | | | | | | | |
| | <i>Inegocia japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Inimicus didactylus</i> | | | | 1 | 80 | | 1 | 10 | | | | | | | |
| | <i>Inimicus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Johnius belangerii</i> | | | | 3 | 70 | | 1 | 10 | | | | | 2 | 11 | |
| | <i>Johnius carona</i> | | | | | | | | | | | | | | | |
| | <i>Johnius poecilnotus</i> | | | | | | | | | | | | | | | |
| | <i>Johnius taiwanense</i> | | | | | | | | | | | | | | | |
| | <i>Konosirus punctatus</i> | | | | | | | | | | | | | | | |
| | <i>Lagocephalus wheeleri</i> | | | | | | | | | | | | | | | |
| | <i>Lepidotrigla japonice</i> | | | | | | | | | | | | | | | |
| | <i>Leiognathus nuchalis</i> | | | | | | | | | | | | | | | |
| | <i>Lisa macrolepis</i> | | | | | | | | | | | | | | | |
| | <i>Monacanthus chinensis</i> | | | | | | | | | | | | | | | |
| | <i>Muraenesox cinereus</i> | | | | | | | | | | | | | 1 | 100 | |
| | <i>Minous monodactylus</i> | | | | | | | | | | | | | | | |
| | <i>Navodon tessellates</i> | | | | | | | | | | | | | | | |
| | <i>Nemipterus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Nibea acuta</i> | | | | | | | | | | | | | | | |
| | <i>Nibea albiflora</i> | | | | | | | | | | | | | | | |

Trawling Data in July 2019

| Group | Species Name | INB | | | | | | | | | | | | | | |
|-----------------|------------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | INB-1 | | | INB-2 | | | INB-3 | | | INB-4 | | | INB-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Oxyurichthys papuensis</i> | | | | | | | | | | | | | | | |
| | <i>Parachaeturichthys polynema</i> | | | | | | | | | | | | | | | |
| | <i>Paraplagusia japonica</i> | | | | | | | | | | | | | | | |
| | <i>Pennahia anea</i> | | | | | | | | | | 1 | 8 | | 5 | 30 | |
| | <i>Pennahia argentata</i> | 1 | 4 | | 4 | 10 | | | | | 2 | 8 | | | | |
| | <i>Platycephalus indicus</i> | 1 | 80 | | | | | | | | | | | | | |
| | <i>Plectorhinchus pictus</i> | | | | | | | | | | | | | | | |
| | <i>Plotosus lineatus</i> | | | | | | | | | | | | | | | |
| | <i>Polydactylus sextarius</i> | | | | | | | | | | | | | | | |
| | <i>Pseudorhombus arsius</i> | | | | | | | | | | | | | | | |
| | <i>Pseudorhombus cinnamomeus</i> | | | | | | | | | | | | | | | |
| | <i>Pseudorhombus oligodon</i> | | | | | | | | | | | | | 1 | 10 | |
| | <i>Saurida elongata</i> | | | | | | | | | | | | | | | |
| | Sciaenidae | | | | | | | | | | | | | | | |
| | <i>Scoliodon sorrakowah</i> | | | | | | | | | | | | | | | |
| | <i>Scorpaenopsis longispina</i> | | | | | | | | | | | | | | | |
| | <i>Secutor ruconius</i> | | | | | | | | | | | | | | | |
| | <i>Sebastes marmoratus</i> | 1 | 50 | | | | | | | | | | | | | |
| | <i>Siganus fuscescens</i> | | | | | | | | | | | | | | | |
| | <i>Sillago sihama</i> | | | | | | | | | | | | | | | |
| | <i>Solea ovata</i> | | | | 2 | 20 | | | | | | | | 1 | 4 | |
| | <i>Takifugu niphobles</i> | | | | | | | | | | | | | | | |
| | <i>Thryssa hamiltonii</i> | | | | | | | | | | | | | | | |
| | <i>Thryssa kammalensis</i> | | | | | | | | | | | | | | | |
| | <i>Trypauchen vagina</i> | | | | | | | | | | | | | | | |
| Prawn or shrimp | <i>Alpheus brevicristatus</i> | | | | | | | | | | | | | 1 | 1 | |
| | <i>Alpheus distinguendus</i> | | | | | | | | | | | | | 1 | 2 | |
| | <i>Exopalaemon carinicauda</i> | | | | | | | | | | | | | | | |
| | <i>Metapenaeopsis barbata</i> | | | | | | | 1 | 3 | | | | | 1 | 3 | |
| | <i>Metapenaeus affinis</i> | | | | 23 | 480 | | 14 | 300 | | | | | 1 | 6 | |
| | <i>Metapenaeus joyneri</i> | | | | | | | | | | | | | | | |
| | <i>Metapenaeus intermedius</i> | | | | | | | | | | | | | | | |
| | <i>Parapenaeopsis cultirostris</i> | | | | | | | | | | | | | | | |
| | <i>Parapenaeopsis cornutus</i> | | | | | | | | | | | | | | | |
| | <i>Parapenaeopsis hardwickii</i> | | | | | | | | | | | | | | | |
| | <i>Parapenaeopsis hungerfordii</i> | | | | | | | | | | | | | | | |
| | <i>Parapenaeopsis tenella</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus merguensis</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus penicillatus</i> | 2 | 10 | | | | | | | | | | | 2 | 20 | |
| | <i>Penaeus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus semisulcatus</i> | | | | | | | 1 | 10 | | | | | 1 | 10 | |
| | <i>Solenocera crassicornis</i> | | | | | | | | | | | | | | | |
| | <i>Trachypenaeus curvirostris</i> | | | | | | | | | | | | | | | |
| Crab | <i>Charybdis affinis</i> | 4 | 70 | | | | | | | | | | | 15 | 230 | |
| | <i>Charybdis annulata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis bimaculata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis cruciata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis japonica</i> | | | | 1 | 10 | | | | | | | | | | |

Trawling Data in July 2019

| Group | Species Name | INB | | | | | | | | | | | | | | |
|---------------|-----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | INB-1 | | | INB-2 | | | INB-3 | | | INB-4 | | | INB-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Charybdis obtusifrons</i> | | | | 1 | 20 | | 2 | 100 | | | | | 4 | 150 | |
| | <i>Charybdis truncata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis variegata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis spp.</i> | 30 | 460 | * | 30 | 470 | * | 40 | 590 | * | 40 | 580 | * | 40 | 560 | * |
| | <i>Doclea sp.</i> | | | | | | | | | | | | | | | |
| | <i>Doclea ovis</i> | | | | | | | | | | | | | | | |
| | <i>Dorippe facchino</i> | | | | 20 | 240 | * | | | | | | | | | |
| | <i>Dorippe polita</i> | | | | | | | | | | | | | | | |
| | <i>Eucrate crenata</i> | | | | | | | | | | | | | 1 | 8.6 | |
| | <i>Galene bispinosa</i> | | | | | | | | | | 2 | 84 | | | | |
| | Hermit crab | 40 | 210 | * | 40 | 230 | * | 40 | 220 | * | 30 | 50 | * | 30 | 50 | * |
| | <i>Leucosia cranioilaris</i> | 20 | 102 | * | | | | 5 | 21 | | | | | 2 | 8.1 | |
| | <i>Macromedaeus distinguendus</i> | | | | | | | | | | | | | | | |
| | <i>Maorophthalmus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Matuta planipes</i> | | | | | | | | | | | | | | | |
| | <i>Ocyrope ceratophthalmus</i> | | | | | | | | | | | | | | | |
| | <i>Parapanope euagora</i> | | | | | | | | | | | | | | | |
| | <i>Parthenope validus</i> | 16 | 174 | | 5 | 63 | | 5 | 63 | | | | | | | |
| | <i>Plummus hirtellus</i> | | | | | | | | | | | | | | | |
| | <i>Porcellanella picta</i> | 4 | 1.6 | | | | | 2 | 0.8 | | | | | 1 | 0.4 | |
| | <i>Portunus hastatooides</i> | | | | | | | | | | | | | | | |
| | <i>Portunus pelagicus</i> | | | | | | | | | | | | | | | |
| | <i>Portunus sanguinolentus</i> | | | | | | | | | | | | | | | |
| | <i>Portunus trituberculatus</i> | | | | | | | | | | | | | | | |
| | <i>Thalamita sima</i> | | | | | | | | | | | | | | | |
| | <i>Typhlocarcinus sp.</i> | | | | | | | | | | | | | | | |
| Mantis shrimp | <i>Anchisquilla fasciata</i> | | | | | | | | | | | | | | | |
| | <i>Dictyosquilla foveolata</i> | | | | | | | | | | | | | | | |
| | <i>Harpisquilla harpax</i> | | | | | | | 1 | 40 | | | | | 1 | 50 | |
| | <i>Oratosquilla interrupta</i> | | | | | | | | | | | | | 4 | 70 | |
| | <i>Oratosquilla oratoria</i> | | | | | | | | | | | | | | | |
| | <i>Oratosquilla kempii</i> | | | | | | | | | | | | | | | |
| Gastropod | <i>Bursa rana</i> | | | | | | | | | | | | | | | |
| | <i>Calyptrea sp.</i> | 2 | 0.6 | | 20 | 12 | * | | | | 2 | 0.9 | | 1 | 0.4 | |
| | <i>Desmaulus extintorius</i> | | | | | | | | | | | | | | | |
| | <i>Drupa musiva</i> | 1 | 4.6 | | | | | | | | | | | | | |
| | <i>Epitonium scalare</i> | | | | | | | | | | | | | | | |
| | <i>Inquistor flavidula</i> | | | | | | | 4 | 5.6 | | | | | | | |
| | <i>Gemmula deshavesii</i> | | | | | | | | | | | | | | | |
| | <i>Lophiotoma leucotropis</i> | | | | 1 | 4.6 | | | | | | | | | | |
| | <i>Murex trapa</i> | 40 | 460 | * | 3 | 11.2 | | 20 | 110 | * | | | | | | |
| | <i>Nassarius hepaticus</i> | | | | | | | | | | | | | | | |
| | <i>Nassarius succinctus</i> | | | | | | | | | | | | | | | |
| | <i>Philine sp.</i> | | | | 2 | 11.8 | | 3 | 18.6 | | 1 | 8.4 | | | | |
| | <i>Pilosabia pilosa</i> | | | | | | | | | | | | | | | |
| | <i>Siphopatella walshi</i> | | | 2 | 0.8 | | | | | | | | | | | |
| | <i>Tonna sulcosa</i> | 2 | 270 | | | | | | | | | | | | | |
| | <i>Turritella terebra</i> | 540 | 5160 | * | 210 | 1940 | * | 190 | 1730 | * | 90 | 780 | * | 90 | 780 | * |

Trawling Data in July 2019

| Group | Species Name | INB-1 | | | INB-2 | | | INB | | | INB-4 | | | INB-5 | | |
|------------|------------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>UNID Nudibranch</i> | | | | | | | | | | | | | | | |
| Bivalve | <i>Anomia chinensis</i> | | | | 3 | 18.6 | | 1 | 6.2 | | | | | | | |
| | <i>Anomalodiscus squamosus</i> | | | | | | | | | | | | | | | |
| | <i>Chione isabellina</i> | | | | | | | | | | | | | | | |
| | <i>Chlamys nobilis</i> | | | | | | | | | | | | | | | |
| | <i>Macoma candida</i> | | | | | | | | | | | | | | | |
| | <i>Ostrea sp.</i> | 40 | 150 | * | 50 | 180 | * | 30 | 112 | * | | | | | | |
| | <i>Paphia undulata</i> | | | | | | | | | | | | | | | |
| | <i>Parahyattissa sp.</i> | | | | | | | | | | | | | | | |
| | <i>Pinctada martensii</i> | 1 | 26 | | | | | | | | | | | | | |
| | <i>Placuna placenta</i> | | | | | | | | | | | | | | | |
| | <i>Ruditapes philippinarum</i> | 20 | 120 | * | | | | | | | 20 | 140 | * | | | |
| | <i>Samacar sp.</i> | | | | | | | | | | | | | | | |
| | <i>Scapharca subcrenata</i> | | | | | | | | | | | | | | | |
| | <i>Tegillarca nodifera</i> | | | | | | | | | | | | | | | |
| | <i>Tolonostrea talonata</i> | 3 | 10.2 | | | | | | | | | | | | | |
| | <i>Vepricardium asiaticum</i> | | | | | | | | | | | | | | | |
| Cephalopod | <i>Lolingid squid</i> | | | | | | | | | | | | | | | |
| | <i>Amphioctopus fangstao</i> | 1 | 30 | | 1 | 80 | | | | | | | | | | |
| | <i>Sepiid cuttlefish</i> | | | | | | | | | | | | | | 1 | 80 |
| Echinoderm | <i>Acaudina molpadioides</i> | 20 | 960 | * | | | | 20 | 690 | * | | | | | 2 | 28 |
| | <i>Colochirus quadrangularis</i> | | | | | | | | | | | | | | | |
| | <i>Protankyra bidentata (c.f.)</i> | | | | | | | | | | | | | | | |
| | <i>Styela plicata</i> | 4 | 28 | | 2 | 11 | | 1 | 5.4 | | 1 | 4.8 | | 2 | 10.4 | |
| | <i>Temnopleurus toreumaticus</i> | 30 | 340 | * | 40 | 620 | * | 30 | 350 | * | 40 | 490 | * | 30 | 420 | * |
| Cnidaria | <i>Carcinactis ichikawai</i> | | | | | | | | | | | | | | | |
| | <i>Cavernularia habereri</i> | 30 | 270 | * | 2 | 6.4 | | 20 | 170 | * | 5 | 21 | | 20 | 160 | * |
| | <i>Comanthus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Edwardsia japonica (c.f.)</i> | 8 | 96 | | | | | | | | | | | | | |
| | <i>Ellisella laevis</i> | 5 | 64 | | 15 | 210 | | 7 | 76 | | | | | 1 | 6.5 | |
| | Jelly fish | | | | 1 | 3100 | | | | | | | | | | |
| | <i>Pteroeides chinense</i> | 15 | 940 | | 8 | 184 | | 14 | 294 | | 3 | 64 | | 10 | 290 | |
| | <i>Gorgonia spp.</i> | | | | | | | 1 | 11 | | 1 | 11.8 | | | | |
| | Sea anomea | 3 | 12 | | 4 | 26 | | 2 | 11 | | | | | | | |
| | <i>Virgularia gustaviana</i> | 6 | 74 | | 1 | 21 | | | | | 1 | 12 | | 1 | 16 | |
| Porifera | Spongiatia | | | | | | | | | | | | | | | |
| Cirripedia | <i>c.f. Bananus spp.</i> | 160 | 94 | * | 320 | 240 | * | 90 | 58 | * | 60 | 52 | * | 70 | 63 | * |

Trawling Data in July 2019

| Group | Species Name | TNA | | | | | | | | | | | | | | |
|-------|--------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | TNA-1 | | | TNA-2 | | | TNA-3 | | | TNA-4 | | | TNA-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| Fish | <i>Arnoglossus waitei</i> | | | | 1 | 10 | | 2 | 20 | | 1 | 10 | | | | |
| | <i>Acentrogobius caninus</i> | 1 | 20 | | | | | | | | | | | | | |
| | <i>Anchoviella commersonii</i> | | | | | | | | | | | | | | | |
| | <i>Apogon kiensis</i> | 1 | 5 | | | | | 1 | 10 | | | | | | | |
| | <i>Apogon lineatus</i> | 1 | 10 | | | | | | | | | | | | | |
| | <i>Arius arius</i> | | | | | | | | | | | | | | | |
| | <i>Callionymus richardsoni</i> | | | | | | | | | | | | | | | |
| | <i>Callionymus curvicornis</i> | | | | | | | | | | | | | | | |
| | <i>Caranx kalla</i> | | | | | | | | | | | | | | | |
| | <i>Clupanodon thrissa</i> | | | | 1 | 30 | | | | | | | | | | |
| | <i>Collichthys lucidus</i> | 12 | 80 | | 16 | 110 | | 10 | 60 | | 2 | 20 | | 2 | 10 | |
| | <i>Coilia mystus</i> | | | | 1 | 20 | | | | | | | | | | |
| | <i>Crossorhombus kanekonis</i> | 1 | 10 | | | | | | | | | | | | | |
| | <i>Cryptocentrus filifer</i> | | | | | | | | | | | | | | | |
| | <i>Cynoglossus abbreviatus</i> | | | | | | | | | | | | | | | |
| | <i>Cynoglossus arel</i> | 3 | 90 | | 3 | 70 | | 1 | 20 | | 2 | 80 | | 4 | 50 | |
| | <i>Cynoglossus puncticeps</i> | | | | | | | 1 | 10 | | 1 | 10 | | | | |
| | <i>Cynoglossus roulei</i> | | | | | | | | | | | | | | | |
| | <i>Dasvatis akajei</i> | | | | | | | | | | | | | | | |
| | <i>Dasvatis zugei</i> | | | | | | | | | | 4 | 680 | | 3 | 390 | |
| | <i>Drepane punctata</i> | | | | | | | | | | | | | | | |
| | <i>Dendrophysa russelii</i> | 3 | 60 | | 5 | 110 | | 37 | 960 | | | | | 2 | 80 | |
| | <i>Epinephelus awoara</i> | | | | | | | | | | | | | | | |
| | <i>Epinephelus bruneus</i> | | | | | | | | | | | | | | | |
| | <i>Erispex pottii</i> | 1 | 10 | | | | | 5 | 20 | | 2 | 10 | | 1 | 5 | |
| | <i>Gerres lucidus</i> | | | | | | | | | | | | | | | |
| | <i>Grammolites scaber</i> | | | | | | | 1 | 40 | | | | | | | |
| | <i>Gymnura japonica</i> | | | | | | | | | | | | | 2 | 760 | |
| | <i>Ilisha elongata</i> | | | | | | | | | | | | | | | |
| | <i>Inegocia japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Inimicus didactylus</i> | | | | | | | | | | | | | | | |
| | <i>Inimicus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Johnius belangerii</i> | | | | | | | | | | | | | | | |
| | <i>Johnius carona</i> | | | | | | | 9 | 160 | | 7 | 60 | | | | |
| | <i>Johnius poecilonotus</i> | | | | | | | | | | | | | | | |
| | <i>Johnius taiwanense</i> | 3 | 30 | | 3 | 60 | | 3 | 60 | | | | | | | |
| | <i>Konosirus punctatus</i> | | | | | | | | | | | | | | | |
| | <i>Lagocephalus wheeleri</i> | | | | | | | | | | | | | | | |
| | <i>Lepidotrigla japonice</i> | 1 | 10 | | | | | 1 | 10 | | | | | | | |
| | <i>Leiognathus nuchalis</i> | | | | | | | 1 | 10 | | | | | 2 | 20 | |
| | <i>Lisa macrolepis</i> | | | | | | | | | | | | | | | |
| | <i>Monacanthus chinensis</i> | | | | | | | | | | | | | | | |
| | <i>Muraenesox cinereus</i> | | | | | | | 1 | 710 | | | | | 1 | 100 | |
| | <i>Minous monodactylus</i> | | | | | | | | | | | | | | | |
| | <i>Navodon tessellates</i> | | | | | | | | | | | | | | | |
| | <i>Nemipterus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Nibea acuta</i> | | | | | | | | | | | | | | | |
| | <i>Nibea albiflora</i> | | | | | | | | | | | | | | | |

Trawling Data in July 2019

| Group | Species Name | TNA | | | | | | | | | | | | | | |
|-----------------|------------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | TNA-1 | | | TNA-2 | | | TNA-3 | | | TNA-4 | | | TNA-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Oxyurichthys papuensis</i> | | | | | | | | | | | | | | | |
| | <i>Parachaeturichthys polynema</i> | 1 | 10 | | | | | | | | | | | | | |
| | <i>Paraplagusia japonica</i> | | | | | | | | | | | | | | | |
| | <i>Pennahia anea</i> | | | | 1 | 7 | | | | | | | | | | |
| | <i>Pennahia argentata</i> | 2 | 10 | | 4 | 30 | | 14 | 140 | | | | 2 | 10 | | |
| | <i>Platycephalus indicus</i> | 1 | 10 | | | | | | | | | | | | | |
| | <i>Plectorhinchus pictus</i> | | | | | | | | | | | | | | | |
| | <i>Plotosus lineatus</i> | | | | | | | | | | | | | | | |
| | <i>Polydactylus sextarius</i> | | | | | | | | | | 2 | 10 | | 1 | 5 | |
| | <i>Pseudorhombus arsius</i> | | | | | | | | | | | | | | | |
| | <i>Pseudorhombus cinnamomeus</i> | | | | | | | 1 | 50 | | | | | | | |
| | <i>Pseudorhombus oligodon</i> | | | | | | | 2 | 40 | | | | | | | |
| | <i>Saurida elongata</i> | | | | | | | | | | | | | | | |
| | <i>Sciaenidae</i> | | | | | | | | | | 4 | 60 | | | | |
| | <i>Scoliodon sorrakowah</i> | | | | 1 | 20 | | 9 | 180 | | | | 1 | 20 | | |
| | <i>Scorpaenopsis longispina</i> | 1 | 10 | | 2 | 10 | | 1 | 10 | | 1 | 5 | | 3 | 10 | |
| | <i>Secutor ruconius</i> | | | | | | | | | | | | | | | |
| | <i>Sebastiscus marmoratus</i> | | | | | | | | | | | | | | | |
| | <i>Siganus fuscescens</i> | | | | | | | | | | | | | | | |
| | <i>Sillago sihama</i> | | | | | | | | | | | | | | | |
| | <i>Solea ovata</i> | | | | 1 | 5 | | 1 | 5 | | 1 | 10 | | 2 | 10 | |
| | <i>Takifugu niphobles</i> | | | | | | | | | | | | | | | |
| | <i>Thryssa hamiltonii</i> | | | | | | | | | | | | | | | |
| | <i>Thryssa kammalensis</i> | | | | | | | | | | | | | | | |
| | <i>Trypauchen vagina</i> | 5 | 70 | | 18 | 250 | | 31 | 400 | | 2 | 20 | | 8 | 110 | |
| Prawn or shrimp | <i>Alpheus brevicristatus</i> | | | | | | | | | | 1 | 2 | | | | |
| | <i>Alpheus distinguendus</i> | | | | | | | 1 | 3 | | | | | | | |
| | <i>Exopalaemon carinicauda</i> | | | | | | | 2 | 10 | | | | | | | |
| | <i>Metapenaeopsis barbata</i> | 1 | 10 | | 4 | 10 | | 3 | 10 | | | | 4 | 16 | | |
| | <i>Metapenaeus affinis</i> | 287 | 1380 | | 385 | 2470 | | 621 | 3560 | | 139 | 900 | | 114 | 820 | |
| | <i>Metapenaeus joyneri</i> | | | | 1 | 10 | | | | | | | | | | |
| | <i>Metapenaeus intermedius</i> | 2 | 20 | | 7 | 30 | | 2 | 10 | | 6 | 30 | | 2 | 10 | |
| | <i>Parapenaeopsis cultirostris</i> | | | | 1 | 10 | | 3 | 10 | | 1 | 8 | | 2 | 10 | |
| | <i>Parapenaeopsis cornutus</i> | 1 | 8 | | | | | | | | | | | | | |
| | <i>Parapenaeopsis hardwickii</i> | 5 | 60 | | 2 | 20 | | 2 | 10 | | 3 | 20 | | 4 | 50 | |
| | <i>Parapenaeopsis hungerfordii</i> | 5 | 30 | | 19 | 90 | | 25 | 120 | | 4 | 30 | | 7 | 40 | |
| | <i>Parapenaeopsis tenella</i> | | | | | | | | | | 1 | 3 | | | | |
| | <i>Penaeus merguensis</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus penicillatus</i> | 4 | 60 | | 17 | 170 | | 6 | 60 | | 10 | 80 | | 16 | 190 | |
| | <i>Penaeus japonicus</i> | | | | 1 | 10 | | 1 | 10 | | | | | | | |
| | <i>Penaeus semisulcatus</i> | 2 | 20 | | 1 | 10 | | | | | | | 3 | 20 | | |
| | <i>Solenocera crassicornis</i> | | | | 1 | 5 | | 4 | 15 | | | | | 1 | 8 | |
| | <i>Trachypenaeus curvirostris</i> | 5 | 20 | | 21 | 70 | | 37 | 130 | | 23 | 80 | | 31 | 90 | |
| Crab | <i>Charybdis affinis</i> | 63 | 1370 | | 101 | 1550 | | 132 | 2010 | | 32 | 450 | | 9 | 140 | |
| | <i>Charybdis annulata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis bimaculata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis cruciata</i> | 1 | 40 | | 2 | 50 | | | | | | | | | | |
| | <i>Charybdis japonica</i> | | | | | | | | | | | | | | | |

Trawling Data in July 2019

| Group | Species Name | TNA | | | | | | | | | | | | | | |
|---------------|----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | TNA-1 | | | TNA-2 | | | TNA-3 | | | TNA-4 | | | TNA-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Charybdis obtusifrons</i> | | | | 5 | 210 | | 9 | 460 | | 2 | 80 | | 2 | 140 | |
| | <i>Charybdis truncata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis variegata</i> | 1 | 10 | | 2 | 10 | | 4 | 20 | | 1 | 10 | | | | |
| | <i>Charybdis spp.</i> | 30 | 410 | * | 30 | 410 | * | 20 | 370 | * | 20 | 360 | * | 30 | 410 | * |
| | <i>Doclea sp.</i> | | | | | | | | | | 20 | 310 | * | | | |
| | <i>Doclea ovis</i> | | | | | | | | | | | | | | | |
| | <i>Dorippe facchino</i> | | | | | | | | | | | | | | | |
| | <i>Dorippe polita</i> | | | | 7 | 56 | | | | | 6 | 62 | | | | |
| | <i>Eucreta crenata</i> | | | | 2 | 9.8 | | 1 | 5.2 | | 3 | 12.8 | | 2 | 12.4 | |
| | <i>Galene bispinosa</i> | | | | | | | | | | | | | | | |
| | Hermit crab | | | | 50 | 260 | * | | | | | | | | | |
| | <i>Leucosia craniolaris</i> | | | | | | | | | | | | | | | |
| | <i>Macromedaeus distinguadus</i> | | | | | | | | | | | | | | | |
| | <i>Maorophthalmus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Matuta planipes</i> | | | | | | | | | | | | | | | |
| | <i>Ocypode ceratophthalmus</i> | | | | | | | | | | | | | | | |
| | <i>Parapanope euagora</i> | | | | | | | | | | | | | | | |
| | <i>Parthenope validus</i> | 5 | 110 | | 20 | 230 | * | 30 | 360 | * | 20 | 230 | * | 3 | 54 | |
| | <i>Plumnus hirtellus</i> | | | | | | | | | | | | | | | |
| | <i>Porcellanella picta</i> | 4 | 1.6 | | 5 | 1.8 | | 4 | 1.5 | | 2 | 0.9 | | 1 | 0.5 | |
| | <i>Portunus hastatoides</i> | 1 | 5 | | 5 | 30 | | 12 | 40 | | | | | 2 | 10 | |
| | <i>Portunus pelagicus</i> | 1 | 10 | | 3 | 110 | | 1 | 15 | | 1 | 80 | | | | |
| | <i>Portunus sanguinolentus</i> | 1 | 10 | | | | | 1 | 10 | | 1 | 110 | | 1 | 60 | |
| | <i>Portunus trituberculatus</i> | 1 | 60 | | | | | | | | | | | | | |
| | <i>Thalamita sima</i> | | | | | | | | | | | | | | | |
| | <i>Typhlocarcinus sp.</i> | | | | | | | | | | 1 | 3.8 | | | | |
| Mantis shrimp | <i>Anchisquilla fasciata</i> | | | | | | | | | | | | | | | |
| | <i>Dictyosquilla foveolata</i> | 3 | 40 | | 4 | 20 | | 2 | 40 | | 1 | 10 | | | | |
| | <i>Harpisquilla harpax</i> | | | | | | | | | | | | | | | |
| | <i>Oratosquilla interrupta</i> | 27 | 530 | | 34 | 630 | | 42 | 870 | | 15 | 310 | | 13 | 280 | |
| | <i>Oratosquilla oratoria</i> | 6 | 70 | | 14 | 190 | | 13 | 140 | | 10 | 130 | | 16 | 160 | |
| | <i>Oratosquilla kempfi</i> | | | | | | | 1 | 20 | | | | | | | |
| Gastropod | <i>Bursa rana</i> | | | | | | | | | | | | | | | |
| | <i>Calyptreaea sp.</i> | 20 | 12 | * | 20 | 12 | * | 20 | 12 | * | 30 | 19 | * | 20 | 13 | * |
| | <i>Desmaulus extinctorius</i> | | | | | | | | | | | | | | | |
| | <i>Drupa musiva</i> | 3 | 14 | | | | | | | | | | | | | |
| | <i>Epitonium scalare</i> | | | | | | | | | | | | | | | |
| | <i>Inquistor flavidula</i> | | | | 20 | 46 | * | | | | 40 | 92 | * | | | |
| | <i>Gemmula deshayesii</i> | | | | | | | 20 | 42 | * | | | | | | |
| | <i>Lophiotoma leucotropis</i> | | | | | | | | | | | | | | | |
| | <i>Murex trapa</i> | 40 | 510 | * | 60 | 770 | * | 30 | 320 | * | 60 | 790 | * | 50 | 610 | * |
| | <i>Nassarius hepaticus</i> | | | | | | | | | | 1 | 4.2 | | | | |
| | <i>Nassarius succinctus</i> | | | | 60 | 130 | * | | | | 60 | 130 | * | | | |
| | <i>Philine sp.</i> | | | | | | | | | | | | | | | |
| | <i>Pilosabia pilosa</i> | | | | 20 | 11 | * | 20 | 11 | * | 20 | 12 | * | 20 | 12 | * |
| | <i>Siphopatella walshi</i> | 20 | 32 | * | | | | | | | | | | | | |
| | <i>Tonna sulcosa</i> | | | | | | | | | | | | | | | |
| | <i>Turritella terebra</i> | 1400 | 13200 | * | 1500 | 14810 | * | 960 | 9230 | * | 920 | 9050 | * | 830 | 8090 | * |

Trawling Data in July 2019

| Group | Species Name | TNA | | | | | | | | | | | | | | |
|------------|------------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | TNA-1 | | | TNA-2 | | | TNA-3 | | | TNA-4 | | | TNA-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>UNID Nudibranch</i> | | | | | | | 2 | 6.4 | | | | | | | |
| Bivalve | <i>Anomia chinensis</i> | | | | | | | | | | | | | | | |
| | <i>Anomalodiscus squamosus</i> | | | | | | | | | | | | | | | |
| | <i>Chione isabellina</i> | | | | | | | | | | | | | | | |
| | <i>Chlamys nobilis</i> | | | | | | | | | | 2 | 21 | | | | |
| | <i>Macoma candida</i> | | | | | | | | | | | | | | | |
| | <i>Ostrea sp.</i> | | | | | | | | | | | | | | | |
| | <i>Paphia undulata</i> | | | | | | | | | | 5 | 32 | | | | |
| | <i>Parahyatisa sp.</i> | | | | | | | | | | | | | | | |
| | <i>Pinctada martensii</i> | | | | | | | | | | | | | | | |
| | <i>Placuna placenta</i> | | | | | | | | | | | | | | | |
| | <i>Ruditapes philippinarum</i> | | | | | | | | | | | | 2 | 10 | | |
| | <i>Samacar sp.</i> | | | | | | | | | | | | | | | |
| | <i>Scapharca subcrenata</i> | 2 | 93 | | 1 | 9 | | 2 | 92 | | 1 | 76 | | 3 | 188 | |
| | <i>Tegillarca nodifera</i> | | | | | | | | | | | | | | | |
| | <i>Tolonostrea talonata</i> | | | | | | | | | | | | | | | |
| | <i>Vepricardium asiaticum</i> | | | | | | | | | | | | | | | |
| Cephalopod | <i>Lolingid squid</i> | | | | 1 | 5 | | | | | | | | | | |
| | <i>Amphioctopus fangsiao</i> | | | | | | | | | | | | | | | |
| | <i>Sepiid cuttlefish</i> | 3 | 90 | | 1 | 20 | | 3 | 110 | | 1 | 30 | | | | |
| Echinoderm | <i>Acaudina molpadioides</i> | | | | 5 | 120 | | 6 | 160 | | | | | 2 | 76 | |
| | <i>Colochirus quadrangularis</i> | | | | | | | | | | | | | | | |
| | <i>Protankyra bidentata (c.f.)</i> | | | | | | | | | | | | | | | |
| | <i>Styela plicata</i> | 2 | 14 | | | | | | | | | | | 3 | 12 | |
| | <i>Temnopleurus toreumaticus</i> | 80 | 1120 | * | 190 | 3460 | * | 110 | 1880 | * | 90 | 1140 | * | 110 | 1840 | |
| Cnidaria | <i>Carcinactis ichikawai</i> | | | | | | | | | | 2 | 1.8 | | | | |
| | <i>Cavernularia habereri</i> | 50 | 410 | * | 70 | 630 | * | 510 | 4110 | * | 50 | 420 | * | 20 | 160 | |
| | <i>Comanthus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Edwardsia japonica (c.f.)</i> | | | | 4 | 26 | | 3 | 16 | | 2 | 16 | | | | |
| | <i>Ellisella laevis</i> | | | | | | | | | | | | | | | |
| | Jelly fish | | | | | | | | | | | | | | | |
| | <i>Pteroeides chinense</i> | 40 | 1630 | * | 40 | 1720 | * | 20 | 870 | * | 30 | 1580 | * | 20 | 780 | |
| | <i>Gorgonia spp.</i> | | | | | | | | | | | | | | | |
| | <i>Sea anomea</i> | 3 | 15 | | 5 | 32 | | 4 | 28 | | 8 | 90 | | 1 | 7 | |
| | <i>Virgularia gustaviana</i> | 4 | 84 | | 1 | 10 | | 2 | 32 | | 4 | 78 | | 1 | 10 | |
| Porifera | <i>Spongiatia</i> | | | | | | | | | | | | | | | |
| Cirripedia | <i>c.f. Bananus spp.</i> | 90 | 63 | * | 80 | 56 | * | 90 | 62 | * | 110 | 63 | * | 60 | 81 | |

Trawling Data in July 2019

| Group | Species Name | TNB | | | | | | | | | | | | | | |
|-------|--------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | TNB-1 | | | TNB-2 | | | TNB-3 | | | TNB-4 | | | TNB-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| Fish | <i>Arnoglossus waitei</i> | 2 | 20 | | 5 | 50 | | | | | 1 | 10 | | 1 | 10 | |
| | <i>Acentrogobius caninus</i> | | | | | | | | | | | | | | | |
| | <i>Anchoviella commersonii</i> | | | | | | | | | | | | | | | |
| | <i>Apogon kiensis</i> | 5 | 20 | | 13 | 60 | | 7 | 20 | | 12 | 50 | | 3 | 10 | |
| | <i>Apogon lineatus</i> | | | | | | | | | | | | | | | |
| | <i>Arius arius</i> | 1 | 30 | | | | | 3 | 120 | | | | | 1 | 280 | |
| | <i>Callionymus richardsoni</i> | | | | | | | | | | | | | | | |
| | <i>Callionymus curvicornis</i> | 2 | 20 | | | | | | | | | | | | | |
| | <i>Caranx kalla</i> | | | | | | | | | | | | | | | |
| | <i>Clupanodon thrissa</i> | 8 | 330 | | 1 | 70 | | 7 | 350 | | | | | | | |
| | <i>Collichthys lucidus</i> | 25 | 140 | | 7 | 30 | | 16 | 86 | | 6 | 30 | | 1 | 10 | |
| | <i>Coilia mystus</i> | | | | | | | | | | | | | | | |
| | <i>Crossorhombus kanekonis</i> | | | | | | | | | | | | | | | |
| | <i>Cryptocentrus filifer</i> | 1 | 10 | | | | | | | | | | | | | |
| | <i>Cynoglossus abbreviatus</i> | | | | | | | | | | | | | | | |
| | <i>Cynoglossus arel</i> | 2 | 50 | | 11 | 240 | | 5 | 140 | | 5 | 110 | | 5 | 90 | |
| | <i>Cynoglossus puncticeps</i> | | | | | | | | | | | | | | | |
| | <i>Cynoglossus roulei</i> | | | | | | | | | | 1 | 70 | | | | |
| | <i>Dasyatis akajei</i> | 1 | 2300 | | 2 | 1890 | | | | | | | | 1 | 150 | |
| | <i>Dasyatis zugei</i> | 14 | 1750 | | 12 | 1690 | | 34 | 4100 | | 14 | 2250 | | 3 | 70 | |
| | <i>Drepane punctata</i> | | | | | | | 1 | 40 | | | | | | | |
| | <i>Dendrophysa russelii</i> | 1 | 20 | | 2 | 50 | | 2 | 20 | | 3 | 60 | | 3 | 90 | |
| | <i>Epinephelus awoara</i> | | | | | | | | | | | | | | | |
| | <i>Epinephelus bruneus</i> | | | | | | | | | | | | | | | |
| | <i>Erisphex pottii</i> | 3 | 30 | | 3 | 20 | | 3 | 20 | | 6 | 50 | | 7 | 40 | |
| | <i>Gerres lucidus</i> | 2 | 60 | | | | | 2 | 20 | | | | | 1 | 20 | |
| | <i>Grammolites scaber</i> | | | | | | | | | | 1 | 10 | | | | |
| | <i>Gymnura japonica</i> | 1 | 3800 | | | | | | | | 1 | 501 | | 3 | 1430 | |
| | <i>Ilisha elongata</i> | | | | | | | | | | | | | | | |
| | <i>Inegocia japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Inimicus didactylus</i> | | | | 1 | 110 | | | | | | | | | | |
| | <i>Inimicus japonicus</i> | | | | | | | 3 | 160 | | | | | | | |
| | <i>Johnius belangerii</i> | | | | | | | | | | | | | | | |
| | <i>Johnius carona</i> | | | | | | | | | | | | | | | |
| | <i>Johnius poecilnotus</i> | | | | | | | | | | | | | | | |
| | <i>Johnius taiwanense</i> | | | | 1 | 30 | | 1 | 30 | | 1 | 20 | | 1 | 30 | |
| | <i>Konosirus punctatus</i> | | | | | | | | | | | | | | | |
| | <i>Lagocephalus wheeleri</i> | | | | | | | 1 | 10 | | 2 | 30 | | | | |
| | <i>Lepidotrigla japonice</i> | | | | | | | | | | | | | | | |
| | <i>Leiognathus nuchalis</i> | 47 | 470 | | 44 | 380 | | 75 | 660 | | 125 | 1190 | | 59 | 580 | |
| | <i>Lisa macrolepis</i> | | | | | | | | | | | | | | | |
| | <i>Monacanthus chinensis</i> | | | | | | | | | | | | | | | |
| | <i>Muraenesox cinereus</i> | | | | | | | | | | | | | | | |
| | <i>Minous monodactylus</i> | | | | 2 | 10 | | | | | | | | | | |
| | <i>Navodon tessellates</i> | | | | 1 | 5 | | | | | | | | | | |
| | <i>Nemipterus japonicus</i> | | | | | | | | | | 1 | 18 | | 2 | 20 | |
| | <i>Nibea acuta</i> | | | | | | | | | | | | | | | |
| | <i>Nibea albiflora</i> | | | | | | | | | | | | | | | |

Trawling Data in July 2019

| Group | Species Name | TNB | | | | | | | | | | | | | | |
|-----------------|------------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | TNB-1 | | | TNB-2 | | | TNB-3 | | | TNB-4 | | | TNB-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Oxurichthys papuensis</i> | | | | | | | 1 | 10 | | 1 | 5 | | | | |
| | <i>Parachaeturichthys polynema</i> | | | | | | | | | | | | | 1 | 10 | |
| | <i>Paraplagusia japonica</i> | | | | | | | | | | | | | | | |
| | <i>Pennahia anea</i> | 2 | 30 | | | | | 2 | 30 | | 7 | 60 | | 3 | 40 | |
| | <i>Pennahia argentata</i> | | | | | | | | | | | | | | | |
| | <i>Platycephalus indicus</i> | 5 | 240 | | 18 | 1290 | | 13 | 830 | | 9 | 390 | | 4 | 170 | |
| | <i>Plectorhinchus pictus</i> | | | | | | | | | | | | | | | |
| | <i>Plotosus lineatus</i> | | | | | | | | | | | | | | | |
| | <i>Polydactylus sextarius</i> | | | | | | | | | | 1 | 10 | | | | |
| | <i>Pseudorhombus arsius</i> | | | | | | | | | | | | | | | |
| | <i>Pseudorhombus cinnamomeus</i> | 2 | 50 | | 1 | 30 | | 4 | 40 | | | | | 1 | 50 | |
| | <i>Pseudorhombus oligodon</i> | | | | 1 | 10 | | 1 | 10 | | 1 | 10 | | 1 | 10 | |
| | <i>Saurida elongata</i> | | | | | | | | | | | | | | | |
| | Sciaenidae | | | | | | | | | | | | | | | |
| | <i>Scoliodon sorrakowah</i> | | | | | | | | | | | | | | | |
| | <i>Scorpaenopsis longispina</i> | 35 | 220 | | 9 | 60 | | 17 | 90 | | 8 | 80 | | 6 | 60 | |
| | <i>Secutor ruconius</i> | | | | | | | | | | | | | | | |
| | <i>Sebastiscus marmoratus</i> | 6 | 250 | | | | | | | | | | | | | |
| | <i>Siganus fuscescens</i> | | | | | | | | | | 1 | 30 | | | | |
| | <i>Sillago sihama</i> | | | | | | | | | | | | | | | |
| | <i>Solea ovata</i> | 1 | 10 | | | | | | | | | | | 1 | 10 | |
| | <i>Takifugu niphobles</i> | | | | | | | | | | | | | | | |
| | <i>Thryssa hamiltonii</i> | | | | | | | | | | | | | 1 | 20 | |
| | <i>Thryssa kammalensis</i> | | | | | | | | | | | | | | | |
| | <i>Trypauchen vagina</i> | 4 | 30 | | 5 | 50 | | 8 | 50 | | 2 | 27 | | 2 | 10 | |
| Prawn or shrimp | <i>Alpheus brevirostratus</i> | | | | | | | | | | | | | | | |
| | <i>Alpheus distinguendus</i> | | | | | | | | | | | | | | | |
| | <i>Exopalaemon carinicauda</i> | | | | | | | | | | | | | | | |
| | <i>Metapenaeopsis barbata</i> | | | | 2 | 8 | | | | | 1 | 10 | | 2 | 10 | |
| | <i>Metapenaeus affinis</i> | 62 | 520 | | 40 | 330 | | 57 | 500 | | 41 | 370 | | 23 | 180 | |
| | <i>Metapenaeus joyneri</i> | | | | 3 | 10 | | 1 | 6 | | | | | 2 | 18 | |
| | <i>Metapenaeus intermedius</i> | 4 | 30 | | 2 | 10 | | 3 | 20 | | 4 | 20 | | 3 | 20 | |
| | <i>Parapenaeopsis cultirostris</i> | | | | | | | | | | | | | | | |
| | <i>Parapenaeopsis cornutus</i> | | | | | | | | | | | | | | | |
| | <i>Parapenaeopsis hardwickii</i> | 1 | 10 | | 2 | 20 | | 2 | 20 | | 3 | 30 | | 2 | 20 | |
| | <i>Parapenaeopsis hungerfordii</i> | 6 | 30 | | 7 | 40 | | 11 | 60 | | 13 | 70 | | | | |
| | <i>Parapenaeopsis tenella</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus merguensis</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus penicillatus</i> | 24 | 190 | | 12 | 110 | | 28 | 190 | | 8 | 60 | | 4 | 30 | |
| | <i>Penaeus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus semisulcatus</i> | 2 | 20 | | | | | | | | 1 | 10 | | | | |
| | <i>Solenocera crassicornis</i> | | | | | | | | | | 6 | 20 | | 4 | 10 | |
| | <i>Trachypenaeus curvirostris</i> | | | | 1 | 5 | | 2 | 10 | | 2 | 10 | | 1 | 10 | |
| Crab | <i>Charybdis affinis</i> | 59 | 1020 | | 33 | 530 | | 20 | 320 | | 11 | 180 | | 23 | 360 | |
| | <i>Charybdis annulata</i> | | | | 1 | 20 | | | | | | | | | | |
| | <i>Charybdis bimaculata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis cruciata</i> | 1 | 10 | | | | | | | | | | | | | |
| | <i>Charybdis japonica</i> | | | | | | | | | | | | | | | |

Trawling Data in July 2019

| Group | Species Name | TNB | | | | | | | | | | | | | | |
|---------------|-----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | TNB-1 | | | TNB-2 | | | TNB-3 | | | TNB-4 | | | TNB-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Charybdis obtusifrons</i> | 4 | 120 | | 2 | 60 | | | | | | | | | | |
| | <i>Charybdis truncata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis variegata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis spp.</i> | 40 | 620 | * | 40 | 430 | * | 30 | 380 | * | 20 | 260 | * | 20 | 290 | * |
| | <i>Doclea sp.</i> | | | | | | | | | | | | | | | |
| | <i>Doclea ovis</i> | | | | | | | | | | | | | | | |
| | <i>Dorippe facchino</i> | | | | | | | | | | 20 | 220 | * | | | |
| | <i>Dorippe polita</i> | 7 | 78 | | 20 | 230 | * | | | | | | | | | |
| | <i>Eucrate crenata</i> | 1 | 5 | | | | | | | | 2 | 11.2 | | | | |
| | <i>Galene bispinosa</i> | | | | | | | | | | | | | | | |
| | Hermit crab | | | | | | | | | | | | | | | |
| | <i>Leucosia craniolepis</i> | | | | 40 | 230 | * | | | | 20 | 105 | * | 5 | 32 | |
| | <i>Macromedaeus distinguendus</i> | | | | | | | | | | | | | | | |
| | <i>Maorophthalmus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Matuta planipes</i> | | | | | | | | | | 1 | 8.2 | | | | |
| | <i>Ocypode ceratophthalmus</i> | | | | | | | | | | | | | | | |
| | <i>Parapanope euagora</i> | | | | | | | | | | | | | | | |
| | <i>Parthenope validus</i> | 20 | 240 | * | 20 | 240 | * | 16 | 210 | | 20 | 260 | * | 7 | 120 | |
| | <i>Plumnus hirtellus</i> | | | | | | | | | | | | | | | |
| | <i>Porcellanella picta</i> | 1 | 10 | | 1 | 5 | | | | | | | | | | |
| | <i>Portunus hastatoides</i> | | | | | | | 8 | 20 | | 7 | 20 | | | | |
| | <i>Portunus pelagicus</i> | | | | | | | 3 | 50 | | 2 | 140 | | | | |
| | <i>Portunus sanguinolentus</i> | | | | 1 | 10 | | | | | | | | | | |
| | <i>Portunus trituberculatus</i> | 1 | 40 | | 1 | 70 | | 4 | 180 | | 2 | 130 | | 2 | 60 | |
| | <i>Thalamita sima</i> | 1 | 10 | | | | | | | | | | | | | |
| | <i>Typhlocarcinus sp.</i> | | | | | | | | | | | | | | | |
| Mantis shrimp | <i>Anchisquilla fasciata</i> | | | | | | | | | | | | | | | |
| | <i>Dictyosquilla foveolata</i> | | | | | | | | | | | | | 1 | 10 | |
| | <i>Harpisquilla harpax</i> | | | | | | | | | | | | | 1 | 30 | |
| | <i>Oratosquilla interrupta</i> | 2 | 60 | | 2 | 50 | | 1 | 10 | | 10 | 170 | | 7 | 120 | |
| | <i>Oratosquilla oratoria</i> | | | | | | | | | | 4 | 30 | | 2 | 20 | |
| | <i>Oratosquilla kempfi</i> | | | | | | | | | | | | | | | |
| Gastropod | <i>Bursa rana</i> | | | | | | | 2 | 16 | | | | | | | |
| | <i>Calyptrea sp.</i> | 20 | 13 | * | 3 | 1.1 | | 2 | 0.9 | | 20 | 13 | * | 30 | 18 | * |
| | <i>Desmaulus extinctorius</i> | | | | | | | | | | | | | | | |
| | <i>Drupa musiva</i> | 4 | 21 | | | | | 2 | 10 | | | | | | | |
| | <i>Epitonium scalare</i> | | | | | | | | | | | | | | | |
| | <i>Inquistor flavidula</i> | | | | | | | | | | | | | | | |
| | <i>Gemmula deshayesii</i> | | | | | | | | | | | | | | | |
| | <i>Lophiotoma leucotropis</i> | | | | 1 | 5.2 | | | | | | | | | | |
| | <i>Murex trapa</i> | 20 | 160 | * | 20 | 170 | * | 20 | 160 | * | 30 | 210 | * | 40 | 410 | * |
| | <i>Nassarius hepaticus</i> | | | | | | | | | | | | | | | |
| | <i>Nassarius succinctus</i> | | | | 30 | 62 | * | | | | 10 | 21 | | 8 | 15 | |
| | <i>Philine sp.</i> | 2 | 11.6 | | | | | 1 | 5.9 | | | | | | | |
| | <i>Pilosabia pilosa</i> | | | | 5 | 3.2 | | 20 | 24 | * | | | | | | |
| | <i>Siphonotus walshi</i> | | | | 7 | 16.2 | | 3 | 4.2 | | | | | | | |
| | <i>Tonna sulcosa</i> | | | | | | | | | | | | | | | |
| | <i>Turritella terebra</i> | 960 | 9120 | * | 1200 | 11800 | * | 860 | 8240 | * | 720 | 7080 | * | 920 | 9130 | * |

Trawling Data in July 2019

| Group | Species Name | TNB | | | | | | | | | | | | | | |
|------------|------------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | TNB-1 | | | TNB-2 | | | TNB-3 | | | TNB-4 | | | TNB-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>UNID Nudibranch</i> | | | | | | | | | | 3 | 10.4 | | 2 | 7.9 | |
| Bivalve | <i>Anomia chinensis</i> | | | | | | | | | | | | | | | |
| | <i>Anomalodiscus squamosus</i> | | | | 1 | 3.4 | | | | | | | | | | |
| | <i>Chione isabellina</i> | | | | | | | | | | | | | | | |
| | <i>Chlamys nobilis</i> | | | | 2 | 19 | | | | | | | | | | |
| | <i>Macoma candida</i> | | | | | | | | | | | | | | | |
| | <i>Ostrea sp.</i> | | | | | | | | | | | | | | | |
| | <i>Paphia undulata</i> | | | | | | | | | | | | | | | |
| | <i>Parahyattisa sp.</i> | | | | | | | | | | | | | | | |
| | <i>Pinctada martensii</i> | 1 | 46 | | | | | 1 | 34 | | | | | | | |
| | <i>Placuna placenta</i> | | | | | | | | | | | | | | | |
| | <i>Ruditapes philippinarum</i> | | | | | | | | | | | | | | | |
| | <i>Samacar sp.</i> | | | | | | | | | | | | | | | |
| | <i>Scapharca subcrenata</i> | | | | | | | 1 | 29 | | | | | | | |
| | <i>Tegillarca nodifera</i> | | | | | | | | | | | | | | | |
| | <i>Tolonostrea talonata</i> | | | | | | | 2 | 5.4 | | | | | | | |
| | <i>Vepricardium asiaticum</i> | | | | | | | | | | | | | | | |
| Cephalopod | <i>Lolingid squid</i> | 1 | 10 | | | | | | | | 1 | 10 | | | | |
| | <i>Amphioctopus fangsiao</i> | 1 | 50 | | | | | 1 | 50 | | | | | | | |
| | <i>Sepiid cuttlefish</i> | | | | 1 | 260 | | | | | 2 | 18 | | | | |
| Echinoderm | <i>Acaudina molpadioides</i> | 7 | 270 | | 6 | 266 | | 6 | 260 | | 5 | 210 | | 5 | 210 | |
| | <i>Colochirus quadrangularis</i> | | | | | | | | | | | | | | | |
| | <i>Protankyra bidentata (c.f.)</i> | | | | | | | | | | | | | | | |
| | <i>Styela plicata</i> | | | | 2 | 10 | | 4 | 24 | | | | | | | |
| | <i>Temnopleurus toreumaticus</i> | 40 | 570 | * | 40 | 580 | * | 30 | 440 | * | 40 | 580 | * | 30 | 460 | * |
| Cnidaria | <i>Carcinactis ichikawai</i> | | | | 3 | 1.6 | | | | | | | | | | |
| | <i>Cavernularia habereri</i> | 6 | 32 | | 50 | 329 | * | 7 | 38 | | 20 | 180 | * | 30 | 280 | * |
| | <i>Comanthus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Edwardsia japonica (c.f.)</i> | 2 | 18 | | 4 | 31 | | 2 | 21 | | | | | | | |
| | <i>Ellisella laevis</i> | | | | 3 | 26 | | | | | | | | | | |
| | Jelly fish | | | | | | | | | | | | | | | |
| | <i>Pteroeides chinense</i> | 13 | 460 | | 19 | 1270 | | 11 | 410 | | | | | 7 | 170 | |
| | <i>Gorgonia spp.</i> | | | | | | | | | | | | | | | |
| | Sea anomea | | | | 2 | 12 | | | | | 2 | 13 | | 3 | 17 | |
| | <i>Virgularia gustaviana</i> | 4 | 120 | | 1 | 22 | | 2 | 110 | | | | | 3 | 22 | |
| Porifera | Spongiatia | | | | | | | | | | | | | | | |
| Cirripedia | <i>c.f. Bananus spp.</i> | 140 | 130 | * | 160 | 140 | * | 280 | 260 | * | 190 | 120 | * | 160 | 110 | * |

Trawling Data in July 2019

| Group | Species Name | TSA-1 | | | TSA-2 | | | TSA | | | TSA-4 | | | TSA-5 | | |
|-------|--------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| Fish | <i>Arnoglossus waitei</i> | | | | | | | | | | | | | | | |
| | <i>Acentrogobius caninus</i> | | | | 1 | 10 | | 3 | 40 | | | | | 1 | 20 | |
| | <i>Anchoviella commersonii</i> | 2 | 10 | | 2 | 10 | | | | | | | | | | |
| | <i>Apogon kiensis</i> | | | | | | | | | | | | | | | |
| | <i>Apogon lineatus</i> | 23 | 50 | | 27 | 90 | | 16 | 50 | | 15 | 40 | | 25 | 70 | |
| | <i>Arius arius</i> | | | | 2 | 110 | | | | | | | | | | |
| | <i>Callionymus richardsoni</i> | | | | | | | | | | | | | | | |
| | <i>Callionymus curvicornis</i> | | | | | | | | | | | | | | | |
| | <i>Caranx kalla</i> | | | | | | | | | | | | | | | |
| | <i>Clupanodon thrissa</i> | | | | | | | | | | 1 | 50 | | | | |
| | <i>Collichthys lucidus</i> | 9 | 40 | | 16 | 88 | | 8 | 60 | | 26 | 150 | | 21 | 130 | |
| | <i>Coilia mystus</i> | | | | | | | | | | | | | | | |
| | <i>Crossorhombus kanekonis</i> | | | | | | | | | | | | | | | |
| | <i>Cryptocentrus filifer</i> | | | | | | | | | | | | | | | |
| | <i>Cynoglossus abbreviatus</i> | | | | | | | | | | | | | 1 | 10 | |
| | <i>Cynoglossus arel</i> | | | | 2 | 80 | | 10 | 270 | | 8 | 310 | | 3 | 50 | |
| | <i>Cynoglossus puncticeps</i> | | | | | | | | | | 1 | 20 | | | | |
| | <i>Cynoglossus roulei</i> | | | | 1 | 650 | | | | | | | | | | |
| | <i>Dasyatis akaiei</i> | | | | | | | | | | | | | | | |
| | <i>Dasyatis zugei</i> | | | | 2 | 1270 | | | | | 1 | 820 | | 3 | 790 | |
| | <i>Drepane punctata</i> | | | | | | | | | | | | | | | |
| | <i>Dendrophysa russelii</i> | 1 | 8 | | 3 | 60 | | 2 | 60 | | 6 | 150 | | 8 | 240 | |
| | <i>Epinephelus awoara</i> | | | | | | | | | | | | | | | |
| | <i>Epinephelus bruneus</i> | | | | | | | | | | | | | | | |
| | <i>Erisphex pottii</i> | 15 | 110 | | 22 | 200 | | 23 | 201 | | 7 | 70 | | 4 | 30 | |
| | <i>Gerres lucidus</i> | 9 | 210 | | 15 | 340 | | 21 | 490 | | 17 | 420 | | 25 | 510 | |
| | <i>Grammoplites scaber</i> | | | | | | | | | | | | | | | |
| | <i>Gymnura japonica</i> | | | | | | | | | | 1 | 680 | | | | |
| | <i>Ilisha elongata</i> | | | | 1 | 5 | | | | | | | | | | |
| | <i>Inegocia japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Inimicus didactylus</i> | | | | 1 | 220 | | | | | | | | | | |
| | <i>Inimicus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Johnius belangerii</i> | | | | | | | | | | | | | | | |
| | <i>Johnius carona</i> | | | | | | | | | | 2 | 40 | | | | |
| | <i>Johnius poecilnotus</i> | | | | | | | | | | | | | | | |
| | <i>Johnius taiwanense</i> | | | | | | | | | | | | | 1 | 50 | |
| | <i>Konosirus punctatus</i> | 1 | 40 | | 1 | 40 | | | | | | | | 3 | 200 | |
| | <i>Lagocephalus wheeleri</i> | | | | | | | | | | | | | | | |
| | <i>Lepidotrigla japonice</i> | | | | | | | | | | | | | | | |
| | <i>Leiognathus nuchalis</i> | 81 | 680 | | 175 | 1670 | | 80 | 650 | | 155 | 1600 | | 95 | 1080 | |
| | <i>Lisa macrolepis</i> | | | | 1 | 40 | | | | | | | | | | |
| | <i>Monacanthus chinensis</i> | | | | | | | | | | | | | | | |
| | <i>Muraenesox cinereus</i> | | | | | | | | | | | | | | | |
| | <i>Minous monodactylus</i> | | | | | | | | | | | | | | | |
| | <i>Navodon tessellates</i> | | | | | | | | | | | | | | | |
| | <i>Nemipterus japonicus</i> | 1 | 10 | | 13 | 90 | | 13 | 110 | | 13 | 100 | | 6 | 50 | |
| | <i>Nibea acuta</i> | | | | | | | 1 | 240 | | 1 | 90 | | | | |
| | <i>Nibea albiflora</i> | | | | | | | | | | | | | 1 | 30 | |

Trawling Data in July 2019

| Group | Species Name | TSA-1 | | | TSA-2 | | | TSA-3 | | | TSA-4 | | | TSA-5 | | |
|-----------------|------------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Oxyurichthys papuensis</i> | 11 | 50 | | 4 | 20 | | 7 | 30 | | 3 | 10 | | 2 | 10 | |
| | <i>Parachaeturichthys polynema</i> | | | | | | | 4 | 30 | | | | | | | |
| | <i>Paraplagusia japonica</i> | | | | 1 | 10 | | | | | 2 | 10 | | | | |
| | <i>Pennahia anea</i> | 2 | 30 | | 5 | 70 | | 6 | 80 | | 11 | 150 | | 11 | 176 | |
| | <i>Pennahia argentata</i> | | | | | | | | | | | | | | | |
| | <i>Platycephalus indicus</i> | 5 | 430 | | 14 | 1090 | | 21 | 1690 | | 9 | 380 | | 14 | 1170 | |
| | <i>Plectorhinchus pictus</i> | | | | | | | | | | | | | | | |
| | <i>Plotosus lineatus</i> | | | | | | | | | | | | | | | |
| | <i>Polydactylus sextarius</i> | | | | | | | | | | | | | | | |
| | <i>Pseudorhombus arsius</i> | | | | | | | | | | | | | | | |
| | <i>Pseudorhombus cinnamomeus</i> | | | | 3 | 20 | | 2 | 10 | | | | | | | |
| | <i>Pseudorhombus oligodon</i> | 8 | 70 | | 10 | 130 | | 11 | 110 | | 6 | 70 | | 16 | 190 | |
| | <i>Saurida elongata</i> | | | | | | | | | | 1 | 20 | | | | |
| | Sciaenidae | | | | | | | | | | | | | | | |
| | <i>Scoliodon sorrakowah</i> | | | | 2 | 30 | | 1 | 20 | | 2 | 60 | | 1 | 20 | |
| | <i>Scorpaenopsis longispina</i> | 8 | 50 | | 4 | 20 | | 2 | 20 | | 1 | 5 | | | | |
| | <i>Secutor ruconius</i> | | | | 1 | 3 | | | | | | | | | | |
| | <i>Sebastiscus marmoratus</i> | | | | | | | | | | | | | | | |
| | <i>Siganus fuscescens</i> | | | | 1 | 2 | | | | | 1 | 5 | | 2 | 30 | |
| | <i>Sillago sihama</i> | | | | 2 | 30 | | | | | | | | 1 | 20 | |
| | <i>Solea ovata</i> | | | | 2 | 20 | | 2 | 15 | | | | | | | |
| | <i>Takifugu niphobles</i> | 5 | 1420 | | 10 | 3010 | | 2 | 370 | | 3 | 1060 | | 1 | 330 | |
| | <i>Thryssa hamiltonii</i> | 1 | 10 | | 1 | 20 | | | | | | | | | | |
| | <i>Thryssa kammalensis</i> | | | | | | | | | | | | | | | |
| | <i>Trypauchen vagina</i> | 6 | 25 | | 11 | 90 | | 15 | 150 | | 2 | 40 | | 7 | 170 | |
| Prawn or shrimp | <i>Alpheus brevicristatus</i> | | | | | | | | | | | | | | | |
| | <i>Alpheus distinguendus</i> | | | | | | | | | | | | | | | |
| | <i>Exopalaemon carinicauda</i> | | | | | | | | | | | | | | | |
| | <i>Metapenaeopsis barbata</i> | | | | | | | | | | | | | | | |
| | <i>Metapenaeus affinis</i> | 39 | 120 | | 66 | 228 | | 40 | 160 | | 17 | 60 | | 26 | 110 | |
| | <i>Metapenaeus joyneri</i> | 8 | 30 | | 7 | 30 | | 4 | 20 | | 12 | 60 | | 1 | 10 | |
| | <i>Metapenaeus intermedius</i> | 1 | 10 | | 6 | 120 | | 10 | 60 | | 16 | 100 | | 13 | 110 | |
| | <i>Parapenaeopsis cultirostris</i> | | | | | | | | | | | | | | | |
| | <i>Parapenaeopsis cornutus</i> | | | | | | | | | | | | | | | |
| | <i>Parapenaeopsis hardwickii</i> | | | | | | | | | | 1 | 5 | | | | |
| | <i>Parapenaeopsis hungerfordii</i> | 36 | 170 | | 80 | 470 | | 47 | 230 | | 40 | 170 | | 46 | 360 | |
| | <i>Parapenaeopsis tenella</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus merguensis</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus penicillatus</i> | 4 | 40 | | 5 | 50 | | 3 | 30 | | | | | 1 | 10 | |
| | <i>Penaeus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus semisulcatus</i> | | | | | | | | | | | | | | | |
| | <i>Solenocera crassicornis</i> | | | | | | | | | | | | | | | |
| | <i>Trachypenaeus curvirostris</i> | | | | | | | | | | | | | | | |
| Crab | <i>Charybdis affinis</i> | 165 | 1810 | | 185 | 2905 | | 95 | 1450 | | 125 | 1450 | | 80 | 1150 | |
| | <i>Charybdis annulata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis bimaculata</i> | | | | | | | | | | | | | 2 | 6 | |
| | <i>Charybdis cruciata</i> | | | | 1 | 80 | | 1 | 30 | | 2 | 20 | | 1 | 20 | |
| | <i>Charybdis japonica</i> | | | | | | | | | | | | | | | |

Trawling Data in July 2019

| Group | Species Name | TSA-1 | | | TSA-2 | | | TSA | | | TSA-3 | | | TSA-4 | | | TSA-5 | | |
|---------------|-----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Charybdis obtusifrons</i> | 1 | 80 | | 1 | 40 | | 1 | 40 | | 2 | 110 | | 4 | 40 | | | | |
| | <i>Charybdis truncata</i> | 40 | 30 | | 4 | 20 | | 3 | 10 | | 12 | 50 | | 7 | 30 | | | | |
| | <i>Charybdis variegata</i> | 1 | 5 | | 4 | 10 | | | | | 1 | 2 | | | | | | | |
| | <i>Charybdis spp.</i> | 30 | 380 | * | 30 | 380 | * | 20 | 310 | * | 40 | 460 | * | 30 | 320 | * | | | |
| | <i>Doclea sp.</i> | | | | | | | | | | | | | | | | | | |
| | <i>Doclea ovis</i> | | | | | | | | | | | | | 1 | 42 | | | | |
| | <i>Dorippe facchino</i> | | | | | | | | | | | | | | | | | | |
| | <i>Dorippe polita</i> | 40 | 430 | * | 30 | 340 | * | 30 | 340 | * | 10 | 78 | | 20 | 230 | * | | | |
| | <i>Eucrate crenata</i> | | | | | | | | | | | | | | | | | | |
| | <i>Galene bispinosa</i> | | | | 1 | 90 | | | | | 6 | 370 | | 5 | 310 | | | | |
| | Hermit crab | 50 | 160 | * | 60 | 230 | * | 50 | 210 | * | 50 | 210 | * | 70 | 190 | * | | | |
| | <i>Leucosia craniolaris</i> | | | | 20 | 110 | * | 20 | 98 | * | 20 | 130 | * | | | | | | |
| | <i>Macromedaeus distinguendus</i> | | | | | | | | | | | | | | | | | | |
| | <i>Maorophthalmus japonicus</i> | | | | | | | | | | 3 | 11.6 | | 1 | 4.1 | | | | |
| | <i>Matuta planipes</i> | | | | | | | | | | | | | | | | | | |
| | <i>Ocypode ceratophthalmus</i> | 4 | 10 | | | | | | | | | | | | | | | | |
| | <i>Parapanope euagora</i> | | | | | | | | | | | | | | | | | | |
| | <i>Parthenope validus</i> | | | | | | | | | | | | | | | | | | |
| | <i>Plumnus hirtellus</i> | | | | | | | | | | | | | | | | | | |
| | <i>Porcellanella picta</i> | | | | | | | | | | | | | | | | | | |
| | <i>Portunus hastatoides</i> | 47 | 90 | | 56 | 130 | | 31 | 60 | | 53 | 110 | | 32 | 100 | | | | |
| | <i>Portunus pelagicus</i> | 1 | 40 | | 4 | 160 | | | | | 3 | 110 | | | | | | | |
| | <i>Portunus sanguinolentus</i> | | | | 1 | 20 | | | | | 1 | 10 | | | | | | | |
| | <i>Portunus trituberculatus</i> | | | | 1 | 60 | | | | | | | | | | | | | |
| | <i>Thalamita sima</i> | 6 | 60 | | 5 | 50 | | 4 | 20 | | 4 | 10 | | 2 | 10 | | | | |
| | <i>Typhlocarcinus sp.</i> | | | | | | | | | | | | | | | | | | |
| Mantis shrimp | <i>Anchisquilla fasciata</i> | | | | | | | | | | | | | | | | | | |
| | <i>Dictyosquilla foveolata</i> | | | | 1 | 10 | | | | | 4 | 30 | | 3 | 40 | | | | |
| | <i>Harpisquilla harpax</i> | | | | 2 | 140 | | | | | 5 | 160 | | 3 | 110 | | | | |
| | <i>Oratosquilla interrupta</i> | | | | 3 | 50 | | 4 | 80 | | 16 | 398 | | 6 | 130 | | | | |
| | <i>Oratosquilla oratoria</i> | | | | | | | | | | 4 | 60 | | | | | | | |
| | <i>Oratosquilla kempfi</i> | | | | | | | | | | | | | | | | | | |
| Gastropod | <i>Bursa rana</i> | | | | | | | | | | | | | 1 | 8.2 | | | | |
| | <i>Calyptreaea sp.</i> | | | | | | | | | | | | | | | | | | |
| | <i>Desmaulus extingtorius</i> | | | | | | | | | | | | | | | | | | |
| | <i>Drupa musiva</i> | | | | 4 | 22 | | | | | | | | | | | | | |
| | <i>Epitonium scalare</i> | | | | | | | | | | | | | 1 | 8.4 | | | | |
| | <i>Inquistor flavidula</i> | 20 | 43 | * | | | | | | | | | | | | | | | |
| | <i>Gemmula deshavesii</i> | | | | | | | | | | | | | | | | | | |
| | <i>Lophiotoma leucotropis</i> | | | | | | | | | | | | | | | | | | |
| | <i>Murex trapa</i> | 90 | 1120 | * | 70 | 730 | * | 180 | 1790 | * | 60 | 640 | * | 80 | 880 | * | | | |
| | <i>Nassarius hepaticus</i> | | | | | | | | | | | | | | | | | | |
| | <i>Nassarius succinctus</i> | 40 | 92 | * | 80 | 170 | * | 30 | 71 | * | | | | | | | | | |
| | <i>Philine sp.</i> | 4 | 1.6 | | 2 | 0.7 | | 5 | 1.6 | | 4 | 1.2 | | | | | | | |
| | <i>Pilosabia pilosa</i> | 20 | 12 | * | | | | 3 | 1 | | | | | | | | | | |
| | <i>Siphopatella walshi</i> | | | | 20 | 11 | * | | | | | | | | | | | | |
| | <i>Tonna sulcosa</i> | | | | | | | | | | | | | | | | | | |
| | <i>Turritella terebra</i> | 390 | 3760 | * | 460 | 4340 | * | 390 | 3780 | * | 420 | 3970 | * | 530 | 5180 | * | | | |

Trawling Data in July 2019

| Group | Species Name | TSA-1 | | | TSA-2 | | | TSA | | | TSA-4 | | | TSA-5 | | |
|------------|------------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>UNID Nudibranch</i> | | | | | | | | | | | | | | | |
| Bivalve | <i>Anomia chinensis</i> | | | | | | | | | | | | | | | |
| | <i>Anomalodiscus squamosus</i> | | | | | | | | | | | | | | | |
| | <i>Chione isabellina</i> | 40 | 260 | * | 30 | 170 | * | | | | | | | | | |
| | <i>Chlamys nobilis</i> | 30 | 360 | * | | | | | | | | | | 20 | 270 | * |
| | <i>Macoma candida</i> | 20 | 130 | * | | | | | | | | | | | | |
| | <i>Ostrea sp.</i> | | | | | | | | | | | | | | | |
| | <i>Paphia undulata</i> | 20 | 130 | * | | | | | | | | | | 20 | 180 | * |
| | <i>Parahyatisa sp.</i> | | | | | | | | | | | | | | | |
| | <i>Pinctada martensii</i> | | | | | | | | | | | | | | | |
| | <i>Placuna placenta</i> | | | | 1 | 11.6 | | | | | | | | | | |
| | <i>Ruditapes philippinarum</i> | 30 | 160 | * | | | | | | | | | | | | |
| | <i>Samacar sp.</i> | | | | | | | | | | | | | | | |
| | <i>Scapharca subcrenata</i> | 3 | 170 | | 2 | 112 | | 1 | 46 | | | | | | | |
| | <i>Tegillarca nodifera</i> | | | | | | | | | | | | | 30 | 180 | * |
| | <i>Tolonostrea talonata</i> | | | | 2 | 6.2 | | | | | | | | | | |
| | <i>Vepricardium asiaticum</i> | | | | 1 | 5.8 | | | | | | | | | | |
| Cephalopod | <i>Lolingid squid</i> | 1 | 10 | | 2 | 30 | | 1 | 20 | | | | | | | |
| | <i>Amphioctopus fangsiao</i> | | | | | | | | | | | | | | | |
| | <i>Sepiid cuttlefish</i> | | | | | | | | | | | | | | | |
| Echinoderm | <i>Acaudina molpadioides</i> | 20 | 620 | * | 20 | 640 | * | 4 | 78 | | 7 | 180 | | | | |
| | <i>Colochirus quadrangularis</i> | | | | | | | | | | | | | | | |
| | <i>Protankyra bidentata (c.f.)</i> | | | | | | | | | | | | | | | |
| | <i>Styela plicata</i> | 1 | 4.2 | | | | | | | | | | | 2 | 9.4 | |
| | <i>Temnopleurus toreumaticus</i> | 30 | 370 | * | 30 | 380 | * | 30 | 370 | * | 20 | 240 | * | 30 | 410 | * |
| Cnidaria | <i>Carcinactis ichikawai</i> | 5 | 4.1 | | 1 | 0.5 | | 1 | 0.6 | | 3 | 1.2 | | 3 | 1.4 | |
| | <i>Cavernularia habereri</i> | 20 | 180 | * | 30 | 170 | * | 20 | 78 | * | 11 | 73 | | 6 | 46 | |
| | <i>Comanthus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Edwardsia japonica (c.f.)</i> | 1 | 6.4 | | | | | | | | 1 | 4.3 | | | | |
| | <i>Ellisella laevis</i> | | | | | | | | | | | | | | | |
| | Jelly fish | | | | | | | | | | | | | | | |
| | <i>Pteroeides chinense</i> | 20 | 760 | * | 20 | 770 | * | 8 | 310 | | 6 | 130 | | 5 | 140 | |
| | <i>Gorgonia spp.</i> | | | | 1 | 19 | | | | | | | | | | |
| | Sea anomea | 2 | 12 | | 1 | 8 | | | | | 2 | 11 | | | | |
| | <i>Virgularia gustaviana</i> | | | | 1 | 9.4 | | 3 | 42 | | 4 | 44 | | 6 | 114 | |
| Porifera | Spongiatia | | | | | | | | | | | | | | | |
| Cirripedia | <i>c.f. Bananus spp.</i> | | | | | | | | | | | | | | | |

Trawling Data in July 2019

| Group | Species Name | TSB | | | | | | | | | | | | | | |
|-------|--------------------------------|-----------------|--------------------|--------------------|-----------------|--------------------|--------------------|-----------------|--------------------|--------------------|-----------------|--------------------|--------------------|-----------------|--------------------|--------------------|
| | | TSB-1 | | | TSB-2 | | | TSB-3 | | | TSB-4 | | | TSB-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| Fish | <i>Arnoglossus waitei</i> | | | | | | | | | | | | | | | |
| | <i>Acentrogobius caninus</i> | | | | 1 | 10 | | 2 | 40 | | 3 | 40 | | 2 | 40 | |
| | <i>Anchoviella commersonii</i> | 1 | 5 | | | | | | | | | | | 1 | 5 | |
| | <i>Apogon kiensis</i> | 1 | 10 | | | | | 5 | 20 | | 3 | 10 | | | | |
| | <i>Apogon lineatus</i> | | | | | | | | | | | | | 3 | 10 | |
| | <i>Arius arius</i> | 1 | 40 | | 1 | 40 | | | | | 4 | 540 | | | | |
| | <i>Callionymus richardsoni</i> | 1 | 6 | | | | | | | | | | | | | |
| | <i>Callionymus curvicornis</i> | 1 | 10 | | | | | | | | 1 | 10 | | | | |
| | <i>Caranx kalla</i> | | | | | | | | | | | | | | | |
| | <i>Clupanodon thrissa</i> | | | | | | | | | | | | | | | |
| | <i>Collichthys lucidus</i> | 20 | 120 | | 24 | 140 | | 18 | 90 | | 22 | 156 | | 23 | 90 | |
| | <i>Coilia mystus</i> | | | | | | | | | | | | | | | |
| | <i>Crossorhombus kanekonis</i> | | | | | | | | | | | | | | | |
| | <i>Cryptocentrus filifer</i> | | | | | | | | | | | | | | | |
| | <i>Cynoglossus abbreviatus</i> | 1 | 10 | | | | | | | | | | | | | |
| | <i>Cynoglossus arel</i> | 3 | 72 | | 6 | 210 | | 8 | 220 | | 2 | 40 | | 5 | 173 | |
| | <i>Cynoglossus puncticeps</i> | | | | | | | | | | | | | 1 | 10 | |
| | <i>Cynoglossus roulei</i> | | | | | | | | | | | | | | | |
| | <i>Dasyatis akaiei</i> | | | | | | | | | | | | | | | |
| | <i>Dasyatis zugei</i> | 6 | 1430 | | 6 | 1570 | | 13 | 2170 | | 2 | 210 | | 2 | 350 | |
| | <i>Drepane punctata</i> | | | | | | | | | | | | | | | |
| | <i>Dendrophysa russelii</i> | 12 | 168 | | 7 | 100 | | 39 | 490 | | 2 | 40 | | | | |
| | <i>Epinephelus awoara</i> | | | | 1 | 10 | | | | | | | | | | |
| | <i>Epinephelus bruneus</i> | | | | | | | | | | 1 | 20 | | | | |
| | <i>Erisphex pottii</i> | | | | 1 | 20 | | 7 | 80 | | 3 | 20 | | 6 | 50 | |
| | <i>Gerres lucidus</i> | 7 | 170 | | 9 | 220 | | 20 | 440 | | 7 | 190 | | 5 | 180 | |
| | <i>Grammoplites scaber</i> | | | | | | | | | | | | | | | |
| | <i>Gymnura japonica</i> | | | | | | | | | | | | | | | |
| | <i>Ilisha elongata</i> | | | | | | | | | | | | | | | |
| | <i>Inegocia japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Inimicus didactylus</i> | | | | | | | | | | | | | | | |
| | <i>Inimicus japonicus</i> | | | | | | | 1 | 30 | | | | | | | |
| | <i>Johnius belangerii</i> | | | | | | | | | | | | | | | |
| | <i>Johnius carona</i> | 1 | 6 | | 1 | 10 | | 4 | 10 | | 2 | 30 | | | | |
| | <i>Johnius poecilonotus</i> | | | | | | | | | | | | | | | |
| | <i>Johnius taiwanense</i> | 2 | 90 | | 2 | 70 | | 1 | 10 | | 2 | 180 | | 4 | 110 | |
| | <i>Konosirus punctatus</i> | | | | 1 | 30 | | 1 | 50 | | 12 | 610 | | 10 | 540 | |
| | <i>Lagocephalus wheeleri</i> | | | | | | | | | | | | | | | |
| | <i>Lepidotrigla japonice</i> | | | | | | | | | | | | | | | |
| | <i>Leiognathus nuchalis</i> | 81 | 850 | | 103 | 1210 | | 67 | 785 | | 158 | 1700 | | 148 | 1610 | |
| | <i>Lisa macrolepis</i> | | | | | | | | | | 1 | 40 | | | | |
| | <i>Monacanthus chinensis</i> | | | | | | | | | | | | | | | |
| | <i>Muraenesox cinereus</i> | | | | 1 | 130 | | | | | | | | | | |
| | <i>Minous monodactylus</i> | | | | | | | | | | | | | | | |
| | <i>Navodon tessellates</i> | | | | | | | | | | | | | | | |
| | <i>Nemipterus japonicus</i> | 6 | 60 | | 3 | 40 | | 13 | 110 | | 3 | 30 | | 1 | 90 | |
| | <i>Nibea acuta</i> | | | | | | | 1 | 130 | | 1 | 90 | | | | |
| | <i>Nibea albiflora</i> | | | | | | | | | | | | | | | |

Trawling Data in July 2019

| Group | Species Name | TSB | | | | | | | | | | | | | | |
|-----------------|------------------------------------|-----------------|--------------------|--------------------|-----------------|--------------------|--------------------|-----------------|--------------------|--------------------|-----------------|--------------------|--------------------|-----------------|--------------------|--------------------|
| | | TSB-1 | | | TSB-2 | | | TSB-3 | | | TSB-4 | | | TSB-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Oxyurichthys papuensis</i> | 1 | 10 | | | | 1 | 5 | | 3 | 10 | | 3 | 10 | | |
| | <i>Parachaeturichthys polynema</i> | 1 | 10 | | | | | | | | | | 1 | 10 | | |
| | <i>Paraplagusia japonica</i> | | | | | | | | | | | | | | | |
| | <i>Pennahia anea</i> | | | | 2 | 30 | | 3 | 50 | | 3 | 30 | | 2 | 20 | |
| | <i>Pennahia argentata</i> | | | | | | | | | | | | | | | |
| | <i>Platycephalus indicus</i> | 2 | 150 | | 6 | 370 | | 9 | 850 | | 2 | 130 | | 12 | 1100 | |
| | <i>Plectorhinchus pictus</i> | | | | | | | | | | | | | | | |
| | <i>Plotosus lineatus</i> | | | | | | | | | | | | | | | |
| | <i>Polydactylus sextarius</i> | | | | | | | | | | | | 2 | 10 | | |
| | <i>Pseudorhombus arsius</i> | | | | | | | | | | | | | | | |
| | <i>Pseudorhombus cinnamomeus</i> | 1 | 10 | | 7 | 90 | | 19 | 210 | | 14 | 170 | | | | |
| | <i>Pseudorhombus oligodon</i> | 4 | 40 | | 2 | 20 | | 2 | 30 | | | | | 6 | 40 | |
| | <i>Saurida elongata</i> | 1 | 20 | | | | | | | | | | | | | |
| | <i>Sciaenidae</i> | | | | | | | | | | | | | | | |
| | <i>Scoliodon sorrakowah</i> | 2 | 30 | | 1 | 30 | | 2 | 60 | | | | | 1 | 20 | |
| | <i>Scorpaenopsis longispina</i> | | | | | | | | | | | | | 1 | 10 | |
| | <i>Secutor ruconius</i> | | | | | | | | | | | | | | | |
| | <i>Sebastiscus marmoratus</i> | | | | | | | | | | | | | | | |
| | <i>Siganus fuscescens</i> | | | | | | | | | | 1 | 10 | | | | |
| | <i>Sillago sihama</i> | 1 | 20 | | | | | 2 | 120 | | | | | | | |
| | <i>Solea ovata</i> | | | | | | | 1 | 10 | | 2 | 20 | | 2 | 20 | |
| | <i>Takifugu niphobles</i> | 5 | 1690 | | 7 | 2180 | | 9 | 2520 | | 7 | 2520 | | 9 | 3160 | |
| | <i>Thryssa hamiltonii</i> | | | | | | | | | | | | | | | |
| | <i>Thryssa kammalensis</i> | 1 | 5 | | | | | | | | | | | | | |
| | <i>Trypauchen vagina</i> | 4 | 60 | | 4 | 60 | | 6 | 60 | | 2 | 20 | | 3 | 30 | |
| Prawn or shrimp | <i>Alpheus brevicristatus</i> | | | | | | | | | | | | | | | |
| | <i>Alpheus distinguendus</i> | | | | | | | | | | 2 | 8 | | | | |
| | <i>Exopalaemon carinicauda</i> | | | | | | | | | | | | | | | |
| | <i>Metapenaeopsis barbata</i> | | | | | | | | | | | | | | | |
| | <i>Metapenaeus affinis</i> | 1 | 5 | | 1 | 10 | | 5 | 20 | | 5 | 20 | | 6 | 23 | |
| | <i>Metapenaeus joyneri</i> | 3 | 10 | | 9 | 40 | | 6 | 24 | | 13 | 70 | | 8 | 40 | |
| | <i>Metapenaeus intermedius</i> | | | | 5 | 20 | | 6 | 60 | | 5 | 50 | | 8 | 70 | |
| | <i>Parapenaeopsis cultirostris</i> | | | | | | | | | | | | | | | |
| | <i>Parapenaeopsis cornutus</i> | | | | | | | | | | | | | | | |
| | <i>Parapenaeopsis hardwickii</i> | | | | | | | | | | | | | | | |
| | <i>Parapenaeopsis hungerfordii</i> | 16 | 100 | | 34 | 200 | | 46 | 260 | | 25 | 145 | | 22 | 120 | |
| | <i>Parapenaeopsis tenella</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus merguensis</i> | 1 | 20 | | | | | | | | 1 | 40 | | 2 | 50 | |
| | <i>Penaeus penicillatus</i> | 2 | 20 | | 5 | 50 | | 3 | 20 | | 4 | 40 | | 4 | 50 | |
| | <i>Penaeus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Penaeus semisulcatus</i> | | | | 1 | 20 | | 1 | 10 | | | | | | | |
| | <i>Solenocera crassicornis</i> | | | | | | | | | | | | | | | |
| | <i>Trachypenaeus curvirostris</i> | | | | | | | | | | | | | | | |
| Crab | <i>Charybdis affinis</i> | 47 | 720 | | 79 | 1250 | | 71 | 970 | | 61 | 970 | | 90 | 1160 | |
| | <i>Charybdis annulata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis bimaculata</i> | | | | | | | | | | 1 | 10 | | | | |
| | <i>Charybdis cruciata</i> | 1 | 70 | | 1 | 50 | | | | | 2 | 20 | | 1 | 20 | |
| | <i>Charybdis japonica</i> | | | | | | | | | | | | | | | |

Trawling Data in July 2019

| Group | Species Name | TSB | | | | | | | | | | | | | | |
|---------------|-----------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | TSB-1 | | | TSB-2 | | | TSB-3 | | | TSB-4 | | | TSB-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>Charybdis obtusifrons</i> | | | | 1 | 40 | | 3 | 170 | | 5 | 70 | | 2 | 20 | |
| | <i>Charybdis truncata</i> | 1 | 10 | | 1 | 10 | | 2 | 10 | | 1 | 10 | | 6 | 20 | |
| | <i>Charybdis variegata</i> | | | | | | | | | | | | | | | |
| | <i>Charybdis spp.</i> | 30 | 370 | * | 30 | 360 | * | 20 | 210 | * | 30 | 370 | * | 20 | 220 | * |
| | <i>Doclea sp.</i> | | | | | | | | | | | | | | | |
| | <i>Doclea ovis</i> | | | | | | | | | | | | | | | |
| | <i>Dorippe facchino</i> | | | | | | | | | | | | | | | |
| | <i>Dorippe polita</i> | 20 | 230 | * | 20 | 240 | * | 30 | 340 | * | 30 | 340 | * | 7 | 86 | |
| | <i>Eucrate crenata</i> | | | | | | | | | | | | | | | |
| | <i>Galene bispinosa</i> | | | | | | | | | | | | | 1 | 10 | |
| | Hermit crab | 50 | 220 | * | 30 | 170 | * | 30 | 64 | * | 30 | 220 | * | 30 | 230 | * |
| | <i>Leucosia craniolaris</i> | | | | | | | 20 | 51 | * | | | | 20 | 96 | * |
| | <i>Macromedaeus distinguendus</i> | | | | | | | | | | | | | | | |
| | <i>Maorophthalmus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Matuta planipes</i> | | | | | | | | | | | | | | | |
| | <i>Ocypode ceratophthalmus</i> | | | | | | | | | | | | | | | |
| | <i>Parapanope euagora</i> | | | | | | | | | | | | | | | |
| | <i>Parthenope validus</i> | | | | | | | 20 | 210 | * | | | | 30 | 410 | * |
| | <i>Plumnus hirtellus</i> | | | | | | | | | | | | | | | |
| | <i>Porcellanella picta</i> | | | | | | | | | | | | | | | |
| | <i>Portunus hastatoides</i> | 13 | 20 | | 6 | 20 | | 22 | 50 | | 27 | 50 | | 33 | 70 | |
| | <i>Portunus pelagicus</i> | 2 | 80 | | 1 | 180 | | 2 | 30 | | | | | | | |
| | <i>Portunus sanguinolentus</i> | 1 | 40 | | | | | | | | | | | | | |
| | <i>Portunus trituberculatus</i> | | | | 1 | 26 | | 1 | 40 | | 2 | 140 | | 2 | 80 | |
| | <i>Thalamita sima</i> | 1 | 10 | | 2 | 18 | | 2 | 20 | | | | | | | |
| | <i>Typhlocarcinus sp.</i> | | | | | | | | | | | | | 1 | 4.2 | |
| Mantis shrimp | <i>Anchisquilla fasciata</i> | | | | | | | 1 | 10 | | | | | 1 | 20 | |
| | <i>Dictyosquilla foveolata</i> | | | | | | | 1 | 10 | | | | | | | |
| | <i>Harpiosquilla harpax</i> | | | | | | | | | | 4 | 200 | | 7 | 290 | |
| | <i>Oratosquilla interrupta</i> | 2 | 40 | | 1 | 20 | | 3 | 50 | | 8 | 180 | | 7 | 160 | |
| | <i>Oratosquilla oratoria</i> | | | | | | | 1 | 30 | | | | | | | |
| | <i>Oratosquilla kempfi</i> | | | | | | | | | | | | | | | |
| Gastropod | <i>Bursa rana</i> | | | | 1 | 8.2 | | | | | | | | | | |
| | <i>Calyptraea sp.</i> | | | | | | | | | | | | | | | |
| | <i>Desmaulus extintorius</i> | | | | | | | | | | | | | | | |
| | <i>Drupa musiva</i> | | | | 2 | 11 | | | | | | | | | | |
| | <i>Epitonium scalare</i> | | | | | | | | | | | | | | | |
| | <i>Inquistor flavidula</i> | | | | | | | | | | | | | | | |
| | <i>Gemmula deshavesii</i> | | | | | | | | | | | | | | | |
| | <i>Lophiotoma leucotropis</i> | | | | | | | | | | | | | | | |
| | <i>Murex trapa</i> | 160 | 1780 | * | 160 | 1610 | * | 70 | 740 | * | 120 | 1190 | * | 150 | 1580 | * |
| | <i>Nassarius hepaticus</i> | | | | | | | | | | | | | | | |
| | <i>Nassarius succinctus</i> | | | | 40 | 86 | * | 20 | 43 | * | | | | 30 | 63 | * |
| | <i>Philine sp.</i> | 20 | 13 | * | 20 | 13 | * | 6 | 2.1 | | | | | 2 | 1.6 | |
| | <i>Pilosabia pilosa</i> | | | | | | | | | | | | | | | |
| | <i>Siphopatella walshi</i> | | | | | | | 5 | 2 | | | | | | | |
| | <i>Tonna sulcosa</i> | | | | | | | | | | | | | | | |
| | <i>Turritella terebra</i> | 390 | 3780 | * | 460 | 4320 | * | 400 | 3840 | * | 560 | 5370 | * | 520 | 5040 | * |

Trawling Data in July 2019

| Group | Species Name | TSB | | | | | | | | | | | | | | |
|------------|------------------------------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|--------------|-----------------|-----------------|
| | | TSB-1 | | | TSB-2 | | | TSB-3 | | | TSB-4 | | | TSB-5 | | |
| | | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| | <i>UNID Nudibranch</i> | | | | | | | | | | | | | | | |
| Bivalve | <i>Anomia chinensis</i> | | | | | | | | | | | | | | | |
| | <i>Anomalodiscus squamosus</i> | | | | | | | 20 | 70 | * | | | | | | |
| | <i>Chione isabellina</i> | | | | | | | 20 | 130 | * | 20 | 130 | * | | | |
| | <i>Chlamys nobilis</i> | | | | | | | | | | | | | | | |
| | <i>Macoma candida</i> | | | | | | | | | | | | | | | |
| | <i>Ostrea sp.</i> | | | | | | | | | | | | | | | |
| | <i>Paphia undulata</i> | | | | | | | | | | | | | | | |
| | <i>Parahyatissa sp.</i> | | | | | | | | | | | | | | | |
| | <i>Pinctada martensii</i> | | | | | | | | | | | | | | | |
| | <i>Placuna placenta</i> | | | | | | | | | | | | | | | |
| | <i>Ruditapes philippinarum</i> | | | | | | | | | | | | | | | |
| | <i>Samacar sp.</i> | | | | | | | | | | | | | | | |
| | <i>Scapharca subcrenata</i> | | | | 1 | 51 | | | | | | | | | | |
| | <i>Tegillarca nodifera</i> | | | | | | | | | | | | | 20 | 110 | * |
| | <i>Tolonostrea talonata</i> | | | | | | | | | | | | | | | |
| | <i>Vepricardium asiaticum</i> | | | | | | | | | | | | | | | |
| Cephalopod | <i>Lolingid squid</i> | | | | 2 | 20 | | 2 | 20 | | 4 | 30 | | 1 | 10 | |
| | <i>Amphioctopus fangsiao</i> | | | | | | | | | | | | | | | |
| | <i>Sepiid cuttlefish</i> | | | | | | | | | | | | | | | |
| Echinoderm | <i>Acaudina molpadioides</i> | | | | | | | | | | 3 | 62 | | | | |
| | <i>Colochirus quadrangularis</i> | | | | | | | | | | | | | | | |
| | <i>Protankyra bidentata (c.f.)</i> | | | | | | | | | | | | | | | |
| | <i>Sivela plicata</i> | | | | | | | | | | 20 | 280 | * | | | |
| | <i>Temnopleurus toreumaticus</i> | 30 | 410 | * | 20 | 280 | * | 30 | 320 | * | 30 | 370 | * | 20 | 260 | * |
| Cnidaria | <i>Carcinactis ichikawai</i> | | | | 2 | 0.9 | | 3 | 1.2 | | 2 | 1.1 | | | | |
| | <i>Cavernularia habereri</i> | 30 | 160 | * | 30 | 160 | * | 20 | 130 | * | 5 | 36 | | 11 | 76 | |
| | <i>Comanthus japonicus</i> | | | | | | | | | | | | | | | |
| | <i>Edwardsia japonica (c.f.)</i> | | | | | | | | | | | | | | | |
| | <i>Ellisella laevis</i> | 2 | 21 | | 3 | 30 | | | | | 1 | 11 | | | | |
| | Jelly fish | | | | | | | | | | | | | | | |
| | <i>Pteroeides chinense</i> | 20 | 760 | * | 11 | 180 | | 7 | 160 | | 6 | 150 | | 5 | 110 | |
| | <i>Gorgonia spp.</i> | | | | | | | | | | | | | | | |
| | Sea anomea | | | | | | | | | | | | | | | |
| | <i>Virgularia gustaviana</i> | | | | | | | 4 | 66 | | | | | | | |
| Porifera | Spongiatia | | | | | | | | | | | | | | | |
| Cirripedia | <i>c.f. Bananus spp.</i> | | | | | | | | | | | | | | | |