Demersal Trawling Data in August 2015

| | | INA | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | INA-1 | | | INA-2 | | | INA-3 | | | INA-4 | | | INA-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| Fish | *Acentrogobius caninus* | | | | | | | | | | | | | | | |
| | *Acanthemblemaria rivasi* | | | | | | | | | | | | | | | |
| | *Apogon kiensis* | | | | 1.0 | 4.0 | | 5.0 | 42.0 | | | | | 1.0 | 2.0 | |
| | *Arius sinensis* | | | | | | | | | | | | | | | |
| | *Arnoglossus waitei* | | | | | | | | | | | | | | | |
| | *Caranx kalla* | | | | | | | | | | | | | | | |
| | *Chrysochir aureus* | | | | | | | | | | | | | | | |
| | *Collichthys lucidus* | | | | | | | | | | | | | | | |
| | *Cynoglossus abbreviatus* | | | | | | | | | | 1.0 | 18.0 | | | | |
| | *Cynoglossus arel* | 1.0 | 62.0 | | 1.0 | 16.0 | | 1.0 | 62.0 | | | | | | | |
| | *Cynoglossus puncticeps* | | | | | | | | | | | | | | | |
| | *Cynoglossus roulei* | | | | | | | | | | | | | | | |
| | *Cynoglossus semilaevis* | | | | | | | | | | | | | | | |
| | *Dasyatis akajei* | 2.0 | 182.0 | | | | | | | | | | | 1.0 | 2178.0 | |
| | *Dasyatis zugei* | | | | | | | | | | | | | | | |
| | *Dendrophysa russelii* | 5.0 | 178.0 | | 2.0 | 54.0 | | 4.0 | 52.0 | | | | | | | |
| | *Drepane punctata* | | | | | | | | | | 2.0 | 478.0 | | 1.0 | 302.0 | |
| | *Epinephelus awoara* | | | | | | | | | | | | | | | |
| | *Epinephelus bruneus* | | | | | | | | | | | | | | | |
| | *Gerres filamentosus* | 1.0 | 38.0 | | | | | | | | | | | 1.0 | 52.0 | |
| | *Grammoplites scaber* | 2.0 | 90.0 | | 4.0 | 152.0 | | | | | 1.0 | 132.0 | | | | |
| | *Gymnothorax chilospilus* | | | | | | | | | | | | | | | |
| | *Gymnura japonica* | | | | | | | | | | | | | | | |
| | *Ilisa elongata* | | | | | | | | | | | | | | | |
| | *Inegocia japonica* | | | | | | | 1.0 | 20.0 | | | | | 2.0 | 44.0 | |
| | *Inegocia japonica* | | | | | | | | | | | | | | | |
| | *Inimicus didactylus* | | | | | | | | | | | | | | | |
| | *Inimicus japonicus* | 3.0 | 463.0 | | | | | | | | 1.0 | 168.0 | | | | |
| | *Johnius belangerii* | 5.0 | 236.0 | | 4.0 | 374.0 | | 3.0 | 200.0 | | | | | | | |
| | *Johnius heterolepis* | | | | | | | | | | | | | | | |
| | *Konosirus punctatus* | | | | | | | | | | | | | | | |
| | *Lagocephalus wheeleri* | | | | | | | | | | | | | | | |
| | *Leiognathus brevirostris* | | | | | | | 2.0 | 28.0 | | 13.0 | 167.0 | | 18.0 | 198.0 | |
| | *Leiognathus nuchalis* | 7.0 | 96.0 | | 18.0 | 262.0 | | | | | 13.0 | 167.0 | | 11.0 | 118.0 | |
| | *Lepturacanthus savala* | | | | | | | 1.0 | 4.0 | | | | | | | |
| | *Liza macrolepis* | | | | | | | | | | 1.0 | 62.0 | | | | |
| | *Minous monodactylus* | | | | | | | | | | | | | | | |
| | *Monacanthus chinensis* | | | | | | | 1.0 | 84.0 | | | | | | | |
| | *Muraenesox cinereus* | 1.0 | 340.0 | | | | | | | | | | | | | |
| | *Nemipterus japonicus* | 3.0 | 48.0 | | 2.0 | 24.0 | | 3.0 | 36.0 | | | | | | | |
| | *Oxyurichthys tentacularis* | | | | | | | | | | | | | | | |
| | *Paracentropogon rubripinnis* | | | | | | | | | | | | | | | |
| | *Parachaeturichthys polynema* | | | | 1.0 | 10.0 | | | | | | | | | | |
| | *Paramonacanthus sulcatus* | | | | 2.0 | 8.0 | | 2.0 | 6.0 | | | | | | | |
| | *Parargyrops edita* | 3.0 | 56.0 | | | | | | | | | | | | | |
| | *Pennahia anea* | 1.0 | 20.0 | | 2.0 | 24.0 | | 1.0 | 10.0 | | | | | | | |
| | *Pennahia macrocephalus* | | | | | | | | | | | | | | | |
| | *Platycephalus indicus* | 1.0 | 14.0 | | | | | 2.0 | 21.0 | | | | | 1.0 | 30.0 | |
| | *Plectorhynchus pictus* | | | | | | | | | | 1.0 | 16.0 | | | | |

Demersal Trawling Data in August 2015

| | | INA | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | INA-1 | | | INA-2 | | | INA-3 | | | INA-4 | | | INA-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| | *Polydactylus sextarius* | 1.0 | 8.0 | | 1.0 | 4.0 | | 2.0 | 10.0 | | | | | 3.0 | 34.0 | |
| | *Pseudorhombus cinnamoneus* | 2.0 | 34.0 | | | | | | | | 1.0 | 10.0 | | | | |
| | *Rhabdosargus sarba* | | | | | | | | | | | | | | | |
| | *Saurida elongata* | | | | | | | | | | | | | | | |
| | *Scatophagus argus* | | | | | | | | | | | | | 1.0 | 202.0 | |
| | *Sebastiscus marmoratus* | 1.0 | 58.0 | | 1.0 | 8.0 | | | | | | | | | | |
| | *Siganus fuscescens* | | | | | | | | | | | | | | | |
| | *Sillago sihama* | 3.0 | 142.0 | | | | | | | | | | | | | |
| | *Solea ovata* | 4.0 | 44.0 | | 11.0 | 166.0 | | 1.0 | 10.0 | | 3.0 | 26.0 | | 2.0 | 18.0 | |
| | *Stolephorus insularis* | | | | | | | | | | | | | | | |
| | *Takifugu niphobles* | | | | | | | | | | | | | | | |
| | *Takifugu oblongus* | | | | | | | | | | | | | 1.0 | 120.0 | |
| | *Takifugu xanthopterus* | | | | | | | | | | | | | 1.0 | 120.0 | |
| | *Terapon jarbua* | | | | | | | | | | | | | | | |
| | *Terapon theraps* | | | | | | | | | | | | | | | |
| | *Thryssa hamiltonii* | | | | | | | | | | | | | | | |
| | *Thryssa kammalensis* | | | | | | | | | | | | | | | |
| | *Trachycephalus uranoscopa* | 3.0 | 34.0 | | 6.0 | 64.0 | | 4.0 | 30.0 | | | | | | | |
| | *Trypauchen vagina* | | | | | | | | | | | | | | | |
| | *Uroconger lepturus* | | | | | | | | | | | | | | | |
| | *Zebrias zebra* | | | | | | | | | | | | | | | |
| Prawn or shrimp | *Alpheus japonicus* | | | | | | | | | | | | | | | |
| | *Metapenaeopsis barbata* | | | | | | | 2.0 | 2.0 | | | | | | | |
| | *Metapenaeus ensis* | | | | | | | | | | | | | | | |
| | *Metapenaeus joyneri* | 1.0 | 2.0 | | | | | | | | | | | | | |
| | *Parapenaeopsis hardwickii* | | | | 2.0 | 18.0 | | | | | | | | | | |
| | *Parapenaeopsis tenella* | 1.0 | 44.0 | | | | | | | | | | | | | |
| | *Penaeus japonicus* | | | | | | | | | | | | | | | |
| | *Penaeus merguiensis* | | | | | | | | | | | | | | | |
| | *Penaeus penicillatus* | | | | | | | 1.0 | 14.0 | | 1.0 | 12.0 | | | | |
| | *Solenocera crassicornis* | | | | | | | | | | | | | | | |
| | *Trachypenaeus curvirostris* | | | | | | | | | | | | | | | |
| Crab | *Arcania heptacontra* | | | | | | | | | | | | | | | |
| | *Arcania undecimspinosa* | 30.0 | 586.0 | * | 1.0 | 14.2 | | | | | | | | 1.0 | 8.2 | |
| | *Calappa philargius* | | | | 3.0 | 142.0 | | 5.0 | 276.0 | | | | | | | |
| | *Charybdis affinis* | 13.0 | 166.0 | | 15.0 | 258.0 | | 9.0 | 120.0 | | 3.0 | 38.0 | | | | |
| | *Charybdis cruciata* | | | | | | | | | | | | | | | |
| | *Charybdis japonica* | | | | 1.0 | 40.0 | | 3.0 | 240.0 | | | | | | | |
| | *Charybdis spp.* | 33.0 | 514.0 | * | 35.0 | 435.0 | * | 40.0 | 423.0 | * | 26.0 | 280.0 | * | 30.0 | 216.0 | * |
| | *Dorippe polita* | 38.0 | 560.0 | * | 25.0 | 259.0 | * | 28.0 | 285.0 | * | 21.0 | 261.0 | * | 25.0 | 267.0 | * |
| | *Eucrate crenata* | | | | | | | | | | | | | 3.0 | 21.7 | |
| | *Galene bispinosa* | | | | 4.0 | 186.0 | | 1.0 | 92.0 | | 20.0 | 1640.0 | * | 2.0 | 120.0 | |
| | Hermit crab | 45.0 | 145.0 | * | 40.0 | 134.0 | * | 40.0 | 180.0 | * | 72.0 | 290.0 | * | 22.0 | 120.0 | * |
| | *Leucosia craniolaris* | 10.0 | 31.0 | | 12.0 | 36.0 | | 15.0 | 44.0 | | 13.0 | 41.0 | | 25.0 | 73.0 | * |
| | *Macrophthalmus japonicus* | | | | | | | | | | | | | | | |
| | *Parapanope euagora* | | | | 1.0 | 26.1 | | | | | | | | | | |
| | *Parthenope validus* | 25.0 | 172.0 | * | 30.0 | 263.0 | * | 22.0 | 165.0 | * | 25.0 | 179.0 | * | 30.0 | 293.0 | * |
| | *Philyra sp.* | | | | 2.0 | 88.2 | | | | | | | | | | |
| | *Porcellanella picta* | 40.0 | 18.2 | * | 30.0 | 12.2 | * | 30.0 | 12.2 | * | 40.0 | 18.6 | * | 30.0 | 14.2 | * |

Demersal Trawling Data in August 2015

| | | INA | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | INA-1 | | | INA-2 | | | INA-3 | | | INA-4 | | | INA-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| | *Portunus pelagicus* | | | | 4.0 | 682.0 | | | | | 1.0 | 226.0 | | | | |
| | *Portunus sanguinolentus* | | | | 1.0 | 114.0 | | | | | | | | | | |
| | *Portunus trituberculatus* | 11.0 | 1216.0 | | 7.0 | 1018.0 | | | | | 3.0 | 468.0 | | | | |
| | *Scylla serrata (mud crab)* | | | | | | | | | | | | | | | |
| | *Tachypleus tridentatus* | | | | | | | | | | | | | | | |
| | *Typhlocarcinus sp.* | | | | 1.0 | 0.8 | | | | | | | | | | |
| Mantis shrimp | *Harpiosquilla harpax* | 1.0 | 64.0 | | 1.0 | 26.0 | | 3.0 | 162.0 | | | | | 1.0 | 46.0 | |
| | *Erugosquilla woodmasoni* | | | | | | | | | | | | | | | |
| | *Oratosquilla kempi* | | | | | | | | | | | | | | | |
| | *Oratosquilla interrupta* | 4.0 | 106.0 | | 13.0 | 236.0 | | 4.0 | 100.0 | | | | | | | |
| | *Oratosquilla oratoria* | 2.0 | 28.0 | | 3.0 | 24.0 | | 1.0 | 6.0 | | | | | | | |
| Gastropod | *Apollon olivator rubustus* | | | | | | | | | | | | | | | |
| | *Bursa rana* | | | | | | | | | | | | | | | |
| | *Calyptraea sp.* | 20.0 | 9.2 | * | 5.0 | 2.1 | | | | | 2.0 | 0.8 | | | | |
| | *Drupa musiva* | 1.0 | 4.1 | | | | | | | | 6.0 | 47.0 | | | | |
| | *Epitonium scalare* | | | | | | | | | | | | | | | |
| | *Inquistor flavidula* | | | | 20.0 | 38.4 | * | | | | | | | | | |
| | *Lophiotoma leucotropis* | | | | | | | | | | | | | | | |
| | *Murex trapa* | 20.0 | 103.0 | | 15.0 | 65.0 | | 24.0 | 126.0 | * | 15.0 | 73.0 | | 16.0 | 67.0 | |
| | *Nassarius hepaticus* | | | | | | | | | | | | | | | |
| | *Nassarius succinctus* | 7.0 | 16.0 | | | | | 30.0 | 62.0 | * | | | | | | |
| | *Natica tigrina* | | | | | | | | | | | | | | | |
| | *Philine sp.* | 20.0 | 210.0 | * | | | | 5.0 | 31.0 | | 3.0 | 12.0 | | 30.0 | 260.0 | * |
| | *Pilosabia pilosa* | 30.0 | 11.8 | * | | | | | | | 2.0 | 0.8 | | | | |
| | *Siphopatella walshi* | | | | 20.0 | 9.2 | * | | | | 5.0 | 2.3 | | | | |
| | *Tonna sulcosa* | | | | | | | | | | | | | | | |
| | *Turritella terebra* | 220.0 | 2100.0 | * | 190.0 | 1720.0 | * | 240.0 | 2350.0 | * | 180.0 | 1730.0 | * | 210.0 | 2030.0 | * |
| Bivalve | Bivalve Juvenil | | | | | | | | | | | | | | | |
| | *Chione isabellina* | | | | | | | | | | | | | | | |
| | *Chlamys nobilis* | 30.0 | 288.0 | * | 20.0 | 210.0 | * | 1.0 | 13.4 | | 2.0 | 16.2 | | | | |
| | *Macoma candida* | | | | | | | | | | | | | | | |
| | *Ostrea sp.* | 20.0 | 36.0 | * | 70.0 | 160.0 | * | 90.0 | 54.0 | * | 120.0 | 186.0 | * | 30.0 | 48.0 | * |
| | *Paphia undulata* | 40.0 | 350.0 | * | | | | 20.0 | 120.0 | * | 30.0 | 260.0 | * | 2.0 | 7.5 | |
| | *Perna viridis* | | | | | | | | | | | | | | | |
| | *Pinctada martensii* | | | | | | | | | | | | | | | |
| | *Placuna placenta* | | | | | | | | | | | | | | | |
| | *Ruditapes philippinarum* | | | | 3.0 | 21.0 | | | | | | | | | | |
| | *Scapharca subcrenata* | | | | | | | | | | | | | | | |
| | *Spondylus sp.* | | | | | | | | | | 1.0 | 20.4 | | | | |
| | *Tegillarca nodifera* | | | | | | | | | | | | | | | |
| | *Vepricardium asiaticum* | | | | | | | | | | | | | | | |
| | *Tolonostrea talonata* | | | | | | | | | | | | | 1.0 | 3.8 | |
| Cephalopod | *Lolingid squid* | | | | | | | 1.0 | 24.0 | | | | | | | |
| | *Sepiid cuttlefish* | | | | | | | | | | | | | | | |
| | *Octopus sp.* | 2.0 | 136.0 | | 1.0 | 18.0 | | 2.0 | 92.0 | | | | | | | |
| Echinoderm | *Acaudina molpadioides* | 4.0 | 141.0 | | 2.0 | 43.2 | | 3.0 | 110.0 | | | | | | | |
| | *Asteroidea* | | | | | | | | | | | | | | | |
| | Brittle star | | | | | | | | | | | | | | | |
| | *Pseudocolochirus sp.* | | | | | | | | | | | | | | | |

Demersal Trawling Data in August 2015

| | | INA | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | INA-1 | | | INA-2 | | | INA-3 | | | INA-4 | | | INA-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| | *Temnopleurus toreumaticus* | 48.0 | 550.0 | * | 25.0 | 497.0 | * | 18.0 | 216.0 | | 46.0 | 770.0 | * | 30.0 | 325.0 | * |
| Cnidaria | *Edwardsia japonica (c.f.)* | | | | | | | | | | | | | | | |
| | *Ellisella laevis* | 5.0 | 290.0 | | 2.0 | 28.4 | | 1.0 | 11.0 | | | | | | | |
| | *Gorgonia spp.* | 2.0 | 64.0 | | 3.0 | 260.0 | | | | | | | | | | |
| | *Carcinactis ichikawai* | 5.0 | 1.1 | | 3.0 | 0.6 | | 3.0 | 0.7 | | 16.0 | 8.2 | | 8.0 | 4.5 | |
| | *Cavernularia habereri* | 30.0 | 140.0 | * | 40.0 | 180.0 | * | 45.0 | 241.0 | * | 30.0 | 150.0 | * | 30.0 | 140.0 | * |
| | Jelly fish | | | | | | | | | | | | | | | |
| | *Pteroeides chinense* | 115.0 | 1660.0 | * | 200.0 | 2900.0 | * | 118.0 | 2193.0 | * | 60.0 | 1420.0 | * | 126.0 | 2270.0 | * |
| | Sea anomea | | | | 2.0 | 18.3 | | 2.0 | 7.4 | | | | | | | |
| | *Virgularia gustaviana* | 22.0 | 210.0 | * | 22.0 | 215.0 | * | 10.0 | 95.0 | | 4.0 | 46.0 | | 3.0 | 27.0 | |
| Urochordata | Sea Squirts | 2.0 | 11.4 | | 1.0 | 4.3 | | | | | | | | | | |
| Cirripedia | *Bananus spp.(c.f.)* | 80.0 | 42.0 | * | 90.0 | 46.0 | * | 80.0 | 42.0 | * | 30.0 | 31.0 | * | 40.0 | 35.0 | * |

Demersal Trawling Data in August 2015

| | | INB | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | INB-1 | | | INB-2 | | | INB-3 | | | INB-4 | | | INB-5 | | |
| Group | *Species Name* | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| Fish | *Acentrogobius caninus* | | | | | | | | | | 3.0 | 52.0 | | 3.0 | 52.0 | |
| | *Acanthemblemaria rivasi* | | | | | | | 1.0 | 4.0 | | | | | | | |
| | *Apogon kiensis* | 1.0 | 4.0 | | 3.0 | 16.0 | | 4.0 | 40.0 | | 1.0 | 4.0 | | | | |
| | *Arius sinensis* | | | | | | | | | | | | | | | |
| | *Arnoglossus waitei* | 1.0 | 8.0 | | | | | | | | 2.0 | 8.0 | | 6.0 | 26.0 | |
| | *Caranx kalla* | | | | | | | | | | | | | | | |
| | *Chrysochir aureus* | | | | | | | | | | | | | | | |
| | *Collichthys lucidus* | | | | | | | | | | | | | | | |
| | *Cynoglossus abbreviatus* | | | | | | | | | | | | | | | |
| | *Cynoglossus arel* | 1.0 | 16.0 | | 3.0 | 192.0 | | 1.0 | 20.0 | | 8.0 | 224.0 | | 3.0 | 130.0 | |
| | *Cynoglossus puncticeps* | | | | | | | | | | | | | | | |
| | *Cynoglossus roulei* | | | | | | | | | | | | | | | |
| | *Cynoglossus semilaevis* | | | | | | | | | | | | | | | |
| | *Dasyatis akajei* | 3.0 | 1364.0 | | 2.0 | 1054.0 | | | | | 4.0 | 2110.0 | | 2.0 | 1040.0 | |
| | *Dasyatis zugei* | 7.0 | 1281.0 | | 2.0 | 658.0 | | 2.0 | 614.0 | | 3.0 | 330.0 | | 2.0 | 94.0 | |
| | *Dendrophysa russelii* | 2.0 | 60.0 | | | | | 3.0 | 142.0 | | 1.0 | 62.0 | | 1.0 | 56.0 | |
| | *Drepane punctata* | 2.0 | 428.0 | | | | | | | | | | | | | |
| | *Epinephelus awoara* | | | | | | | | | | | | | 1.0 | 44.0 | |
| | *Epinephelus bruneus* | | | | | | | | | | 1.0 | 32.0 | | | | |
| | *Gerres filamentosus* | | | | | | | | | | 3.0 | 46.0 | | 1.0 | 54.0 | |
| | *Grammoplites scaber* | 1.0 | 16.0 | | 1.0 | 46.0 | | 1.0 | 130.0 | | | | | 2.0 | 28.0 | |
| | *Gymnothorax chilospilus* | | | | | | | | | | | | | | | |
| | *Gymnura japonica* | | | | | | | | | | 5.0 | 3640.0 | | 3.0 | 1940.0 | |
| | *Ilisa elongata* | | | | | | | | | | | | | | | |
| | *Inegocia japonica* | | | | 2.0 | 50.0 | | 3.0 | 36.0 | | 4.0 | 126.0 | | 2.0 | 124.0 | |
| | *Inegocia japonica* | | | | | | | | | | | | | | | |
| | *Inimicus didactylus* | | | | | | | 1.0 | 180.0 | | | | | | | |
| | *Inimicus japonicus* | | | | | | | | | | | | | | | |
| | *Johnius belangerii* | | | | | | | 2.0 | 68.0 | | 1.0 | 42.0 | | | | |
| | *Johnius heterolepis* | | | | | | | | | | | | | | | |
| | *Konosirus punctatus* | | | | | | | | | | | | | | | |
| | *Lagocephalus wheeleri* | | | | | | | | | | | | | | | |
| | *Leiognathus brevirostris* | 14.0 | 150.0 | | 10.0 | 105.0 | | 9.0 | 102.0 | | 16.0 | 150.0 | | 16.0 | 141.0 | |
| | *Leiognathus nuchalis* | 48.0 | 510.0 | | 188.0 | 1949.0 | | 214.0 | 2490.0 | | 84.0 | 790.0 | | 111.0 | 980.0 | |
| | *Lepturacanthus savala* | | | | | | | | | | | | | | | |
| | *Liza macrolepis* | | | | | | | | | | | | | | | |
| | *Minous monodactylus* | | | | | | | | | | | | | | | |
| | *Monacanthus chinensis* | | | | | | | | | | | | | | | |
| | *Muraenesox cinereus* | | | | 1.0 | 210.0 | | | | | | | | 1.0 | 64.0 | |
| | *Nemipterus japonicus* | 1.0 | 24.0 | | 1.0 | 12.0 | | 1.0 | 10.0 | | | | | | | |
| | *Oxyurichthys tentacularis* | | | | | | | | | | 1.0 | 6.0 | | 1.0 | 10.0 | |
| | *Paracentropogon rubripinnis* | | | | | | | | | | | | | | | |
| | *Parachaeturichthys polynema* | | | | | | | | | | | | | | | |
| | *Paramonacanthus sulcatus* | | | | | | | 1.0 | 6.0 | | | | | | | |
| | *Parargyrops edita* | | | | | | | | | | | | | | | |
| | *Pennahia anea* | 6.0 | 172.0 | | 21.0 | 654.0 | | 19.0 | 562.0 | | 2.0 | 94.0 | | | | |
| | *Pennahia macrocephalus* | | | | | | | | | | | | | | | |
| | *Platycephalus indicus* | 1.0 | 16.0 | | 5.0 | 876.0 | | 4.0 | 584.0 | | 2.0 | 96.0 | | 5.0 | 210.0 | |
| | *Plectorhynchus pictus* | | | | | | | | | | | | | | | |

Demersal Trawling Data in August 2015

| | | INB | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | INB-1 | | | INB-2 | | | INB-3 | | | INB-4 | | | INB-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| | *Polydactylus sextarius* | 1.0 | 18.0 | | 2.0 | 46.0 | | 23.0 | 384.0 | | 1.0 | 14.0 | | | | |
| | *Pseudorhombus cinnamoneus* | 2.0 | 74.0 | | 3.0 | 62.0 | | 1.0 | 30.0 | | 4.0 | 66.0 | | 2.0 | 44.0 | |
| | *Rhabdosargus sarba* | | | | | | | | | | | | | 1.0 | 134.0 | |
| | *Saurida elongata* | | | | | | | | | | | | | | | |
| | *Scatophagus argus* | | | | | | | | | | | | | | | |
| | *Sebastiscus marmoratus* | | | | | | | 1.0 | 18.0 | | | | | | | |
| | *Siganus fuscescens* | | | | | | | 2.0 | 88.0 | | 2.0 | 112.0 | | | | |
| | *Sillago sihama* | | | | | | | | | | | | | | | |
| | *Solea ovata* | | | | | | | 4.0 | 48.0 | | 5.0 | 54.0 | | 4.0 | 48.0 | |
| | *Stolephorus insularis* | | | | | | | | | | | | | | | |
| | *Takifugu niphobles* | | | | | | | | | | | | | | | |
| | *Takifugu oblongus* | | | | | | | | | | | | | | | |
| | *Takifugu xanthopterus* | | | | | | | | | | | | | | | |
| | *Terapon jarbua* | | | | | | | | | | | | | 1.0 | 14.0 | |
| | *Terapon theraps* | | | | | | | | | | | | | | | |
| | *Thryssa hamiltonii* | | | | | | | | | | | | | | | |
| | *Thryssa kammalensis* | | | | | | | | | | | | | | | |
| | *Trachycephalus uranoscopa* | 1.0 | 12.0 | | 9.0 | 148.0 | | 1.0 | 10.0 | | 8.0 | 152.0 | | 2.0 | 36.0 | |
| | *Trypauchen vagina* | | | | 2.0 | 30.0 | | | | | 1.0 | 8.0 | | | | |
| | *Uroconger lepturus* | | | | | | | | | | | | | | | |
| | *Zebrias zebra* | | | | | | | | | | | | | | | |
| Prawn or shrimp | *Alpheus japonicus* | | | | | | | | | | | | | | | |
| | *Metapenaeopsis barbata* | | | | | | | | | | | | | | | |
| | *Metapenaeus ensis* | | | | 4.0 | 30.0 | | 2.0 | 14.0 | | 3.0 | 32.0 | | | | |
| | *Metapenaeus joyneri* | 1.0 | 6.0 | | | | | | | | | | | | | |
| | *Parapenaeopsis hardwickii* | | | | | | | | | | | | | | | |
| | *Parapenaeopsis tenella* | | | | | | | | | | | | | | | |
| | *Penaeus japonicus* | | | | | | | | | | | | | 1.0 | 18.0 | |
| | *Penaeus merguiensis* | | | | | | | | | | | | | | | |
| | *Penaeus penicillatus* | | | | | | | | | | | | | | | |
| | *Solenocera crassicornis* | | | | | | | | | | | | | | | |
| | *Trachypenaeus curvirostris* | | | | | | | | | | | | | | | |
| Crab | *Arcania heptacontra* | | | | | | | | | | | | | | | |
| | *Arcania undecimspinosa* | | | | 1.0 | 7.4 | | | | | | | | | | |
| | *Calappa philargius* | | | | | | | | | | | | | | | |
| | *Charybdis affinis* | 8.0 | 142.0 | | 63.0 | 954.0 | | 27.0 | 400.0 | | 5.0 | 112.0 | | 10.0 | 228.0 | |
| | *Charybdis cruciata* | 1.0 | 306.0 | | | | | | | | 1.0 | 226.0 | | | | |
| | *Charybdis japonica* | | | | | | | | | | | | | | | |
| | *Charybdis spp.* | 42.0 | 568.0 | * | 80.0 | 1260.0 | * | 50.0 | 630.0 | * | 60.0 | 640.0 | * | 73.0 | 790.0 | * |
| | *Dorippe polita* | 40.0 | 480.0 | * | 80.0 | 1480.0 | * | 50.0 | 630.0 | * | 70.0 | 660.0 | * | 60.0 | 590.0 | * |
| | *Eucrate crenata* | | | | 1.0 | 4.8 | | 4.0 | 21.2 | | | | | 8.0 | 56.0 | |
| | *Galene bispinosa* | 2.0 | 184.0 | | | | | | | | 30.0 | 1190.0 | * | 3.0 | 182.0 | |
| | Hermit crab | 70.0 | 380.0 | * | 50.0 | 308.0 | * | 60.0 | 310.0 | * | 50.0 | 210.0 | * | 40.0 | 190.0 | * |
| | *Leucosia craniolaris* | 11.0 | 54.0 | | | | | 2.0 | 7.8 | | 30.0 | 160.0 | * | 20.0 | 120.0 | * |
| | *Macrophthalmus japonicus* | 20.0 | 170.0 | * | | | | | | | | | | | | |
| | *Parapanope euagora* | | | | | | | | | | | | | | | |
| | *Parthenope validus* | 30.0 | 460.0 | * | 43.0 | 770.0 | * | 11.0 | 94.0 | | 20.0 | 210.0 | * | 30.0 | 360.0 | * |
| | *Philyra sp.* | | | | | | | | | | | | | | | |
| | *Porcellanella picta* | 36.0 | 13.1 | * | 30.0 | 11.6 | * | 40.0 | 21.2 | * | 20.0 | 10.0 | * | 26.0 | 11.0 | * |

Demersal Trawling Data in August 2015

| | | INB | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | INB-1 | | | INB-2 | | | INB-3 | | | INB-4 | | | INB-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| | *Portunus pelagicus* | | | | | | | 3.0 | 472.0 | | 7.0 | 1354.0 | | 5.0 | 794.0 | |
| | *Portunus sanguinolentus* | | | | | | | | | | 1.0 | 4.0 | | | | |
| | *Portunus trituberculatus* | 1.0 | 86.0 | | 1.0 | 168.0 | | 11.0 | 1344.0 | | 8.0 | 956.0 | | 9.0 | 990.0 | |
| | *Scylla serrata (mud crab)* | | | | | | | | | | | | | | | |
| | *Tachypleus tridentatus* | 1.0 | 43.0 | | | | | | | | | | | | | |
| | *Typhlocarcinus sp.* | | | | | | | | | | | | | | | |
| Mantis shrimp | *Harpiosquilla harpax* | 4.0 | 148.0 | | 10.0 | 482.0 | | 5.0 | 268.0 | | 4.0 | 218.0 | | 7.0 | 330.0 | |
| | *Erugosquilla woodmasoni* | | | | | | | | | | | | | | | |
| | *Oratosquilla kempi* | | | | | | | | | | | | | | | |
| | *Oratosquilla interrupta* | 6.0 | 158.0 | | 8.0 | 220.0 | | 15.0 | 420.0 | | 10.0 | 286.0 | | 4.0 | 122.0 | |
| | *Oratosquilla oratoria* | | | | 1.0 | 12.0 | | | | | 1.0 | 14.0 | | 1.0 | 22.0 | |
| Gastropod | *Apollon olivator rubustus* | | | | | | | | | | | | | | | |
| | *Bursa rana* | | | | 1.0 | 6.2 | | | | | 1.0 | 17.5 | | | | |
| | *Calyptraea sp.* | 5.0 | 2.3 | | | | | 3.0 | 1.2 | | | | | 1.0 | 0.3 | |
| | *Drupa musiva* | | | | | | | 3.0 | 11.4 | | | | | | | |
| | *Epitonium scalare* | | | | | | | | | | | | | | | |
| | *Inquistor flavidula* | | | | 30.0 | 68.0 | * | 11.0 | 23.0 | | | | | | | |
| | *Lophiotoma leucotropis* | | | | | | | | | | | | | | | |
| | *Murex trapa* | 60.0 | 370.0 | * | 40.0 | 210.0 | * | 50.0 | 290.0 | * | 40.0 | 302.0 | * | 30.0 | 260.0 | * |
| | *Nassarius hepaticus* | | | | | | | | | | | | | | | |
| | *Nassarius succinctus* | 40.0 | 76.0 | * | 30.0 | 52.0 | * | 30.0 | 52.0 | * | | | | | | |
| | *Natica tigrina* | | | | | | | | | | | | | | | |
| | *Philine sp.* | | | | 5.0 | 37.0 | | 40.0 | 370.0 | * | 50.0 | 420.0 | * | 50.0 | 420.0 | * |
| | *Pilosabia pilosa* | 1.0 | 0.4 | | | | | | | | | | | 1.0 | 0.2 | |
| | *Siphopatella walshi* | | | | | | | 1.0 | 0.3 | | | | | | | |
| | *Tonna sulcosa* | | | | | | | | | | | | | | | |
| | *Turritella terebra* | 3360.0 | 31000.0 | * | 2680.0 | 24160.0 | * | 2170.0 | 20810.0 | * | 420.0 | 4040.0 | * | 390.0 | 3760.0 | * |
| Bivalve | Bivalve Juvenil | | | | | | | | | | | | | | | |
| | *Chione isabellina* | | | | | | | | | | 1.0 | 3.6 | | 3.0 | 19.6 | |
| | *Chlamys nobilis* | 1.0 | 4.3 | | 20.0 | 290.0 | * | | | | | | | | | |
| | *Macoma candida* | | | | | | | | | | 2.0 | 12.0 | | | | |
| | *Ostrea sp.* | 30.0 | 52.0 | * | 40.0 | 66.0 | * | 20.0 | 38.0 | * | 50.0 | 36.0 | * | 30.0 | 48.0 | * |
| | *Paphia undulata* | 40.0 | 330.0 | * | 9.0 | 56.0 | | 130.0 | 1240.0 | * | 90.0 | 760.0 | * | 110.0 | 1040.0 | * |
| | *Perna viridis* | | | | | | | | | | | | | | | |
| | *Pinctada martensii* | | | | | | | | | | | | | | | |
| | *Placuna placenta* | | | | | | | | | | 1.0 | 8.6 | | | | |
| | *Ruditapes philippinarum* | | | | | | | | | | | | | | | |
| | *Scapharca subcrenata* | 3.0 | 112.0 | | | | | | | | | | | | | |
| | *Spondylus sp.* | | | | | | | | | | | | | | | |
| | *Tegillarca nodifera* | | | | | | | | | | | | | | | |
| | *Vepricardium asiaticum* | | | | 2.0 | 41.8 | | | | | 1.0 | 21.3 | | | | |
| | *Tolonostrea talonata* | | | | 1.0 | 2.2 | | | | | | | | | | |
| Cephalopod | *Lolingid squid* | | | | 1.0 | 40.0 | | | | | | | | | | |
| | *Sepiid cuttlefish* | | | | 1.0 | 6.0 | | 3.0 | 228.0 | | 4.0 | 294.0 | | 2.0 | 44.0 | |
| | *Octopus sp.* | | | | | | | 2.0 | 312.0 | | | | | | | |
| Echinoderm | *Acaudina molpadioides* | 20.0 | 610.0 | * | 30.0 | 1240.0 | * | 20.0 | 720.0 | * | 30.0 | 910.0 | * | 26.0 | 880.0 | * |
| | *Asteroidea* | | | | | | | | | | | | | 1.0 | 19.3 | |
| | Brittle star | | | | | | | | | | | | | | | |
| | *Pseudocolochirus sp.* | | | | | | | 1.0 | 317.0 | | | | | | | |

Demersal Trawling Data in August 2015

| | | INB | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | INB-1 | | | INB-2 | | | INB-3 | | | INB-4 | | | INB-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| | *Temnopleurus toreumaticus* | 18.0 | 164.0 | | 16.0 | 94.0 | | 40.0 | 830.0 | * | 70.0 | 690.0 | * | 60.0 | 610.0 | * |
| Cnidaria | *Edwardsia japonica (c.f.)* | | | | | | | | | | | | | | | |
| | *Ellisella laevis* | 2.0 | 26.2 | | 1.0 | 18.2 | | 4.0 | 360.0 | | 3.0 | 36.0 | | 2.0 | 22.0 | |
| | *Gorgonia spp.* | | | | | | | 3.0 | 210.0 | | 1.0 | 12.0 | | | | |
| | *Carcinactis ichikawai* | 16.0 | 8.1 | | 42.0 | 16.1 | * | 18.0 | 9.9 | | 20.0 | 11.0 | * | 20.0 | 12.0 | * |
| | *Cavernularia habereri* | 60.0 | 260.0 | * | 50.0 | 220.0 | * | 90.0 | 410.0 | * | 50.0 | 210.0 | * | 60.0 | 240.0 | * |
| | Jelly fish | | | | | | | | | | | | | | | |
| | *Pteroeides chinense* | 1980.0 | 21000.0 | * | 2130.0 | 26000.0 | * | 2410.0 | 31000.0 | * | 70.0 | 2100.0 | * | 80.0 | 2300.0 | * |
| | Sea anomea | | | | 1.0 | 6.2 | | 2.0 | 16.0 | | | | | 1.0 | 7.2 | |
| | *Virgularia gustaviana* | 5.0 | 172.0 | | 8.0 | 141.0 | | 20.0 | 206.0 | * | 6.0 | 46.0 | | 7.0 | 55.0 | |
| Urochordata | Sea Squirts | | | | | | | | | | | | | | | |
| Cirripedia | *Bananus spp.(c.f.)* | 70.0 | 38.0 | * | 120.0 | 61.0 | * | 90.0 | 47.0 | * | 110.0 | 62.0 | * | 120.0 | 63.0 | * |

Demersal Trawling Data in August 2015

| | | TNA | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | TNA-1 | | | TNA-2 | | | TNA-3 | | | TNA-4 | | | TNA-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| Fish | *Acentrogobius caninus* | | | | | | | | | | | | | | | |
| | *Acanthemblemaria rivasi* | | | | | | | | | | | | | | | |
| | *Apogon kiensis* | 3.0 | 14.0 | | | | | 2.0 | 18.0 | | 1.0 | 10.0 | | | | |
| | *Arius sinensis* | | | | | | | | | | | | | | | |
| | *Arnoglossus waitei* | | | | | | | | | | | | | | | |
| | *Caranx kalla* | | | | | | | | | | | | | | | |
| | *Chrysochir aureus* | | | | | | | | | | | | | | | |
| | *Collichthys lucidus* | | | | 1.0 | 12.0 | | 1.0 | 18.0 | | 5.0 | 82.0 | | 8.0 | 92.0 | |
| | *Cynoglossus abbreviatus* | | | | | | | | | | | | | | | |
| | *Cynoglossus arel* | 1.0 | 4.0 | | | | | 9.0 | 54.0 | | 3.0 | 40.0 | | 8.0 | 136.0 | |
| | *Cynoglossus puncticeps* | | | | | | | | | | | | | 1.0 | 4.0 | |
| | *Cynoglossus roulei* | 3.0 | 218.0 | | | | | 6.0 | 64.0 | | 2.0 | 36.0 | | 2.0 | 42.0 | |
| | *Cynoglossus semilaevis* | | | | | | | | | | | | | | | |
| | *Dasyatis akajei* | | | | | | | | | | 2.0 | 264.0 | | 1.0 | 100.0 | |
| | *Dasyatis zugei* | | | | | | | | | | | | | | | |
| | *Dendrophysa russelii* | | | | 9.0 | 116.0 | | | | | 1.0 | 26.0 | | 1.0 | 20.0 | |
| | *Drepane punctata* | | | | | | | | | | | | | | | |
| | *Epinephelus awoara* | | | | | | | | | | | | | | | |
| | *Epinephelus bruneus* | | | | | | | | | | | | | | | |
| | *Gerres filamentosus* | | | | | | | | | | | | | | | |
| | *Grammoplites scaber* | 2.0 | 66.0 | | 1.0 | 60.0 | | 23.0 | 888.0 | | 8.0 | 532.0 | | 17.0 | 524.0 | |
| | *Gymnothorax chilospilus* | | | | | | | | | | | | | | | |
| | *Gymnura japonica* | | | | | | | | | | | | | | | |
| | *Ilisa elongata* | | | | | | | | | | | | | | | |
| | *Inegocia japonica* | | | | | | | | | | | | | | | |
| | *Inegocia japonica* | | | | | | | | | | | | | | | |
| | *Inimicus didactylus* | | | | | | | | | | | | | | | |
| | *Inimicus japonicus* | | | | | | | | | | | | | | | |
| | *Johnius belangerii* | 15.0 | 196.0 | | | | | 72.0 | 1504.0 | | 44.0 | 754.0 | | 44.0 | 952.0 | |
| | *Johnius heterolepis* | | | | | | | | | | | | | | | |
| | *Konosirus punctatus* | | | | | | | | | | | | | | | |
| | *Lagocephalus wheeleri* | | | | | | | | | | | | | | | |
| | *Leiognathus brevirostris* | | | | | | | | | | 7.0 | 72.0 | | 9.0 | 120.0 | |
| | *Leiognathus nuchalis* | | | | 3.0 | 26.0 | | | | | | | | | | |
| | *Lepturacanthus savala* | | | | | | | | | | | | | | | |
| | *Liza macrolepis* | | | | | | | | | | | | | | | |
| | *Minous monodactylus* | 1.0 | 24.0 | | | | | | | | | | | | | |
| | *Monacanthus chinensis* | | | | | | | | | | | | | | | |
| | *Muraenesox cinereus* | | | | | | | 1.0 | 172.0 | | | | | | | |
| | *Nemipterus japonicus* | | | | | | | | | | | | | | | |
| | *Oxyurichthys tentacularis* | | | | | | | | | | | | | | | |
| | *Paracentropogon rubripinnis* | | | | | | | | | | | | | | | |
| | *Parachaeturichthys polynema* | 2.0 | 16.0 | | | | | | | | | | | 1.0 | 10.0 | |
| | *Paramonacanthus sulcatus* | 2.0 | 8.0 | | 2.0 | 10.0 | | 1.0 | 4.0 | | 1.0 | 4.0 | | | | |
| | *Parargyrops edita* | | | | | | | | | | | | | | | |
| | *Pennahia anea* | | | | | | | | | | | | | | | |
| | *Pennahia macrocephalus* | | | | | | | | | | | | | | | |
| | *Platycephalus indicus* | 1.0 | 12.0 | | 1.0 | 14.0 | | 2.0 | 120.0 | | | | | | | |
| | *Plectorhynchus pictus* | | | | | | | | | | | | | | | |

Demersal Trawling Data in August 2015

| | | TNA | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | TNA-1 | | | TNA-2 | | | TNA-3 | | | TNA-4 | | | TNA-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| | *Polydactylus sextarius* | 1.0 | 16.0 | | 2.0 | 28.0 | | 1.0 | 6.0 | | 3.0 | 48.0 | | | | |
| | *Pseudorhombus cinnamoneus* | 1.0 | 22.0 | | | | | | | | | | | | | |
| | *Rhabdosargus sarba* | | | | | | | | | | | | | | | |
| | *Saurida elongata* | | | | | | | | | | | | | | | |
| | *Scatophagus argus* | | | | | | | | | | | | | | | |
| | *Sebastiscus marmoratus* | | | | | | | | | | | | | | | |
| | *Siganus fuscescens* | | | | | | | | | | | | | | | |
| | *Sillago sihama* | | | | | | | | | | | | | | | |
| | *Solea ovata* | | | | | | | 24.0 | 196.0 | | 45.0 | 366.0 | | 9.0 | 370.0 | |
| | *Stolephorus insularis* | | | | | | | | | | | | | | | |
| | *Takifugu niphobles* | | | | | | | | | | | | | | | |
| | *Takifugu oblongus* | | | | | | | | | | | | | | | |
| | *Takifugu xanthopterus* | | | | | | | | | | | | | | | |
| | *Terapon jarbua* | | | | | | | | | | | | | | | |
| | *Terapon theraps* | | | | | | | | | | | | | | | |
| | *Thryssa hamiltonii* | | | | | | | | | | | | | | | |
| | *Thryssa kammalensis* | | | | | | | | | | | | | | | |
| | *Trachycephalus uranoscopa* | | | | | | | | | | | | | | | |
| | *Trypauchen vagina* | | | | | | | 2.0 | 42.0 | | 1.0 | 20.0 | | 2.0 | 20.0 | |
| | *Uroconger lepturus* | | | | | | | | | | | | | | | |
| | *Zebrias zebra* | 1.0 | 18.0 | | | | | | | | | | | | | |
| Prawn or shrimp | *Alpheus japonicus* | | | | | | | | | | | | | | | |
| | *Metapenaeopsis barbata* | | | | | | | 1.0 | 4.0 | | | | | 1.0 | 6.0 | |
| | *Metapenaeus ensis* | 69.0 | 698.0 | | 52.0 | 538.0 | | 200.0 | 2240.0 | | 187.0 | 2264.0 | | 153.0 | 3158.0 | |
| | *Metapenaeus joyneri* | 11.0 | 58.0 | | 2.0 | 8.0 | | 35.0 | 180.0 | | | | | 5.0 | 16.0 | |
| | *Parapenaeopsis hardwickii* | 5.0 | 52.0 | | 7.0 | 64.0 | | 2.0 | 26.0 | | 1.0 | 12.0 | | 5.0 | 60.0 | |
| | *Parapenaeopsis tenella* | 40.0 | 20.0 | | 4.0 | 16.0 | | 1.0 | 10.0 | | 4.0 | 104.0 | | 3.0 | 48.0 | |
| | *Penaeus japonicus* | | | | | | | 4.0 | 56.0 | | | | | 1.0 | 12.0 | |
| | *Penaeus merguiensis* | | | | 2.0 | 36.0 | | 2.0 | 28.0 | | 4.0 | 76.0 | | | | |
| | *Penaeus penicillatus* | 1.0 | 14.0 | | | | | 10.0 | 20.0 | | 4.0 | 44.0 | | 3.0 | 38.0 | |
| | *Solenocera crassicornis* | 1.0 | 6.0 | | 2.0 | 8.0 | | | | | 1.0 | 4.0 | | 1.0 | 2.0 | |
| | *Trachypenaeus curvirostris* | 3.0 | 6.0 | | 1.0 | 4.0 | | 8.0 | 30.0 | | 17.0 | 78.0 | | 5.0 | 26.0 | |
| Crab | *Arcania heptacontra* | | | | 1.0 | 6.4 | | | | | | | | | | |
| | *Arcania undecimspinosa* | | | | | | | | | | | | | | | |
| | *Calappa philargius* | | | | | | | | | | | | | | | |
| | *Charybdis affinis* | 4.0 | 66.0 | | 59.0 | 564.0 | | 58.0 | 682.0 | | 35.0 | 426.0 | | 15.0 | 158.0 | |
| | *Charybdis cruciata* | 1.0 | 2.0 | | | | | | | | | | | 1.0 | 206.0 | |
| | *Charybdis japonica* | | | | | | | | | | 1.0 | 24.0 | | 1.0 | 12.0 | |
| | *Charybdis spp.* | 70.0 | 760.0 | * | 83.0 | 1102.0 | * | 143.0 | 1818.0 | * | 84.0 | 1094.0 | * | 153.0 | 2012.0 | * |
| | *Dorippe polita* | 60.0 | 570.0 | * | 40.0 | 410.0 | * | 70.0 | 730.0 | * | 50.0 | 504.0 | * | 80.0 | 930.0 | * |
| | *Eucrate crenata* | | | | 3.0 | 12.4 | | | | | | | | | | |
| | *Galene bispinosa* | | | | | | | | | | | | | | | |
| | Hermit crab | 50.0 | 140.0 | * | 40.0 | 110.0 | * | 42.0 | 116.0 | * | 50.0 | 130.0 | * | 70.0 | 190.0 | * |
| | *Leucosia craniolaris* | | | | 1.0 | 3.9 | | | | | | | | | | |
| | *Macrophthalmus japonicus* | | | | | | | | | | | | | | | |
| | *Parapanope euagora* | | | | | | | | | | | | | | | |
| | *Parthenope validus* | 7.0 | 90.0 | | 30.0 | 480.0 | * | 20.0 | 360.0 | * | | | | | | |
| | *Philyra sp.* | | | | | | | | | | | | | | | |
| | *Porcellanella picta* | | | | 2.0 | 0.4 | | | | | | | | | | |

Demersal Trawling Data in August 2015

| | | TNA | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | TNA-1 | | | TNA-2 | | | TNA-3 | | | TNA-4 | | | TNA-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| | *Portunus pelagicus* | | | | | | | 1.0 | 54.0 | | | | | | | |
| | *Portunus sanguinolentus* | | | | | | | | | | | | | | | |
| | *Portunus trituberculatus* | 1.0 | 78.0 | | 1.0 | 52.0 | | 8.0 | 784.0 | | 4.0 | 414.0 | | | | |
| | *Scylla serrata (mud crab)* | | | | | | | | | | | | | | | |
| | *Tachypleus tridentatus* | | | | | | | | | | | | | | | |
| | *Typhlocarcinus sp.* | | | | | | | 1.0 | 1.1 | | | | | | | |
| Mantis shrimp | *Harpiosquilla harpax* | 1.0 | 30.0 | | 1.0 | 52.0 | | | | | | | | 1.0 | 50.0 | |
| | *Erugosquilla woodmasoni* | | | | | | | | | | | | | | | |
| | *Oratosquilla kempi* | 1.0 | 4.0 | | | | | 1.0 | 34.0 | | 1.0 | 34.0 | | | | |
| | *Oratosquilla interrupta* | 13.0 | 268.0 | | 10.0 | 276.0 | | 5.0 | 114.0 | | 14.0 | 344.0 | | 9.0 | 228.0 | |
| | *Oratosquilla oratoria* | 2.0 | 20.0 | | 3.0 | 34.0 | | 7.0 | 84.0 | | 8.0 | 120.0 | | 13.0 | 166.0 | |
| Gastropod | *Apollon olivator rubustus* | | | | 1.0 | 4.2 | | | | | | | | | | |
| | *Bursa rana* | | | | | | | | | | | | | | | |
| | *Calyptraea sp.* | 1.0 | 0.4 | | 2.0 | 0.8 | | | | | | | | | | |
| | *Drupa musiva* | | | | | | | | | | | | | | | |
| | *Epitonium scalare* | | | | | | | | | | | | | | | |
| | *Inquistor flavidula* | | | | 20.0 | 38.0 | * | | | | | | | | | |
| | *Lophiotoma leucotropis* | | | | 1.0 | 4.2 | | | | | | | | | | |
| | *Murex trapa* | 50.0 | 320.0 | * | 40.0 | 290.0 | * | 40.0 | 320.0 | * | 60.0 | 420.0 | * | 50.0 | 340.0 | * |
| | *Nassarius hepaticus* | | | | | | | | | | 7.0 | 8.2 | | | | |
| | *Nassarius succinctus* | | | | 30.0 | 63.0 | * | | | | 5.0 | 11.2 | | 20.0 | 42.0 | * |
| | *Natica tigrina* | | | | | | | | | | | | | 1.0 | 2.4 | |
| | *Philine sp.* | 30.0 | 260.0 | * | | | | 20.0 | 264.0 | * | 5.0 | 38.0 | | | | |
| | *Pilosabia pilosa* | 1.0 | 0.2 | | | | | | | | | | | | | |
| | *Siphopatella walshi* | 3.0 | 0.8 | | | | | | | | | | | | | |
| | *Tonna sulcosa* | | | | | | | | | | | | | | | |
| | *Turritella terebra* | 420.0 | 4160 | * | 610.0 | 6030.0 | * | 590.0 | 5740.0 | * | 720.0 | 7170.0 | * | 710.0 | 6980.0 | * |
| Bivalve | Bivalve Juvenil | | | | | | | | | | | | | | | |
| | *Chione isabellina* | | | | | | | | | | | | | | | |
| | *Chlamys nobilis* | 1.0 | 4.2 | | | | | | | | | | | | | |
| | *Macoma candida* | | | | | | | | | | | | | | | |
| | *Ostrea sp.* | 40.0 | 32.0 | * | 50.0 | 41.0 | * | | | | 12.0 | 8.2 | | 60.0 | 72.0 | * |
| | *Paphia undulata* | | | | | | | | | | | | | | | |
| | *Perna viridis* | | | | | | | | | | | | | | | |
| | *Pinctada martensii* | | | | | | | | | | | | | | | |
| | *Placuna placenta* | | | | | | | | | | | | | | | |
| | *Ruditapes philippinarum* | | | | 2.0 | 7.2 | | 20.0 | 110.0 | * | | | | 1.0 | 4.1 | |
| | *Scapharca subcrenata* | | | | 2.0 | 192.0 | | | | | 1.0 | 42.0 | | | | |
| | *Spondylus sp.* | | | | | | | | | | | | | | | |
| | *Tegillarca nodifera* | | | | | | | | | | | | | | | |
| | *Vepricardium asiaticum* | | | | | | | | | | | | | | | |
| | *Tolonostrea talonata* | | | | | | | | | | | | | | | |
| Cephalopod | *Lolingid squid* | | | | | | | | | | | | | | | |
| | *Sepiid cuttlefish* | | | | | | | | | | 1.0 | 76.0 | | | | |
| | *Octopus sp.* | | | | | | | 5.0 | 328.0 | | 1.0 | 70.0 | | | | |
| Echinoderm | *Acaudina molpadioides* | 30.0 | 1510.0 | * | 20.0 | 890.0 | * | 30.0 | 840.0 | * | 40.0 | 720.0 | * | 20.0 | 810.0 | * |
| | *Asteroidea* | | | | | | | | | | | | | | | |
| | Brittle star | | | | | | | | | | | | | | | |
| | *Pseudocolochirus sp.* | | | | | | | | | | | | | | | |

Demersal Trawling Data in August 2015

| | | TNA | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | TNA-1 | | | TNA-2 | | | TNA-3 | | | TNA-4 | | | TNA-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| | *Temnopleurus toreumaticus* | 40.0 | 630.0 | * | 30.0 | 510.0 | * | 70.0 | 780.0 | * | 90.0 | 1260.0 | * | 50.0 | 720.0 | * |
| Cnidaria | *Edwardsia japonica (c.f.)* | | | | | | | | | | | | | | | |
| | *Ellisella laevis* | | | | 4.0 | 141.0 | | | | | 1.0 | 22.0 | | | | |
| | *Gorgonia spp.* | | | | 1.0 | 12.0 | | 1.0 | 14.2 | | | | | 1.0 | 17.0 | |
| | *Carcinactis ichikawai* | 40.0 | 21.2 | * | 30.0 | 14.2 | * | 50.0 | 26.2 | * | 8.0 | 4.2 | | 11.0 | 6.1 | |
| | *Cavernularia habereri* | 20.0 | 110.0 | * | 30.0 | 140.0 | * | 7.0 | 31.0 | | 60.0 | 440.0 | * | 19.0 | 110.0 | |
| | Jelly fish | 1.0 | 210.0 | | | | | | | | | | | | | |
| | *Pteroeides chinense* | 6.0 | 130.0 | | 20.0 | 480.0 | * | 20.0 | 280.0 | * | 11.0 | 260.0 | | 2.0 | 38.0 | |
| | Sea anomea | 4.0 | 63.0 | | 2.0 | 16.3 | | | | | | | | 3.0 | 21.0 | |
| | *Virgularia gustaviana* | 1.0 | 11.2 | | | | | 2.0 | 18.4 | | 3.0 | 41.2 | | | | |
| Urochordata | Sea Squirts | 1.0 | 21.0 | | | | | | | | | | | | | |
| Cirripedia | *Bananus spp.(c.f.)* | 130.0 | 64.0 | * | 90.0 | 42.0 | * | 120.0 | 58.0 | * | 110.0 | 63.0 | * | 80.0 | 36.0 | * |

Demersal Trawling Data in August 2015

| | | TNB | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | TNB-1 | | | TNB-2 | | | TNB-3 | | | TNB-4 | | | TNB-5 | | |
| Group | *Species Name* | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| Fish | *Acentrogobius caninus* | | | | | | | | | | | | | | | |
| | *Acanthemblemaria rivasi* | | | | | | | | | | | | | | | |
| | *Apogon kiensis* | 1.0 | 4.0 | | 1.0 | 6.0 | | | | | | | | | | |
| | *Arius sinensis* | | | | | | | | | | | | | | | |
| | *Arnoglossus waitei* | | | | 3.0 | 28.0 | | 2.0 | 20.0 | | | | | | | |
| | *Caranx kalla* | | | | | | | | | | | | | | | |
| | *Chrysochir aureus* | | | | | | | | | | 2.0 | 758.0 | | | | |
| | *Collichthys lucidus* | | | | | | | | | | | | | | | |
| | *Cynoglossus abbreviatus* | | | | | | | | | | | | | | | |
| | *Cynoglossus arel* | 7.0 | 240.0 | | 6.0 | 140.0 | | | | | 16.0 | 318.0 | | 1.0 | 18.0 | |
| | *Cynoglossus puncticeps* | 1.0 | 10.0 | | | | | | | | | | | | | |
| | *Cynoglossus roulei* | | | | | | | | | | | | | 1.0 | 180.0 | |
| | *Cynoglossus semilaevis* | | | | | | | | | | | | | | | |
| | *Dasyatis akajei* | | | | | | | 4.0 | 1224.0 | | | | | | | |
| | *Dasyatis zugei* | 4.0 | 222.0 | | 1.0 | 276.0 | | | | | 8.0 | 545.0 | | 1.0 | 66.0 | |
| | *Dendrophysa russelii* | | | | 1.0 | 16.0 | | 3.0 | 98.0 | | | | | | | |
| | *Drepane punctata* | | | | | | | | | | | | | | | |
| | *Epinephelus awoara* | | | | | | | | | | | | | | | |
| | *Epinephelus bruneus* | | | | | | | | | | | | | | | |
| | *Gerres filamentosus* | | | | | | | | | | | | | | | |
| | *Grammoplites scaber* | | | | | | | | | | | | | | | |
| | *Gymnothorax chilospilus* | | | | | | | | | | | | | | | |
| | *Gymnura japonica* | | | | | | | | | | | | | | | |
| | *Ilisa elongata* | | | | | | | | | | | | | | | |
| | *Inegocia japonica* | | | | | | | | | | | | | | | |
| | *Inegocia japonica* | | | | | | | | | | | | | | | |
| | *Inimicus didactylus* | | | | | | | | | | | | | | | |
| | *Inimicus japonicus* | | | | | | | | | | | | | | | |
| | *Johnius belangerii* | | | | | | | | | | 1.0 | 6.0 | | | | |
| | *Johnius heterolepis* | | | | | | | | | | | | | | | |
| | *Konosirus punctatus* | | | | 2.0 | 146.0 | | | | | | | | | | |
| | *Lagocephalus wheeleri* | | | | | | | | | | 2.0 | 126.0 | | | | |
| | *Leiognathus brevirostris* | 17.0 | 172.0 | | 17.0 | 175.0 | | 9.0 | 114.0 | | 21.0 | 186.0 | | 20.0 | 188.0 | |
| | *Leiognathus nuchalis* | 42.0 | 420.0 | | 38.0 | 391.0 | | | | | 17.0 | 182.0 | | 7.0 | 66.0 | |
| | *Lepturacanthus savala* | | | | | | | | | | | | | | | |
| | *Liza macrolepis* | 10.0 | 156.0 | | 3.0 | 106.0 | | 1.0 | 16.0 | | | | | | | |
| | *Minous monodactylus* | | | | | | | | | | | | | | | |
| | *Monacanthus chinensis* | | | | | | | | | | | | | | | |
| | *Muraenesox cinereus* | 1.0 | 250.0 | | | | | | | | | | | 1.0 | 196.0 | |
| | *Nemipterus japonicus* | | | | 1.0 | 14.0 | | | | | | | | | | |
| | *Oxyurichthys tentacularis* | | | | | | | 2.0 | 18.0 | | | | | | | |
| | *Paracentropogon rubripinnis* | 2.0 | 12.0 | | 1.0 | 6.0 | | | | | 1.0 | 10.0 | | | | |
| | *Parachaeturichthys polynema* | | | | | | | 1.0 | 10.0 | | | | | | | |
| | *Paramonacanthus sulcatus* | | | | | | | | | | 3.0 | 22.0 | | | | |
| | *Parargyrops edita* | | | | | | | | | | | | | | | |
| | *Pennahia anea* | 1.0 | 12.0 | | | | | 1.0 | 24.0 | | | | | | | |
| | *Pennahia macrocephalus* | | | | | | | | | | | | | | | |
| | *Platycephalus indicus* | 16.0 | 792.0 | | 6.0 | 236.0 | | 9.0 | 228.0 | | 10.0 | 174.0 | | 7.0 | 374.0 | |
| | *Plectorhynchus pictus* | | | | | | | | | | | | | | | |

Demersal Trawling Data in August 2015

| | | TNB | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | TNB-1 | | | TNB-2 | | | TNB-3 | | | TNB-4 | | | TNB-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| | *Polydactylus sextarius* | 7.0 | 106.0 | | 11.0 | 182.0 | | 9.0 | 130.0 | | 12.0 | 180.0 | | 3.0 | 34.0 | |
| | *Pseudorhombus cinnamoneus* | | | | | | | 1.0 | 14.0 | | | | | | | |
| | *Rhabdosargus sarba* | | | | | | | | | | | | | | | |
| | *Saurida elongata* | | | | | | | | | | | | | | | |
| | *Scatophagus argus* | | | | | | | | | | | | | | | |
| | *Sebastiscus marmoratus* | | | | | | | | | | | | | | | |
| | *Siganus fuscescens* | | | | | | | | | | | | | | | |
| | *Sillago sihama* | | | | | | | | | | | | | | | |
| | *Solea ovata* | | | | | | | 1.0 | 4.0 | | 3.0 | 34.0 | | 1.0 | 10.0 | |
| | *Stolephorus insularis* | | | | | | | | | | | | | | | |
| | *Takifugu niphobles* | | | | | | | | | | | | | | | |
| | *Takifugu oblongus* | | | | | | | | | | | | | | | |
| | *Takifugu xanthopterus* | | | | | | | | | | | | | | | |
| | *Terapon jarbua* | | | | | | | | | | | | | | | |
| | *Terapon theraps* | | | | | | | | | | | | | | | |
| | *Thryssa hamiltonii* | | | | | | | | | | | | | | | |
| | *Thryssa kammalensis* | | | | | | | | | | | | | | | |
| | *Trachycephalus uranoscopa* | | | | 1.0 | 14.0 | | | | | | | | | | |
| | *Trypauchen vagina* | 4.0 | 18.0 | | 1.0 | 6.0 | | 5.0 | 36.0 | | | | | | | |
| | *Uroconger lepturus* | | | | | | | | | | | | | | | |
| | *Zebrias zebra* | | | | | | | | | | | | | | | |
| Prawn or shrimp | *Alpheus japonicus* | | | | | | | | | | | | | | | |
| | *Metapenaeopsis barbata* | | | | | | | | | | | | | | | |
| | *Metapenaeus ensis* | 29.0 | 342.0 | | 18.0 | 152.0 | | 21.0 | 238.0 | | 43.0 | 434.0 | | 29.0 | 264.0 | |
| | *Metapenaeus joyneri* | | | | | | | | | | 14.0 | 82.0 | | | | |
| | *Parapenaeopsis hardwickii* | | | | 3.0 | 16.0 | | 1.0 | 6.0 | | | | | | | |
| | *Parapenaeopsis tenella* | 1.0 | 8.0 | | | | | | | | 8.0 | 66.0 | | 3.0 | 14.0 | |
| | *Penaeus japonicus* | | | | | | | | | | | | | | | |
| | *Penaeus merguiensis* | 7.0 | 94.0 | | | | | | | | 10.0 | 190.0 | | 1.0 | 14.0 | |
| | *Penaeus penicillatus* | 13.0 | 178.0 | | 19.0 | 244.0 | | 18.0 | 216.0 | | 14.0 | 208.0 | | 1.0 | 10.0 | |
| | *Solenocera crassicornis* | | | | | | | | | | | | | | | |
| | *Trachypenaeus curvirostris* | | | | | | | | | | | | | | | |
| Crab | *Arcania heptacontra* | | | | | | | | | | | | | | | |
| | *Arcania undecimspinosa* | | | | | | | | | | | | | | | |
| | *Calappa philargius* | | | | | | | | | | | | | | | |
| | *Charybdis affinis* | 16.0 | 184.0 | | 72.0 | 762.0 | | 19.0 | 556.0 | | 12.0 | 142.0 | | 30.0 | 392.0 | |
| | *Charybdis cruciata* | | | | 1.0 | 2.0 | | | | | | | | | | |
| | *Charybdis japonica* | | | | | | | | | | | | | | | |
| | *Charybdis spp.* | 160.0 | 2130.0 | * | 170.0 | 2490.0 | * | 90.0 | 1210.0 | * | 111.0 | 1676.0 | * | 171.0 | 2476.0 | * |
| | *Dorippe polita* | 120.0 | 1360.0 | * | 80.0 | 920.0 | * | 80.0 | 920.0 | * | 90.0 | 1170.0 | * | 80.0 | 930.0 | * |
| | *Eucrate crenata* | | | | | | | | | | | | | 11.0 | 68.0 | |
| | *Galene bispinosa* | | | | | | | | | | | | | | | |
| | Hermit crab | 70.0 | 190.0 | * | 60.0 | 140.0 | * | 60.0 | 140.0 | * | 50.0 | 160.0 | * | 30.0 | 160.0 | * |
| | *Leucosia craniolaris* | 1.0 | 4.1 | | | | | 30.0 | 146.0 | * | 2.0 | 7.2 | | | | |
| | *Macrophthalmus japonicus* | | | | | | | 1.0 | 8.1 | | 2.0 | 11.2 | | | | |
| | *Parapanope euagora* | | | | | | | | | | | | | | | |
| | *Parthenope validus* | | | | | | | | | | | | | | | |
| | *Philyra sp.* | | | | | | | | | | | | | | | |
| | *Porcellanella picta* | | | | 1.0 | 0.2 | | | | | | | | | | |

Demersal Trawling Data in August 2015

| | | TNB | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | TNB-1 | | | TNB-2 | | | TNB-3 | | | TNB-4 | | | TNB-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| | *Portunus pelagicus* | | | | 1.0 | 30.0 | | | | | | | | | | |
| | *Portunus sanguinolentus* | | | | | | | | | | | | | | | |
| | *Portunus trituberculatus* | | | | 2.0 | 182.0 | | 1.0 | 168.0 | | 1.0 | 98.0 | | 3.0 | 208.0 | |
| | *Scylla serrata (mud crab)* | | | | | | | 1.0 | 106.0 | | | | | | | |
| | *Tachypleus tridentatus* | | | | | | | | | | | | | | | |
| | *Typhlocarcinus sp.* | | | | | | | | | | | | | | | |
| Mantis shrimp | *Harpiosquilla harpax* | 2.0 | 90.0 | | | | | | | | 2.0 | 136.0 | | 1.0 | 68.0 | |
| | *Erugosquilla woodmasoni* | | | | | | | | | | 1.0 | 20.0 | | | | |
| | *Oratosquilla kempi* | | | | | | | | | | | | | | | |
| | *Oratosquilla interrupta* | 3.0 | 56.0 | | | | | 5.0 | 96.0 | | 4.0 | 104.0 | | | | |
| | *Oratosquilla oratoria* | 2.0 | 30.0 | | | | | | | | | | | | | |
| Gastropod | *Apollon olivator rubustus* | | | | | | | | | | | | | | | |
| | *Bursa rana* | 2.0 | 23.4 | | | | | | | | | | | | | |
| | *Calyptraea sp.* | | | | | | | 4.0 | 1.6 | | | | | | | |
| | *Drupa musiva* | | | | | | | | | | 2.0 | 8.2 | | | | |
| | *Epitonium scalare* | | | | | | | | | | | | | | | |
| | *Inquistor flavidula* | | | | | | | | | | | | | | | |
| | *Lophiotoma leucotropis* | | | | | | | | | | | | | | | |
| | *Murex trapa* | 60.0 | 370.0 | * | 50.0 | 350.0 | * | 40.0 | 310.0 | * | 50.0 | 350.0 | * | 40.0 | 310.0 | * |
| | *Nassarius hepaticus* | | | | | | | | | | | | | | | |
| | *Nassarius succinctus* | | | | | | | | | | | | | 50.0 | 110.0 | * |
| | *Natica tigrina* | | | | | | | | | | | | | | | |
| | *Philine sp.* | | | | 20.0 | 180.0 | * | 7.0 | 48.0 | | | | | | | |
| | *Pilosabia pilosa* | | | | | | | | | | | | | | | |
| | *Siphopatella walshi* | 2.0 | 0.8 | | | | | 20.0 | 9.4 | * | | | | 1.0 | 0.3 | |
| | *Tonna sulcosa* | | | | | | | | | | | | | | | |
| | *Turritella terebra* | 710.0 | 6950.0 | * | 820.0 | 8140.0 | * | 720.0 | 6980.0 | * | 640.0 | 6310.0 | * | 610.0 | 5840.0 | * |
| Bivalve | Bivalve Juvenil | | | | | | | | | | | | | | | |
| | *Chione isabellina* | | | | | | | | | | | | | | | |
| | *Chlamys nobilis* | 2.0 | 16.2 | | 1.0 | 8.1 | | | | | | | | 3.0 | 21.2 | |
| | *Macoma candida* | | | | | | | 5.0 | 41.0 | | | | | | | |
| | *Ostrea sp.* | 40.0 | 38.0 | * | | | | 30.0 | 31.0 | * | | | | | | |
| | *Paphia undulata* | | | | | | | | | | | | | | | |
| | *Perna viridis* | | | | | | | | | | | | | | | |
| | *Pinctada martensii* | | | | | | | | | | | | | | | |
| | *Placuna placenta* | | | | | | | | | | | | | | | |
| | *Ruditapes philippinarum* | | | | 7.0 | 29.0 | | | | | | | | | | |
| | *Scapharca subcrenata* | 9.0 | 1310.0 | | 1.0 | 132.0 | | 3.0 | 304.0 | | 4.0 | 530.0 | | | | |
| | *Spondylus sp.* | | | | | | | | | | | | | | | |
| | *Tegillarca nodifera* | | | | 4.0 | 18.0 | | 2.0 | 8.4 | | | | | | | |
| | *Vepricardium asiaticum* | | | | | | | | | | | | | | | |
| | *Tolonostrea talonata* | | | | | | | | | | | | | | | |
| Cephalopod | *Lolingid squid* | | | | | | | | | | 1.0 | 24.0 | | | | |
| | *Sepiid cuttlefish* | | | | 1.0 | 248.0 | | | | | | | | | | |
| | *Octopus sp.* | | | | | | | | | | 1.0 | 46.0 | | | | |
| Echinoderm | *Acaudina molpadioides* | 0.0 | 870.0 | * | 28.0 | 1210.0 | * | 5.0 | 270.0 | | 30.0 | 1460.0 | * | 2.0 | 41.0 | |
| | *Asteroidea* | | | | | | | | | | | | | | | |
| | Brittle star | | | | | | | | | | | | | | | |
| | *Pseudocolochirus sp.* | | | | | | | | | | | | | | | |

Demersal Trawling Data in August 2015

| | | TNB | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | TNB-1 | | | TNB-2 | | | TNB-3 | | | TNB-4 | | | TNB-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| | *Temnopleurus toreumaticus* | 80.0 | 1410.0 | * | 130.0 | 1880.0 | * | 260.0 | 3940.0 | * | 250.0 | 3870.0 | * | 310.0 | 4660.0 | * |
| Cnidaria | *Edwardsia japonica (c.f.)* | | | | | | | | | | | | | | | |
| | *Ellisella laevis* | | | | 1.0 | 11.2 | | | | | | | | | | |
| | *Gorgonia spp.* | 2.0 | 62.0 | | 4.0 | 210.0 | | | | | | | | 1.0 | 14.0 | |
| | *Carcinactis ichikawai* | 20.0 | 11.2 | * | 30.0 | 14.6 | * | 30.0 | 15.0 | * | 40.0 | 19.0 | * | 30.0 | 16.2 | * |
| | *Cavernularia habereri* | 30.0 | 150.0 | * | 40.0 | 170.0 | * | 30.0 | 160.0 | * | 50.0 | 210.0 | * | 50.0 | 210.0 | * |
| | Jelly fish | | | | | | | 1.0 | 340.0 | | | | | | | |
| | *Pteroeides chinense* | 3.0 | 72.0 | | 4.0 | 82.0 | | 3.0 | 70.0 | | | | | | | |
| | Sea anomea | | | | 2.0 | 13.8 | | 1.0 | 7.2 | | | | | | | |
| | *Virgularia gustaviana* | 1.0 | 11.2 | | | | | | | | 2.0 | 26.0 | | | | |
| Urochordata | Sea Squirts | | | | | | | | | | | | | | | |
| Cirripedia | *Bananus spp.(c.f.)* | 90.0 | 53.0 | * | 110.0 | 68.0 | * | 140.0 | 82.0 | * | 40.0 | 26.0 | * | 50.0 | 24.0 | * |

Demersal Trawling Data in August 2015

| | | TSA | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | TSA-1 | | | TSA-2 | | | TSA-3 | | | TSA-4 | | | TSA-5 | | |
| Group | *Species Name* | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) | Total number | Total weight(g) | Subsampling (*) |
| Fish | *Acentrogobius caninus* | | | | | | | | | | 1.0 | 2.0 | | | | |
| | *Acanthemblemaria rivasi* | | | | | | | | | | | | | | | |
| | *Apogon kiensis* | | | | | | | | | | | | | | | |
| | *Arius sinensis* | 41.0 | 1560.0 | | 91.0 | 3108.0 | | 59.0 | 2022.0 | | 27.0 | 1038.0 | | 36.0 | 1592.0 | |
| | *Arnoglossus waitei* | | | | | | | | | | | | | | | |
| | *Caranx kalla* | | | | | | | | | | 1.0 | 10.0 | | 1.0 | 12.0 | |
| | *Chrysochir aureus* | 1.0 | 444.0 | | | | | | | | 1.0 | 352.0 | | 1.0 | 388.0 | |
| | *Collichthys lucidus* | | | | | | | 8.0 | 110.0 | | 9.0 | 114.0 | | 8.0 | 80.0 | |
| | *Cynoglossus abbreviatus* | | | | | | | | | | | | | | | |
| | *Cynoglossus arel* | | | | | | | | | | 2.0 | 96.0 | | 3.0 | 38.0 | |
| | *Cynoglossus puncticeps* | | | | | | | | | | | | | | | |
| | *Cynoglossus roulei* | 1.0 | 138.0 | | 1.0 | 156.0 | | 2.0 | 302.0 | | 1.0 | 130.0 | | 1.0 | 118.0 | |
| | *Cynoglossus semilaevis* | 10.0 | 1000.0 | | | | | | | | | | | | | |
| | *Dasyatis akajei* | | | | 2.0 | 206.0 | | 1.0 | 440.0 | | | | | | | |
| | *Dasyatis zugei* | | | | | | | | | | 1.0 | 436.0 | | 1.0 | 274.0 | |
| | *Dendrophysa russelii* | 1.0 | 28.0 | | 2.0 | 52.0 | | 6.0 | 92.0 | | | | | | | |
| | *Drepane punctata* | | | | | | | | | | | | | | | |
| | *Epinephelus awoara* | 1.0 | 22.0 | | | | | | | | | | | | | |
| | *Epinephelus bruneus* | | | | 1.0 | 56.0 | | | | | | | | | | |
| | *Gerres filamentosus* | 24.0 | 626.0 | | 71.0 | 2032.0 | | 41.0 | 872.0 | | | | | 1.0 | 14.0 | |
| | *Grammoplites scaber* | 1.0 | 8.0 | | | | | 2.0 | 18.0 | | 2.0 | 84.0 | | 1.0 | 68.0 | |
| | *Gymnothorax chilospilus* | | | | | | | | | | | | | 1.0 | 138.0 | |
| | *Gymnura japonica* | 2.0 | 1850.0 | | | | | | | | | | | | | |
| | *Ilisa elongata* | | | | 1.0 | 24.0 | | | | | | | | | | |
| | *Inegocia japonica* | | | | | | | | | | | | | | | |
| | *Inegocia japonica* | | | | 3.0 | 34.0 | | | | | | | | | | |
| | *Inimicus didactylus* | | | | | | | | | | | | | | | |
| | *Inimicus japonicus* | | | | | | | | | | | | | | | |
| | *Johnius belangerii* | 6.0 | 232.0 | | 6.0 | 170.0 | | 13.0 | 322.0 | | 6.0 | 110.0 | | 1.0 | 18.0 | |
| | *Johnius heterolepis* | | | | | | | | | | | | | | | |
| | *Konosirus punctatus* | | | | 1.0 | 42.0 | | | | | | | | 1.0 | 40.0 | |
| | *Lagocephalus wheeleri* | | | | | | | | | | | | | 3.0 | 114.0 | |
| | *Leiognathus brevirostris* | 77.0 | 568.0 | | 200.0 | 1500.0 | | 75.0 | 660.0 | | 45.0 | 390.0 | | 87.0 | 716.0 | |
| | *Leiognathus nuchalis* | 49.0 | 360.0 | | 30.0 | 372.0 | | 32.0 | 288.0 | | 128.0 | 1174.0 | | 74.0 | 676.0 | |
| | *Lepturacanthus savala* | | | | | | | | | | | | | | | |
| | *Liza macrolepis* | 3.0 | 162.0 | | 18.0 | 936.0 | | | | | 12.0 | 676.0 | | 1.0 | 46.0 | |
| | *Minous monodactylus* | | | | | | | | | | | | | | | |
| | *Monacanthus chinensis* | | | | | | | | | | | | | | | |
| | *Muraenesox cinereus* | 2.0 | 186.0 | | | | | 1.0 | 108.0 | | | | | | | |
| | *Nemipterus japonicus* | 4.0 | 54.0 | | 3.0 | 42.0 | | 2.0 | 40.0 | | | | | 1.0 | 16.0 | |
| | *Oxyurichthys tentacularis* | | | | | | | | | | | | | | | |
| | *Paracentropogon rubripinnis* | | | | | | | | | | | | | | | |
| | *Parachaeturichthys polynema* | | | | | | | | | | | | | | | |
| | *Paramonacanthus sulcatus* | 6.0 | 28.0 | | 11.0 | 54.0 | | 9.0 | 48.0 | | 2.0 | 10.0 | | 1.0 | 8.0 | |
| | *Parargyrops edita* | | | | | | | | | | | | | | | |
| | *Pennahia anea* | 3.0 | 90.0 | | 8.0 | 216.0 | | 11.0 | 202.0 | | 2.0 | 44.0 | | 2.0 | 32.0 | |
| | *Pennahia macrocephalus* | | | | | | | | | | | | | | | |
| | *Platycephalus indicus* | 2.0 | 138.0 | | 8.0 | 460.0 | | 6.0 | 226.0 | | 4.0 | 58.0 | | 1.0 | 114.0 | |
| | *Plectorhynchus pictus* | | | | | | | | | | | | | | | |

Demersal Trawling Data in August 2015

| | | TSA | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | TSA-1 | | | TSA-2 | | | TSA-3 | | | TSA-4 | | | TSA-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| | *Polydactylus sextarius* | 5.0 | 100.0 | | 11.0 | 252.0 | | 13.0 | 278.0 | | 1.0 | 22.0 | | 2.0 | 42.0 | |
| | *Pseudorhombus cinnamoneus* | | | | 1.0 | 18.0 | | 2.0 | 38.0 | | | | | | | |
| | *Rhabdosargus sarba* | | | | | | | | | | | | | | | |
| | *Saurida elongata* | 3.0 | 234.0 | | 3.0 | 314.0 | | | | | | | | | | |
| | *Scatophagus argus* | | | | | | | | | | | | | | | |
| | *Sebastiscus marmoratus* | | | | | | | | | | | | | | | |
| | *Siganus fuscescens* | | | | | | | | | | | | | | | |
| | *Sillago sihama* | | | | 3.0 | 130.0 | | 30.0 | 156.0 | | 4.0 | 178.0 | | 2.0 | 110.0 | |
| | *Solea ovata* | 32.0 | 308.0 | | 46.0 | 470.0 | | 25.0 | 278.0 | | 12.0 | 70.0 | | 2.0 | 20.0 | |
| | *Stolephorus insularis* | | | | | | | | | | 4.0 | 24.0 | | 6.0 | 34.0 | |
| | *Takifugu niphobles* | 1.0 | 218.0 | | | | | | | | | | | | | |
| | *Takifugu oblongus* | | | | | | | | | | | | | | | |
| | *Takifugu xanthopterus* | | | | | | | | | | | | | | | |
| | *Terapon jarbua* | | | | 1.0 | 12.0 | | 2.0 | 20.0 | | | | | | | |
| | *Terapon theraps* | | | | 1.0 | 78.0 | | | | | | | | 4.0 | 96.0 | |
| | *Thryssa hamiltonii* | | | | | | | 1.0 | 14.0 | | | | | | | |
| | *Thryssa kammalensis* | 1.0 | 8.0 | | 1.0 | 8.0 | | 1.0 | 12.0 | | | | | | | |
| | *Trachycephalus uranoscopa* | | | | | | | | | | | | | | | |
| | *Trypauchen vagina* | | | | 3.0 | 64.0 | | 2.0 | 22.0 | | | | | 1.0 | 20.0 | |
| | *Uroconger lepturus* | 1.0 | 8.0 | | 1.0 | 8.0 | | | | | | | | | | |
| | *Zebrias zebra* | | | | | | | | | | | | | | | |
| Prawn or shrimp | *Alpheus japonicus* | | | | 1.0 | 4.0 | | | | | | | | | | |
| | *Metapenaeopsis barbata* | | | | | | | | | | | | | | | |
| | *Metapenaeus ensis* | 3.0 | 40.0 | | 3.0 | 24.0 | | | | | 3.0 | 28.0 | | | | |
| | *Metapenaeus joyneri* | 1.0 | 8.0 | | 6.0 | 34.0 | | 6.0 | 38.0 | | 15.0 | 100.0 | | 11.0 | 70.0 | |
| | *Parapenaeopsis hardwickii* | | | | 1.0 | 6.0 | | 2.0 | 26.0 | | | | | | | |
| | *Parapenaeopsis tenella* | | | | | | | | | | 1.0 | 6.0 | | 3.0 | 22.0 | |
| | *Penaeus japonicus* | 1.0 | 18.0 | | | | | | | | | | | | | |
| | *Penaeus merguiensis* | | | | 6.0 | 114.0 | | | | | 3.0 | 70.0 | | 8.0 | 128.0 | |
| | *Penaeus penicillatus* | 3.0 | 40.0 | | 12.0 | 152.0 | | 7.0 | 110.0 | | 13.0 | 186.0 | | 1.0 | 12.0 | |
| | *Solenocera crassicornis* | | | | | | | | | | | | | | | |
| | *Trachypenaeus curvirostris* | | | | | | | | | | | | | | | |
| Crab | *Arcania heptacontra* | | | | | | | | | | | | | | | |
| | *Arcania undecimspinosa* | 5.0 | 152.0 | | 30.0 | 440.0 | * | 20.0 | 560.0 | * | | | | 1.0 | 28.4 | |
| | *Calappa philargius* | 2.0 | 94.0 | | | | | | | | | | | | | |
| | *Charybdis affinis* | 81.0 | 1008.0 | | 115.0 | 1310.0 | | 89.0 | 904.0 | | 43.0 | 576.0 | | 28.0 | 312.0 | |
| | *Charybdis cruciata* | | | | | | | | | | | | | 1.0 | 34.0 | |
| | *Charybdis japonica* | | | | | | | 2.0 | 56.0 | | | | | 1.0 | 18.0 | |
| | *Charybdis spp.* | 33.0 | 396.0 | * | 31.0 | 489.0 | * | 40.0 | 417.0 | * | 44.0 | 299.0 | * | 43.0 | 518.0 | * |
| | *Dorippe polita* | 28.0 | 275.0 | * | 25.0 | 178.0 | * | 23.0 | 266.0 | * | 18.0 | 56.0 | * | 30.0 | 108.0 | * |
| | *Eucrate crenata* | | | | | | | | | | | | | | | |
| | *Galene bispinosa* | 1.0 | 96.4 | | | | | 20.0 | 1370.0 | * | 1.0 | 98.0 | | 3.0 | 260.0 | |
| | Hermit crab | 30.0 | 110.0 | * | 36.0 | 130.0 | * | 47.0 | 420.0 | * | 21.0 | 92.0 | * | 32.0 | 120.0 | * |
| | *Leucosia craniolaris* | 15.0 | 42.0 | * | 23.0 | 168.0 | * | 16.0 | 43.0 | * | 21.0 | 62.0 | * | 18.0 | 55.0 | * |
| | *Macrophthalmus japonicus* | | | | 1.0 | 4.1 | | | | | | | | | | |
| | *Parapanope euagora* | | | | | | | | | | | | | | | |
| | *Parthenope validus* | 25.0 | 383.0 | * | 31.0 | 413.0 | * | 32.0 | 422.0 | * | 35.0 | 432.0 | * | 30.0 | 418.0 | * |
| | *Philyra sp.* | | | | | | | | | | | | | | | |
| | *Porcellanella picta* | | | | | | | | | | | | | | | |

Demersal Trawling Data in August 2015

| | | TSA | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | TSA-1 | | | TSA-2 | | | TSA-3 | | | TSA-4 | | | TSA-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| | *Portunus pelagicus* | 1.0 | 110.0 | | 2.0 | 242.0 | | | | | 7.0 | 994.0 | | | | |
| | *Portunus sanguinolentus* | 5.0 | 590.0 | | | | | | | | | | | | | |
| | *Portunus trituberculatus* | | | | 6.0 | 910.0 | | 1.0 | 62.0 | | 4.0 | 392.0 | | 6.0 | 634.0 | |
| | *Scylla serrata (mud crab)* | | | | | | | | | | | | | | | |
| | *Tachypleus tridentatus* | | | | | | | | | | | | | | | |
| | *Typhlocarcinus sp.* | | | | | | | | | | | | | | | |
| Mantis shrimp | *Harpiosquilla harpax* | | | | 1.0 | 36.0 | | | | | | | | 1.0 | 46.0 | |
| | *Erugosquilla woodmasoni* | | | | | | | | | | | | | | | |
| | *Oratosquilla kempi* | | | | | | | | | | | | | | | |
| | *Oratosquilla interrupta* | 2.0 | 40.0 | | 7.0 | 86.0 | | 4.0 | 40.0 | | 2.0 | 16.0 | | 1.0 | 6.0 | |
| | *Oratosquilla oratoria* | | | | | | | | | | | | | | | |
| Gastropod | *Apollon olivator rubustus* | | | | | | | | | | | | | | | |
| | *Bursa rana* | | | | | | | | | | | | | | | |
| | *Calyptraea sp.* | | | | 1.0 | 0.3 | | | | | 3.0 | 1.6 | | | | |
| | *Drupa musiva* | | | | | | | | | | | | | 3.0 | 11.4 | |
| | *Epitonium scalare* | 30.0 | 364.0 | * | 20.0 | 216.0 | * | | | | | | | | | |
| | *Inquistor flavidula* | | | | | | | | | | | | | | | |
| | *Lophiotoma leucotropis* | | | | | | | | | | | | | | | |
| | *Murex trapa* | 25.0 | 213.0 | * | 11.0 | 34.0 | | 6.0 | 56.0 | | 19.0 | 126.0 | | 15.0 | 125.0 | |
| | *Nassarius hepaticus* | | | | | | | | | | | | | 1.0 | 3.2 | |
| | *Nassarius succinctus* | | | | 30.0 | 62.0 | * | 20.0 | 43.0 | * | | | | | | |
| | *Natica tigrina* | | | | | | | | | | | | | | | |
| | *Philine sp.* | 46.0 | 418.0 | * | 40.0 | 315.2 | * | 40.0 | 64.3 | * | 30.0 | 260.0 | * | 15.0 | 132.0 | |
| | *Pilosabia pilosa* | | | | 3.0 | 1.1 | | | | | | | | | | |
| | *Siphopatella walshi* | | | | 4.0 | 2.2 | | 2.0 | 0.6 | | 1.0 | 0.4 | | | | |
| | *Tonna sulcosa* | | | | | | | | | | | | | | | |
| | *Turritella terebra* | 400.0 | 4200.0 | * | 330.0 | 3400.0 | * | 500.0 | 5100.0 | * | 600.0 | 6300.0 | * | 400.0 | 3700.0 | * |
| Bivalve | Bivalve Juvenil | 40.0 | 21.0 | * | 55.0 | 28.0 | * | 35.0 | 18.0 | * | 20.0 | 11.0 | * | 15.0 | 7.8 | |
| | *Chione isabellina* | | | | 1.0 | 4.4 | | | | | | | | | | |
| | *Chlamys nobilis* | 20.0 | 194.0 | * | | | | | | | | | | | | |
| | *Macoma candida* | | | | | | | | | | | | | | | |
| | *Ostrea sp.* | 40.0 | 26.0 | * | 30.0 | 42.0 | * | | | | | | | 40.0 | 64.0 | * |
| | *Paphia undulata* | 3.0 | 16.0 | | | | | | | | 30.0 | 238.0 | * | | | |
| | *Perna viridis* | | | | | | | | | | | | | | | |
| | *Pinctada martensii* | | | | | | | | | | | | | | | |
| | *Placuna placenta* | | | | | | | | | | 2.0 | 41.2 | | | | |
| | *Ruditapes philippinarum* | | | | 2.0 | 9.3 | | | | | | | | | | |
| | *Scapharca subcrenata* | | | | 1.0 | 5.3 | | | | | | | | 4.0 | 280.0 | |
| | *Spondylus sp.* | | | | | | | | | | | | | | | |
| | *Tegillarca nodifera* | | | | | | | | | | | | | | | |
| | *Vepricardium asiaticum* | | | | | | | | | | | | | | | |
| | *Tolonostrea talonata* | | | | | | | | | | | | | | | |
| Cephalopod | *Lolingid squid* | | | | | | | | | | | | | | | |
| | *Sepiid cuttlefish* | | | | 1.0 | 108.0 | | | | | | | | | | |
| | *Octopus sp.* | | | | | | | | | | | | | | | |
| Echinoderm | *Acaudina molpadioides* | | | | 2.0 | 29.0 | | | | | | | | 3.0 | 82.0 | |
| | *Asteroidea* | | | | | | | | | | | | | | | |
| | Brittle star | | | | | | | | | | | | | | | |
| | *Pseudocolochirus sp.* | | | | | | | | | | | | | | | |

Demersal Trawling Data in August 2015

| | | TSA | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | TSA-1 | | | TSA-2 | | | TSA-3 | | | TSA-4 | | | TSA-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| | *Temnopleurus toreumaticus* | 2800.0 | 28500.0 | * | 2300.0 | 23200.0 | * | 2100.0 | 21500.0 | * | 2500.0 | 26700.0 | * | 2600.0 | 27800.0 | * |
| Cnidaria | *Edwardsia japonica (c.f.)* | | | | | | | | | | | | | | | |
| | *Ellisella laevis* | | | | | | | | | | 1.0 | 21.0 | | | | |
| | *Gorgonia spp.* | | | | | | | | | | | | | | | |
| | *Carcinactis ichikawai* | 3.0 | 0.7 | | 11.0 | 4.9 | | 2.0 | 0.4 | | 2.0 | 0.5 | | 1.0 | 0.2 | |
| | *Cavernularia habereri* | | | | 5.0 | 21.0 | | 11.0 | 43.0 | | | | | | | |
| | Jelly fish | | | | | | | | | | | | | | | |
| | *Pteroeides chinense* | | | | 6.0 | 94.0 | | 5.0 | 81.0 | | | | | 4.0 | 72.0 | |
| | Sea anomea | | | | | | | | | | | | | 1.0 | 15.2 | |
| | *Virgularia gustaviana* | | | | 1.0 | 9.2 | | | | | | | | | | |
| Urochordata | Sea Squirts | | | | | | | | | | | | | | | |
| Cirripedia | *Bananus spp.(c.f.)* | 80.0 | 43.0 | * | 70.0 | 36.0 | * | 60.0 | 34.0 | * | 42.0 | 40.8 | * | 80.0 | 44.7 | * |

Demersal Trawling Data in August 2015

| | | TSB | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | TSB-1 | | | TSB-2 | | | TSB-3 | | | TSB-4 | | | TSB-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| Fish | *Acentrogobius caninus* | | | | | | | 1.0 | 4.0 | | | | | | | |
| | *Acanthemblemaria rivasi* | | | | | | | | | | | | | | | |
| | *Apogon kiensis* | | | | | | | 1.0 | 14.0 | | | | | | | |
| | *Arius sinensis* | 2.0 | 188.0 | | | | | | | | 1.0 | 40.0 | | | | |
| | *Arnoglossus waitei* | | | | | | | | | | | | | | | |
| | *Caranx kalla* | | | | 1.0 | 10.0 | | 1.0 | 12.0 | | | | | 1.0 | 10.0 | |
| | *Chrysochir aureus* | | | | | | | | | | | | | | | |
| | *Collichthys lucidus* | 12.0 | 150.0 | | 11.0 | 164.0 | | 4.0 | 44.0 | | 2.0 | 16.0 | | 4.0 | 50.0 | |
| | *Cynoglossus abbreviatus* | | | | | | | | | | 1.0 | 12.0 | | | | |
| | *Cynoglossus arel* | 1.0 | 18.0 | | 11.0 | 358.0 | | 3.0 | 76.0 | | 2.0 | 108.0 | | 2.0 | 64.0 | |
| | *Cynoglossus puncticeps* | | | | | | | | | | | | | | | |
| | *Cynoglossus roulei* | | | | | | | | | | | | | | | |
| | *Cynoglossus semilaevis* | | | | | | | | | | | | | | | |
| | *Dasyatis akajei* | | | | | | | | | | | | | | | |
| | *Dasyatis zugei* | | | | 1.0 | 218.0 | | 2.0 | 788.0 | | | | | | | |
| | *Dendrophysa russelii* | 1.0 | 36.0 | | 2.0 | 46.0 | | 4.0 | 68.0 | | | | | 3.0 | 56.0 | |
| | *Drepane punctata* | | | | | | | | | | | | | | | |
| | *Epinephelus awoara* | | | | | | | | | | | | | | | |
| | *Epinephelus bruneus* | | | | | | | | | | | | | | | |
| | *Gerres filamentosus* | 1.0 | 12.0 | | | | | 1.0 | 10.0 | | | | | | | |
| | *Grammoplites scaber* | | | | | | | | | | 1.0 | 12.0 | | | | |
| | *Gymnothorax chilospilus* | | | | | | | | | | | | | | | |
| | *Gymnura japonica* | 1.0 | 2500.0 | | | | | | | | | | | | | |
| | *Ilisa elongata* | | | | | | | | | | | | | | | |
| | *Inegocia japonica* | | | | | | | | | | | | | | | |
| | *Inegocia japonica* | | | | | | | | | | | | | | | |
| | *Inimicus didactylus* | | | | | | | | | | | | | | | |
| | *Inimicus japonicus* | | | | | | | | | | | | | | | |
| | *Johnius belangerii* | | | | 12.0 | 362.0 | | 6.0 | 184.0 | | 3.0 | 78.0 | | | | |
| | *Johnius heterolepis* | 3.0 | 184.0 | | | | | | | | | | | | | |
| | *Konosirus punctatus* | | | | | | | | | | | | | 1.0 | 44.0 | |
| | *Lagocephalus wheeleri* | | | | | | | | | | | | | | | |
| | *Leiognathus brevirostris* | 15.0 | 205.0 | | 10.0 | 140.0 | | 18.0 | 234.0 | | 39.0 | 498.0 | | 15.0 | 161.0 | |
| | *Leiognathus nuchalis* | 182.0 | 2357.0 | | 164.0 | 2426.0 | | 60.0 | 862.0 | | 3.0 | 22.0 | | 23.0 | 279.0 | |
| | *Lepturacanthus savala* | | | | | | | | | | | | | | | |
| | *Liza macrolepis* | 4.0 | 32.0 | | 1.0 | 18.0 | | 1.0 | 16.0 | | 6.0 | 368.0 | | 12.0 | 718.0 | |
| | *Minous monodactylus* | | | | | | | | | | | | | | | |
| | *Monacanthus chinensis* | | | | | | | | | | | | | | | |
| | *Muraenesox cinereus* | | | | | | | 1.0 | 32.0 | | | | | | | |
| | *Nemipterus japonicus* | 1.0 | 12.0 | | | | | | | | | | | | | |
| | *Oxyurichthys tentacularis* | | | | | | | | | | | | | | | |
| | *Paracentropogon rubripinnis* | | | | | | | | | | | | | | | |
| | *Parachaeturichthys polynema* | | | | | | | | | | | | | | | |
| | *Paramonacanthus sulcatus* | | | | 4.0 | 15.3 | | 1.0 | 6.0 | | | | | 2.0 | 16.0 | |
| | *Parargyrops edita* | | | | | | | | | | | | | | | |
| | *Pennahia anea* | 3.0 | 30.0 | | 2.0 | 44.0 | | | | | 1.0 | 8.0 | | | | |
| | *Pennahia macrocephalus* | | | | | | | 1.0 | 4.0 | | | | | | | |
| | *Platycephalus indicus* | 4.0 | 162.0 | | 3.0 | 68.0 | | 4.0 | 178.0 | | 1.0 | 26.0 | | 1.0 | 208.0 | |
| | *Plectorhynchus pictus* | | | | | | | | | | | | | | | |

Demersal Trawling Data in August 2015

| | | TSB | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | TSB-1 | | | TSB-2 | | | TSB-3 | | | TSB-4 | | | TSB-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| | *Polydactylus sextarius* | 10.0 | 184.0 | | 7.0 | 128.0 | | 1.0 | 18.0 | | 1.0 | 18.0 | | 1.0 | 12.0 | |
| | *Pseudorhombus cinnamoneus* | | | | | | | 1.0 | 24.0 | | 1.0 | 22.0 | | | | |
| | *Rhabdosargus sarba* | | | | | | | | | | | | | | | |
| | *Saurida elongata* | | | | 1.0 | 110.0 | | 1.0 | 90.0 | | 1.0 | 64.0 | | | | |
| | *Scatophagus argus* | | | | | | | | | | | | | | | |
| | *Sebastiscus marmoratus* | | | | | | | | | | | | | | | |
| | *Siganus fuscescens* | | | | | | | | | | | | | | | |
| | *Sillago sihama* | | | | | | | 1.0 | 28.0 | | 1.0 | 38.0 | | | | |
| | *Solea ovata* | 3.0 | 34.0 | | 1.0 | 12.0 | | 2.0 | 18.0 | | | | | 1.0 | 10.0 | |
| | *Stolephorus insularis* | | | | | | | | | | | | | | | |
| | *Takifugu niphobles* | | | | | | | | | | | | | | | |
| | *Takifugu oblongus* | | | | | | | | | | | | | | | |
| | *Takifugu xanthopterus* | | | | | | | | | | | | | | | |
| | *Terapon jarbua* | | | | | | | | | | | | | | | |
| | *Terapon theraps* | | | | | | | 1.0 | 18.0 | | | | | | | |
| | *Thryssa hamiltonii* | | | | | | | | | | | | | | | |
| | *Thryssa kammalensis* | | | | | | | | | | | | | | | |
| | *Trachycephalus uranoscopa* | 1.0 | 12.0 | | 1.0 | 12.0 | | 1.0 | 16.0 | | 2.0 | 16.0 | | | | |
| | *Trypauchen vagina* | | | | | | | | | | | | | | | |
| | *Uroconger lepturus* | | | | | | | | | | | | | | | |
| | *Zebrias zebra* | | | | | | | | | | | | | | | |
| Prawn or shrimp | *Alpheus japonicus* | | | | | | | | | | | | | | | |
| | *Metapenaeopsis barbata* | | | | | | | | | | | | | | | |
| | *Metapenaeus ensis* | 2.0 | 28.0 | | 4.0 | 38.0 | | 3.0 | 18.0 | | | | | | | |
| | *Metapenaeus joyneri* | 11.0 | 82.0 | | 8.0 | 38.0 | | 4.0 | 18.0 | | 5.0 | 24.0 | | 4.0 | 28.0 | |
| | *Parapenaeopsis hardwickii* | 4.0 | 46.0 | | 2.0 | 20.0 | | 7.0 | 58.0 | | 2.0 | 14.0 | | 1.0 | 12.0 | |
| | *Parapenaeopsis tenella* | 3.0 | 12.0 | | 2.0 | 8.0 | | 5.0 | 20.0 | | | | | 8.0 | 26.0 | |
| | *Penaeus japonicus* | | | | | | | | | | | | | | | |
| | *Penaeus merguiensis* | | | | | | | | | | 5.0 | 90.0 | | 2.0 | 36.0 | |
| | *Penaeus penicillatus* | 5.0 | 78.0 | | 5.0 | 68.0 | | 5.0 | 72.0 | | | | | | | |
| | *Solenocera crassicornis* | | | | | | | 1.0 | 4.0 | | | | | 2.0 | 32.0 | |
| | *Trachypenaeus curvirostris* | | | | | | | | | | | | | | | |
| Crab | *Arcania heptacontra* | | | | | | | | | | | | | | | |
| | *Arcania undecimspinosa* | | | | | | | 1.0 | 8.6 | | | | | 2.0 | 48.2 | |
| | *Calappa philargius* | 1.0 | 32.5 | | 4.0 | 328.5 | | | | | | | | | | |
| | *Charybdis affinis* | 17.0 | 186.0 | | 18.0 | 176.0 | | 14.0 | 156.0 | | 9.0 | 106.0 | | 8.0 | 80.0 | |
| | *Charybdis cruciata* | | | | | | | | | | | | | | | |
| | *Charybdis japonica* | | | | | | | | | | | | | | | |
| | *Charybdis spp.* | 32.0 | 191.0 | * | 35.0 | 237.0 | * | 38.0 | 205.0 | * | 35.0 | 189.0 | * | 33.0 | 187.0 | * |
| | *Dorippe polita* | 25.0 | 177.0 | * | 30.0 | 185.0 | * | 20.0 | 163.0 | * | 27.0 | 174.0 | * | 30.0 | 188.0 | * |
| | *Eucrate crenata* | | | | | | | | | | | | | | | |
| | *Galene bispinosa* | | | | 3.0 | 169.0 | | | | | 3.0 | 116.0 | | 8.0 | 764.0 | |
| | Hermit crab | 45.0 | 190.0 | * | 42.0 | 780.0 | * | 30.0 | 130.0 | * | 46.0 | 190.0 | * | 48.0 | 210.0 | * |
| | *Leucosia craniolaris* | 11.0 | 35.0 | | 20.0 | 66.0 | * | 7.0 | 23.0 | | 5.0 | 23.0 | | 9.0 | 28.0 | |
| | *Macrophthalmus japonicus* | | | | | | | | | | | | | | | |
| | *Parapanope euagora* | | | | | | | | | | | | | | | |
| | *Parthenope validus* | 21.0 | 179.0 | * | 18.0 | 171.0 | * | 28.0 | 291.0 | * | 20.0 | 178.0 | * | 25.0 | 182.0 | * |
| | *Philyra sp.* | | | | | | | | | | | | | | | |
| | *Porcellanella picta* | | | | | | | | | | | | | | | |

Demersal Trawling Data in August 2015

| | | TSB | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | TSB-1 | | | TSB-2 | | | TSB-3 | | | TSB-4 | | | TSB-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| | *Portunus pelagicus* | | | | | | | | | | | | | | | |
| | *Portunus sanguinolentus* | | | | | | | | | | | | | | | |
| | *Portunus trituberculatus* | 3.0 | 272.0 | | 1.0 | 78.0 | | 2.0 | 308.0 | | | | | 2.0 | 282.0 | |
| | *Scylla serrata (mud crab)* | | | | | | | | | | | | | | | |
| | *Tachypleus tridentatus* | | | | | | | | | | | | | | | |
| | *Typhlocarcinus sp.* | | | | | | | | | | 1.0 | 0.9 | | | | |
| Mantis shrimp | *Harpiosquilla harpax* | 3.0 | 122.0 | | 3.0 | 140.0 | | 4.0 | 172.0 | | | | | 3.0 | 108.0 | |
| | *Erugosquilla woodmasoni* | | | | | | | | | | | | | | | |
| | *Oratosquilla kempi* | | | | | | | | | | | | | | | |
| | *Oratosquilla interrupta* | 4.0 | 68.0 | | 1.0 | 12.0 | | | | | 4.0 | 46.0 | | 3.0 | 34.0 | |
| | *Oratosquilla oratoria* | | | | 1.0 | 8.0 | | 1.0 | 4.0 | | 1.0 | 10.0 | | | | |
| Gastropod | *Apollon olivator rubustus* | | | | | | | | | | | | | | | |
| | *Bursa rana* | | | | | | | | | | | | | | | |
| | *Calyptraea sp.* | | | | 1.0 | 0.3 | | 5.0 | 2.2 | | | | | 5.0 | 2.1 | |
| | *Drupa musiva* | | | | | | | | | | 1.0 | 4.2 | | | | |
| | *Epitonium scalare* | | | | 5.0 | 49.0 | | 3.0 | 26.0 | | | | | | | |
| | *Inquistor flavidula* | | | | | | | | | | | | | | | |
| | *Lophiotoma leucotropis* | | | | | | | | | | | | | | | |
| | *Murex trapa* | 30.0 | 292.0 | * | 13.0 | 112.0 | | 15.0 | 119.0 | | 20.0 | 186.0 | * | 30.0 | 290.0 | * |
| | *Nassarius hepaticus* | 1.0 | 3.3 | | | | | | | | | | | 2.0 | 6.2 | |
| | *Nassarius succinctus* | | | | | | | 30.0 | 64.0 | * | | | | | | |
| | *Natica tigrina* | | | | | | | | | | | | | | | |
| | *Philine sp.* | 20.0 | 180.0 | * | 5.0 | 26.1 | | 8.0 | 64.0 | | 30.0 | 270.0 | * | 26.0 | 196.0 | * |
| | *Pilosabia pilosa* | | | | 2.0 | 1.6 | | | | | | | | 1.0 | 0.3 | |
| | *Siphopatella walshi* | | | | | | | 5.0 | 2.2 | | | | | | | |
| | *Tonna sulcosa* | | | | | | | | | | | | | 2.0 | 148.0 | |
| | *Turritella terebra* | 1300.0 | 12820.0 | * | 1380.0 | 13700.0 | * | 700.0 | 6800.0 | * | 1500.0 | 14300.0 | * | 1200.0 | 11600.0 | * |
| Bivalve | Bivalve Juvenil | 30.0 | 16.0 | * | 25.0 | 12.5 | * | | | | | | | | | |
| | *Chione isabellina* | | | | | | | | | | | | | | | |
| | *Chlamys nobilis* | | | | 2.0 | 12.4 | | 1.0 | 4.7 | | 1.0 | 4.9 | | | | |
| | *Macoma candida* | 20.0 | 214.0 | * | | | | | | | | | | | | |
| | *Ostrea sp.* | | | | 6.0 | 5.2 | | 30.0 | 26.0 | * | | | | | | |
| | *Paphia undulata* | | | | | | | 4.0 | 19.2 | | 30.0 | 260.0 | * | | | |
| | *Perna viridis* | | | | 6.0 | 46.0 | | | | | | | | | | |
| | *Pinctada martensii* | | | | 1.0 | 51.4 | | | | | | | | | | |
| | *Placuna placenta* | | | | | | | | | | | | | 1.0 | 19.4 | |
| | *Ruditapes philippinarum* | | | | | | | | | | | | | | | |
| | *Scapharca subcrenata* | 1.0 | 21.6 | | 1.0 | 147.0 | | | | | | | | | | |
| | *Spondylus sp.* | | | | | | | | | | | | | | | |
| | *Tegillarca nodifera* | | | | | | | | | | | | | | | |
| | *Vepricardium asiaticum* | | | | | | | | | | | | | | | |
| | *Tolonostrea talonata* | | | | | | | | | | | | | | | |
| Cephalopod | *Lolingid squid* | | | | | | | | | | | | | | | |
| | *Sepiid cuttlefish* | | | | | | | | | | 1.0 | 16.0 | | | | |
| | *Octopus sp.* | | | | | | | | | | | | | | | |
| Echinoderm | *Acaudina molpadioides* | 1.0 | 8.4 | | | | | 5.0 | 109.0 | | | | | 3.0 | 170.0 | |
| | *Asteroidea* | | | | | | | | | | | | | | | |
| | Brittle star | 1.0 | 2.2 | | | | | | | | | | | | | |
| | *Pseudocolochirus sp.* | | | | | | | | | | | | | | | |

Demersal Trawling Data in August 2015

| | | TSB | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | TSB-1 | | | TSB-2 | | | TSB-3 | | | TSB-4 | | | TSB-5 | | |
| Group | *Species Name* | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling | Total | Total | Subsampling |
| | | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) | number | weight(g) | (*) |
| | *Temnopleurus toreumaticus* | 15.0 | 135.0 | | 20.0 | 202.0 | | 126.0 | 1258.0 | * | 25.0 | 280.0 | | 42.0 | 438.0 | * |
| Cnidaria | *Edwardsia japonica (c.f.)* | 1.0 | 2.4 | | | | | | | | | | | | | |
| | *Ellisella laevis* | | | | | | | | | | | | | | | |
| | *Gorgonia spp.* | 1.0 | 6.4 | | | | | | | | | | | | | |
| | *Carcinactis ichikawai* | 2.0 | 0.7 | | | | | 2.0 | 0.8 | | 4.0 | 2.1 | | | | |
| | *Cavernularia habereri* | 20.0 | 110.0 | * | | | | 7.0 | 29.0 | | | | | | | |
| | Jelly fish | | | | | | | | | | | | | | | |
| | *Pteroeides chinense* | | | | 2.0 | 12.8 | | | | | | | | | | |
| | Sea anomea | | | | 1.0 | 15.4 | | | | | | | | | | |
| | *Virgularia gustaviana* | | | | 2.0 | 16.2 | | | | | | | | | | |
| Urochordata | Sea Squirts | | | | | | | | | | | | | | | |
| Cirripedia | *Bananus spp.(c.f.)* | 120.0 | 70.0 | * | 90.0 | 46.0 | * | 80.0 | 40.1 | * | 70.0 | 38.0 | * | 60.0 | 38.0 | * |